

# Energy-Momentum Tensor on the Lattice via Gradient Flow

Masakiyo Kitazawa  
(Osaka University)

FlowQCD Collab.


M. Asakawa, T. Hatsuda, **T. Iritani**, H. Suzuki, **R. Yanagihara**  
PRD**94**,114512(2016); PRD**96**,111502(2017); arXiv:1803.05656

WHOT-QCD Collab.

S. Ejiri, K. Kanaya, H. Suzuki, **Y. Taniguchi**, T. Umeda, ...  
PRD**96**,014509(2017); arXiv:1710.10015; arXiv:1711.02262

**Takaura, Suzuki**, Iritani, MK, in preparation

MK, Mogliacci, et al., in preparation



# Energy-Momentum Tensor

One of the **most fundamental** quantities in physics

$T_{\mu\nu}$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}$$
$$\partial_{\mu} T^{\mu\nu} = 0$$

# Energy-Momentum Tensor

$$T_{\mu\nu} = \begin{bmatrix} \text{energy} & & & \\ T_{00} & T_{01} & T_{02} & T_{03} \\ T_{10} & T_{11} & T_{12} & T_{13} \\ T_{20} & T_{21} & T_{22} & T_{23} \\ T_{30} & T_{31} & T_{32} & T_{33} \\ & & \text{stress} & \end{bmatrix}$$


The diagram illustrates the components of the Energy-Momentum Tensor  $T_{\mu\nu}$ . The tensor is represented as a 4x4 matrix. The components are categorized as follows:

- $T_{00}$  is labeled "energy".
- The components  $T_{01}, T_{02}, T_{03}$  are labeled "momentum".
- The components  $T_{11}, T_{22}, T_{33}$  are labeled "pressure".
- The components  $T_{31}, T_{32}, T_{33}$  are labeled "stress".

All components are important physical observables!


$T_{\mu\nu}$  : nontrivial observable  
on the lattice

- ① Definition of the operator is nontrivial because of the explicit breaking of Lorentz symmetry



ex:  $T_{\mu\nu} = F_{\mu\rho}F_{\nu\rho} - \frac{1}{4}\delta_{\mu\nu}FF$

$F_{\mu\nu} =$




- ② Its measurement is extremely noisy due to high dimensionality and etc.

# Thermodynamics

direct measurement of  
expectation values

$$\langle T_{00} \rangle, \langle T_{ii} \rangle$$



# Fluctuations and Correlations


viscosity, specific heat, ...

$$c_V \sim \langle \delta T_{00}^2 \rangle$$

$$\eta = \langle T_{12}; T_{12} \rangle$$


If we have

$$T_{\mu\nu}$$



- flux tube / hadrons
- stress distribution

## Hadron Structure



- vacuum configuration
- mixed state on 1<sup>st</sup> transition

## Vacuum Structure

# Contents




## Constructing EMT on the lattice

### Thermodynamics

direct measurement of  
expectation values

$$\langle T_{00} \rangle, \langle T_{ii} \rangle$$



## Thermodynamics

### Fluctuations and Correlations


viscosity, specific heat, ...

$$\eta = \int_0^\infty dt \langle T_{12}; T_{12} \rangle$$
$$c_V \sim \langle \delta T_{00}^2 \rangle$$

## EMT Correlation Function

### Hadron Structure

- flux tube / hadrons
- stress distribution



## Stress distribution in $\bar{q}q$ system

# Yang-Mills Gradient Flow

Luscher 2010

Narayanan, Neuberger, 2006

Luscher, Weiss, 2011

$$\frac{\partial}{\partial t} A_\mu(t, x) = - \frac{\partial S_{\text{YM}}}{\partial A_\mu}$$


$$A_\mu(0, x) = A_\mu(x)$$

t: "flow time"  
dim:[length<sup>2</sup>]

leading

$$\partial_t A_\mu = D_\nu G_{\mu\nu} = \partial_\nu \partial_\nu A_\mu + \dots$$

- diffusion equation in 4-dim space
- diffusion distance  $d \sim \sqrt{8t}$
- "continuous" cooling/smearing
- No UV divergence at  $t > 0$



# Yang-Mills Gradient Flow

Luscher 2010

Narayanan, Neuberger, 2006

Luscher, Weiss, 2011

$$\frac{\partial}{\partial t} A_\mu(t, x) = - \frac{\partial S_{\text{YM}}}{\partial A_\mu}$$

$$A_\mu(0, x) = A_\mu(x)$$

t: "flow time"  
dim:[length<sup>2</sup>]



leading

$$\partial_t A_\mu = D_\nu G_{\mu\nu} = \partial_\nu \partial_\nu A_\mu + \dots$$

## Applications

scale setting / topological charge / running coupling  
noise reduction / **defining operators** / ...




# Small Flow-Time Expansion

Luescher, Weisz, 2011  
Suzuki, 2013

$$\tilde{\mathcal{O}}(t, x) \xrightarrow[t \rightarrow 0]{} \sum_i c_i(t) \mathcal{O}_i^R(x)$$

an operator at  $t > 0$


remormalized operators  
of original theory



# Constructing EMT 1

Suzuki, 2013

$$\tilde{\mathcal{O}}(t, x) \xrightarrow{t \rightarrow 0} \sum_i c_i(t) \mathcal{O}_i^R(x)$$



## □ Gauge-invariant dimension 4 operators


$$\left\{ \begin{array}{l} U_{\mu\nu}(t, x) = G_{\mu\rho}(t, x)G_{\nu\rho}(t, x) - \frac{1}{4}\delta_{\mu\nu}G_{\mu\nu}(t, x)G_{\mu\nu}(t, x) \\ E(t, x) = \frac{1}{4}\delta_{\mu\nu}G_{\mu\nu}(t, x)G_{\mu\nu}(t, x) \end{array} \right.$$

# Constructing EMT 2

Suzuki, 2013

$$U_{\mu\nu}(t, x) = \alpha_U(t) \left[ T_{\mu\nu}^R(x) - \frac{1}{4} \delta_{\mu\nu} T_{\rho\rho}^R(x) \right] + \mathcal{O}(t)$$

$$E(t, x) = \langle E(t, x) \rangle + \alpha_E(t) T_{\rho\rho}^R(x) + \mathcal{O}(t)$$



Suzuki coeffs.  $\left\{ \begin{array}{l} \alpha_U(t) = g^2 [1 + 2b_0 s_1 g^2 + O(g^4)] \\ \alpha_E(t) = \frac{1}{2b_0} [1 + 2b_0 s_2 g^2 + O(g^4)] \end{array} \right.$

$$g = g(1/\sqrt{8t})$$

$$s_1 = 0.03296 \dots$$

$$s_2 = 0.19783 \dots$$

## Remormalized EMT

$$T_{\mu\nu}^R(x) = \lim_{t \rightarrow 0} \left[ \frac{1}{\alpha_U(t)} U_{\mu\nu}(t, x) + \frac{\delta_{\mu\nu}}{4\alpha_E(t)} E(t, x)_{\text{subt.}} \right]$$

# Gradient Flow for Fermions

$$\partial_t \psi(t, x) = D_\mu D_\mu \psi(t, x)$$

$$\partial_t \bar{\psi}(t, x) = \bar{\psi}(t, x) \overleftarrow{D}_\mu \overleftarrow{D}_\mu$$

$$D_\mu = \partial_\mu + A_\mu(t, x)$$

Luscher, 2013

Makino, Suzuki, 2014

Taniguchi+ (WHOT)

2016; 2017

□ Not “gradient” flow but a “diffusion” equation.

□ Divergence in field renormalization of fermions.

□ All observables are finite at  $t > 0$  once  $Z(t)$  is fixed.

$$\tilde{\psi}(t, x) = Z(t)\psi(t, x)$$

□ Energy-momentum tensor from SFTE Makino, Suzuki, 2014

# Higher Order Analysis

The 2-loop analysis of  $\alpha_U, \alpha_E$  is now available!  
(full QCD / pure gauge)

Harlander+, 1808.09837

	LO	1-loop	2-loop	3-loop
$e - 3p$	× zero	○	○	
$e + p$	○	○		

Suzuki (2013)

Effect on thermodynamics  
(pure gauge)

- $e-3p$ : negligible (<0.5%)
- $e+p$ : 2-3% increase

# Higher Order Analysis

The 2-loop analysis of  $\alpha_U, \alpha_E$  is now available!  
(full QCD / pure gauge)

Harlander+, 1808.09837

	LO	1-loop	2-loop	3-loop
$e - 3p$	× zero	○	○	
$e + p$	○	○	○	

Suzuki (2013) Harlander+(2018)

Effect on thermodynamics  
(pure gauge)

- $e-3p$ : negligible ( $<0.5\%$ )
- $e+p$ : 2-3% increase

# Higher Order Analysis

The 2-loop analysis of  $\alpha_U, \alpha_E$  is now available!  
(full QCD / pure gauge)

Harlander+, 1808.09837

	LO	1-loop	2-loop	3-loop
$e - 3p$	× zero	○	○	○
$e + p$	○	○	○	○

Takaura, Suzuki,  
Iritani, MK, in prep.

Suzuki (2013) Harlander+(2018)

Effect on thermodynamics  
(pure gauge)

- $e-3p$ : negligible (<0.5%)
- $e+p$ : 2-3% increase

# Contents




## Constructing EMT on the lattice

### Thermodynamics

direct measurement of  
expectation values

$$\langle T_{00} \rangle, \langle T_{ii} \rangle$$



## Thermodynamics

### Fluctuations and Correlations


viscosity, specific heat, ...

$$\eta = \int_0^\infty dt \langle T_{12}; T_{12} \rangle$$
$$c_V \sim \langle \delta T_{00}^2 \rangle$$

## EMT Correlation Function

### Hadron Structure

- flux tube / hadrons
- stress distribution





## Stress distribution in $\bar{q}q$ system



# t, a Dependence

- Budapest-Wuppertal
- ..... Bielefeld
- $\beta = 6.719, 64^3 \times 12$
- $\beta = 6.941, 96^3 \times 16$
- $\beta = 7.117, 128^3 \times 20$



- $\left\{ \begin{array}{l} \sqrt{8t} < a : \text{strong discretization effect} \\ \sqrt{8t} > 1/(2T) : \text{over smeared} \end{array} \right.$


$a < \sqrt{8t} < 1/(2T) : \text{Linear } t \text{ dependence}$

# Double Extrapolation

$$t \rightarrow 0, a \rightarrow 0$$

$$\langle T_{\mu\nu}(t) \rangle_{\text{latt}} = \langle T_{\mu\nu}(t) \rangle_{\text{phys}} + C_{\mu\nu} t + D_{\mu\nu}(t) \frac{a^2}{t}$$

$O(t)$  terms in SFTE    lattice discretization




Continuum extrapolation

$$\langle T_{\mu\nu}(t) \rangle_{\text{cont}} = \langle T_{\mu\nu}(t) \rangle_{\text{lat}} + C(t)a^2$$

Small t extrapolation


$$\langle T_{\mu\nu} \rangle = \langle T_{\mu\nu}(t) \rangle + C' t$$

# Double Extrapolation



Black line: continuum extrapolated

# Double Extrapolation



Black line: continuum extrapolated

□ Fitting ranges:

□ range-1:  $0.01 < tT^2 < 0.015$



□ range-2:  $0.005 < tT^2 < 0.015$

□ range-3:  $0.01 < tT^2 < 0.02$



Systematic error from the choice of fitting range  
 $\approx$  statistical error

# Temperature Dependence



Error includes

- statistical error
- choice of  $t$  range for  $t \rightarrow 0$  limit
- uncertainty in  $a\Lambda_{\text{MS}}$

total error <1.5% for  $T > 1.1T_c$

- Excellent agreement with integral method
- High accuracy only with  $\sim 2000$  confs.

# Thermodynamics of SU(3) YM

## □ Integral method

- Most conventional / established
- Use thermodynamic relations  
Boyd+ 1995; Borsanyi, 2012

$$p = \frac{T}{V} \ln Z$$

$$T \frac{\partial(p/T^4)}{\partial T} = \frac{\varepsilon - 3p}{T^4}$$

## □ Gradient-flow method

- Take expectation values of EMT  
FlowQCD, 2014, 2016

$$\begin{cases} \varepsilon = \langle T_{00} \rangle \\ p = \langle T_{11} \rangle \end{cases}$$

## □ Moving-frame method

Giusti, Pepe, 2014~

## □ Non-equilibrium method

- Use Jarzynski's equality Caselle+, 2016;2018

## □ Differential method

Shirogane+(WHOT-QCD), 2016~

# SU(3) YM EoS: Comparison

$$\frac{e - 3p}{T^4}$$






Figure from  
Caselle+,  
1801.03110

- Measurement of thermodynamics with various methods.
- All results are in good agreement.
- But, non-negligible discrepancy at  $T/T_c \approx 1-1.3$ ?

# Effect of Higher-Order Coeffs.

Takaura, Suzuki, Iritani, MK, in prep.



Effect on thermodynamics  
(pure gauge)


- e-3p: negligible (<0.5%)
- e+p: 2-3% increase



# 2+1 QCD EoS from Gradient Flow

Taniguchi+ (WHOT-QCD), PRD**96**, 014509 (2017)

$m_{PS}/m_V \approx 0.63$




- Agreement with integral method except for  $N_t=4, 6$
- No stable extrapolation for  $N_t=4, 6$
- Statistical error is substantially suppressed!

Physical mass: Kanaya+ (WHOT-QCD), 1710.10015

# Pressure anisotropy in finite system

## Casimir effect



## Finite system at nonzero $T$

MK, Mogliacci, et al.  
in preparation

$$V = L_x \times L_y \times L_z$$

$$L_x \ll L_y = L_z$$



pressure anisotropy

$$T_{11} \neq T_{22} = T_{33}$$



# Pressure Anisotropy

MK, Mogliacci, et al. in prep.

## Free scalar field


□  $L_2=L_3=\infty$

Mogliacci+, 1807.07871



# Pressure Anisotropy

MK, Mogliacci, et al. in prep.



## Free scalar field

$$\square L_2 = L_3 = \infty$$

Mogliacci+, 1807.07871

## Lattice results

$\square$  Periodic BC

$\square N_s^2 \times N_x \times N_t = 72^2 \times N_x \times 12$

$\square N_x = 12, 14, 16, 18$

$\square$  Only  $t \rightarrow 0$  limit (fixed  $a$ )

Medium near  $T_c$  is remarkably insensitive to finite size!  
How do we understand??

# Contents




## Constructing EMT on the lattice

### Thermodynamics

direct measurement of  
expectation values

$$\langle T_{00} \rangle, \langle T_{ii} \rangle$$



## Thermodynamics

### Fluctuations and Correlations


viscosity, specific heat, ...

$$\eta = \int_0^\infty dt \langle T_{12}; T_{12} \rangle$$
$$c_V \sim \langle \delta T_{00}^2 \rangle$$

## EMT Correlation Function

### Hadron Structure

- flux tube / hadrons
- stress distribution



## Stress distribution in $\bar{q}q$ system

# EMT Correlator: Motivation

## □ Transport Coefficient

Kubo formula  $\rightarrow$  viscosity

$$\eta = \int_0^\infty dt \int_0^{1/T} d\tau \int d^3x \langle T_{12}(x, -i\tau) T_{12}(0, t) \rangle$$

Karsch, Wyld, 1987  
Nakamura, Sakai, 2005  
Meyer; 2007, 2008  
...  
Borsanyi+, 2018  
Astrakhantsev+, 2018

## □ Energy/Momentum Conservation





$\langle \bar{T}_{0\mu}(\tau) \bar{T}_{\rho\sigma}(0) \rangle$  :  $\tau$ -independent constant

## □ Fluctuation-Response Relations

$$c_V = \frac{\langle \delta E^2 \rangle}{VT^2} \quad E + p = \frac{\langle \bar{T}_{01}^2 \rangle}{VT} = \frac{\langle \bar{T}_{11} \bar{T}_{00} \rangle}{VT}$$

# EMT Euclidean Correlator




FlowQCD, PR D96, 111502 (2017)

	$tT^2 = 0.0024$
	$tT^2 = 0.0035$
	$tT^2 = 0.0052$
	$tT^2 = 0.0069$

$$\langle \bar{T}_{44}(\tau) \bar{T}_{44}(0) \rangle$$

$$\langle \bar{T}_{44}(\tau) \bar{T}_{11}(0) \rangle$$

$$\langle \bar{T}_{41}(\tau) \bar{T}_{41}(0) \rangle$$




- $\tau$ -independent plateau in all channels  $\rightarrow$  conservation law
- Confirmation of fluctuation-response relations
- New method to measure  $c_v$

- Similar result for (41;41) channel: Borsanyi+, 2018
- Perturbative analysis: Eller, Moore, 2018


# Fluctuation-Response Relations

$$\langle T_{44}(\tau)T_{44}(0) \rangle$$




New measurement of  $c_v$

$$\langle T_{44}(\tau)T_{11}(0) \rangle$$



$$\langle T_{41}(\tau)T_{41}(0) \rangle$$



Confirmation of FRR

$$E + p = \frac{\langle \bar{T}_{01}^2 \rangle}{VT} = \frac{\langle \bar{T}_{11} \bar{T}_{00} \rangle}{VT}$$

2+1 QCD:

Taniguchi+ (WHOT-QCD), 1711.02262

$c_v/T^3$

$T/T_c$	$C_{44;44}(\tau_m)$	Ref.[19]	Ref.[11]	ideal gas
1.68	17.7(8) $^{(+2.1)}_{(-0.4)}$	22.8(7)*	17.7	21.06
2.24	17.5(0.8) $^{(+0)}_{(-0.1)}$	17.9(7)**	18.2	21.06



# Contents




## Constructing EMT on the lattice

### Thermodynamics

direct measurement of  
expectation values

$$\langle T_{00} \rangle, \langle T_{ii} \rangle$$



## Thermodynamics

### Fluctuations and Correlations


viscosity, specific heat, ...

$$\eta = \int_0^\infty dt \langle T_{12}; T_{12} \rangle$$
$$c_V \sim \langle \delta T_{00}^2 \rangle$$

## EMT Correlation Function

### Hadron Structure

- flux tube / hadrons
- stress distribution




## Stress distribution in $\bar{q}q$ system

Stress = Force per Unit Area

# Stress = Force per Unit Area


Pressure



$$\vec{P} = P\vec{n}$$

# Stress = Force per Unit Area

Pressure





$$\vec{P} = P\vec{n}$$

In thermal medium

$$T_{ij} = P\delta_{ij}$$

Generally,  $F$  and  $n$  are not parallel



$$\frac{F_i}{S} = \sigma_{ij}n_j$$


Stress Tensor

$$\sigma_{ij} = -T_{ij}$$

Landau  
Lifshitz


# Force

## Action-at-a-distance



Newton  
1687

$m_1, q_1$



$m_2, q_2$




$$F = -G \frac{m_1 m_2}{r^2} \quad F = -\frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2}$$

## Local interaction




Faraday  
1839



# Maxwell Stress

(in Maxwell Theory)




Maxwell

$$\sigma_{ij} = \varepsilon_0 E_i E_j + \frac{1}{\mu_0} B_i B_j - \frac{1}{2} \delta_{ij} \left( \varepsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right)$$

$$\vec{E} = (E, 0, 0)$$


$$T_{ij} = \begin{pmatrix} -E^2 & 0 & 0 \\ 0 & E^2 & 0 \\ 0 & 0 & E^2 \end{pmatrix}$$

- Parallel to field: **Pulling**
- Vertical to field: **Pushing**



# Maxwell Stress


(in Maxwell Theory)



$$T_{ij} v_j^{(k)} = \lambda_k v_i^{(k)}$$

$(k = 1, 2, 3)$

length:  $\sqrt{|\lambda_k|}$




## Definite physical meaning

- Distortion of field, line of the field
- Propagation of the force as local interaction

# Quark—Anti-quark system


Formation of the flux tube  $\rightarrow$  confinement




## Previous Studies on Flux Tube

- Potential
- Action density
- Color-electric field

so many studies...




Cea+ (2012)



Cardoso+ (2013)



# Stress Tensor in $Q\bar{Q}$ System




Yanagihara+, 1803.05656

Lattice simulation  
SU(3) Yang-Mills

$a=0.029$  fm

$R=0.69$  fm

$t/a^2=2.0$




**Definite physical meaning**

- Distortion of field, line of the field
- Propagation of the force as local interaction
- Manifestly gauge invariant


# Lattice Setup

- ❑ SU(3) Yang-Mills (Quenched)
- ❑ Wilson gauge action
- ❑ Clover operator
- ❑ APE smearing / multi-hit
- ❑ fine lattices ( $a=0.029-0.06$  fm)
- ❑ continuum extrapolation
- ❑ Simulation: bluegene/Q@KEK



Yanagihara+, 1803.05656




$\beta$	$a$ [fm]	$N_{\text{size}}^4$	$N_{\text{conf}}$	$R/a$		
6.304	0.058	$48^4$	140	8	12	16
6.465	0.046	$48^4$	440	10	–	20
6.513	0.043	$48^4$	600	–	16	–
6.600	0.038	$48^4$	1,500	12	18	24
6.819	0.029	$64^4$	1,000	16	24	32
$R$ [fm]				0.46	0.69	0.92





# Continuum Extrapolation at mid-point




□  $a \rightarrow 0$  extrapolation with fixed  $t$



# $t \rightarrow 0$ Extrapolation at mid-point



- $a \rightarrow 0$  extrapolation with fixed  $t$
- Then,  $t \rightarrow 0$  with three ranges



# Stress Distribution on Mid-Plane


From rotational symm. & parity

EMT is diagonalized  
in Cylindrical Coordinates

$$T_{cc'}(r) = \begin{pmatrix} T_{rr} & & & \\ & T_{\theta\theta} & & \\ & & T_{zz} & \\ & & & T_{44} \end{pmatrix}$$

$$T_{rr} = \vec{e}_r^T T \vec{e}_r$$



$$T_{\theta\theta} = \vec{e}_\theta^T T \vec{e}_\theta$$



Degeneracy  
in Maxwell theory

$$T_{rr} = T_{\theta\theta} = -T_{zz} = -T_{44}$$

# Mid-Plane






**Continuum  
Extrapolated!**

In Maxwell theory

$$T_{rr} = T_{\theta\theta} = -T_{zz} = -T_{44}$$


- Degeneracy:  $T_{44} \simeq T_{zz}, \quad T_{rr} \simeq T_{\theta\theta}$
- Separation:  $T_{zz} \neq T_{rr}$
- Nonzero trace anomaly  $\sum T_{cc} \neq 0$

# Mid-Plane




- Degeneracy:  $T_{44} \simeq T_{zz}, \quad T_{rr} \simeq T_{\theta\theta}$
- Separation:  $T_{zz} \neq T_{rr}$
- Nonzero trace anomaly  $\sum T_{cc} \neq 0$

# Force



Force from Potential

$$F_{\text{pot}} = -\frac{dV}{dR}$$




Force from Stress

$$F_{\text{stress}} = \int_{\text{mid.}} d^2x T_{zz}(x)$$




# Force




Force from Potential

$$F_{\text{pot}} = -\frac{dV}{dR}$$



Force from Stress

$$F_{\text{stress}} = \int_{\text{mid.}} d^2x T_{zz}(x)$$




Newton  
1687




Faraday  
1839

# Force




Force from Potential

$$F_{\text{pot}} = -\frac{dV}{dR}$$




Force from Stress

$$F_{\text{stress}} = \int_{\text{mid.}} d^2x T_{zz}(x)$$



Newton  
1687



Faraday  
1839

# Abelian-Higgs Model

Yanagihara, Iritani, MK, in prep.

## Abelian-Higgs Model

$$\mathcal{L}_{\text{AH}} = -\frac{1}{4}F_{\mu\nu}^2 + |(\partial_\mu + igA_\mu)\phi|^2 - \lambda(\phi^2 - v^2)^2$$

**GL parameter:**  $\kappa = \sqrt{\lambda}/g$


- type-I :  $\kappa < 1/\sqrt{2}$
- type-II :  $\kappa > 1/\sqrt{2}$
- Bogomol'nyi bound :  
 $\kappa = 1/\sqrt{2}$

**Infinitely long tube**

- degeneracy  
 $T_{zz}(r) = T_{44}(r)$  Luscher, 1981
- momentum conservation  
 $\frac{d}{dr}(rT_{rr}) = T_{\theta\theta}$

# Stress Tensor in AH Model infinitely-long flux tube

Bogomol'nyi bound :  $\kappa = 1/\sqrt{2}$



$$T_{rr} = T_{\theta\theta} = 0$$


de Vega, Schaposnik, PRD**14**, 1100 (1976).

# Stress Tensor in AH Model

## infinitely-long flux tube


Type-I

$\kappa = 0.1$



Type-II

$\kappa = 3.0$



- No degeneracy bw  $T_{rr}$  &  $T_{\theta\theta}$
- $T_{\theta\theta}$  changes sign

conservation law



$$\frac{d}{dr} (rT_{rr}) = T_{\theta\theta}$$

# Stress Tensor in AH Model

## infinitely-long flux tube

Type-I

$$\kappa = 0.1$$



- No degeneracy bw  $T_{rr}$  &  $T_{\theta\theta}$
- $T_{\theta\theta}$  changes sign





Inconsistent with  
lattice result

$$T_{rr} \simeq T_{\theta\theta}$$

# Flux Tube with Finite Length

$R=0.92$  fm




**Left:**  $T_{zz}(o), T_{rr}(o)$  reproduce lattice result

**Right:** A parameter satisfying  $T_{rr} \approx T_{\theta\theta}$






No parameter can reproduce lattice data at  $R=0.92$  fm.



# Summary




- The analysis of energy-momentum tensor on the lattice is now available, and various studies are ongoing!
  - gradient flow method
  - higher-order perturbative coefficients





# Summary

- The analysis of energy-momentum tensor on the lattice is now available, and various studies are ongoing!
  - gradient flow method
  - higher-order perturbative coefficients



## □ So many future studies

- Flux tube at nonzero temperature
- EMT distribution inside hadrons
- viscosity / other operators / instantons / full QCD

Equations and Correlations

specific heat, ...

$$c_V \sim \langle \delta T_{00}^2 \rangle$$

$$\eta = \langle T_{12}; T_{12} \rangle$$

If we have  $T_{\mu\nu}$

flux tube / hadrons

stress distribution

hadron structure

vacuum configuration

mixed state on 1<sup>st</sup> transition

Vacuum

# A Naïve Question



$$\times T_{\mu\nu}(x)$$

Put **a single quark** in QCD vacuum

How does **energy density and stress** behave in this system?

backup

# EMT on the Lattice: Conventional

## Lattice EMT Operator Caracciolo+, 1990

$$T_{\mu\nu} = Z_6 T_{\mu\nu}^{[6]} + Z_3 T_{\mu\nu}^{[3]} + Z_1 (T_{\mu\nu}^{[1]} - \langle T_{\mu\nu}^{[1]} \rangle)$$

$$T_{\mu\nu}^{[6]} = (1 - \delta_{\mu\nu}) F_{\mu\rho}^a F_{\nu\rho}^a, \quad T_{\mu\nu}^{[3]} = \delta_{\mu\nu} \left( F_{\mu\rho}^a F_{\nu\rho}^a - \frac{1}{4} F_{\rho\sigma}^a F_{\rho\sigma}^a \right), \quad T_{\mu\nu}^{[1]} = \delta_{\mu\nu} F_{\rho\sigma}^a F_{\rho\sigma}^a$$


□ Fit to thermodynamics:  $Z_3, Z_1$

□ Shifted-boundary method:  $Z_6, Z_3$  Giusti, Meyer, 2011; 2013;  
Giusti, Pepe, 2014~; Borsanyi+, 2018

## Multi-level algorithm

□ effective in reducing statistical error of correlator Meyer, 2007;  
Borsanyi, 2018;  
Astrakhantsev+, 2018

# Gradient Flow Method



Take Extrapolation  $(t, a) \rightarrow (0, 0)$

$$\langle T_{\mu\nu}(t) \rangle_{\text{latt}} = \langle T_{\mu\nu}(t) \rangle_{\text{phys}} + C_{\mu\nu} t + D_{\mu\nu} \frac{a^2}{t} + \dots$$


$O(t)$  terms in SFTE lattice discretization

# Numerical Simulation

FlowQCD,  
PRD94, 114512 (2016)

- Expectation values of  $T_{\mu\nu}$
- SU(3) YM theory
- Wilson gauge action
- Parameters:
  - $N_t = 12, 16, 20-24$
  - aspect ratio  $5.3 < N_s/N_t < 8$
  - 1500~2000 configurations
- Scale from gradient flow
  - $aT_c$  and  $a\Lambda_{\overline{MS}}$




FlowQCD, 1503.06516



$T/T_c$	$\beta$	$N_s$	$N_t$	Configurations
0.93	6.287	64	12	2125
	6.495	96	16	1645
	6.800	128	24	2040
1.02	6.349	64	12	2000
	6.559	96	16	1600
	6.800	128	22	2290
1.12	6.418	64	12	1875
	6.631	96	16	1580
	6.800	128	20	2000
1.40	6.582	64	12	2080
	6.800	128	16	900
	7.117	128	24	2000
1.68	6.719	64	12	2000
	6.941	96	16	1680
	7.117	128	20	2000
2.10	6.891	64	12	2250
	7.117	128	16	840
	7.296	128	20	2040
2.31	7.200	96	16	1490
	7.376	128	20	2020
	7.519	128	24	1970
2.69	7.086	64	12	2000
	7.317	96	16	1560
	7.500	128	20	2040

# Mid-Point Correlator

$$\langle T_{44}(\tau_{\text{mid}})T_{44}(0) \rangle \quad \langle T_{44}(\tau_{\text{mid}})T_{11}(0) \rangle \quad \langle T_{41}(\tau_{\text{mid}})T_{41}(0) \rangle$$



- (44;11), (41;41) channels : confirmation of FRR
- (44;44) channel: **new** measurement of  $c_V$

$$c_V = \frac{\langle \delta E^2 \rangle}{VT^2}$$



$c_V/T^3$				
$T/T_c$	$C_{44;44}(\tau_m)$	Ref.[19]	Ref.[11]	ideal gas
1.68	17.7(8) $^{+2.1}_{-0.4}$	22.8(7)*	17.7	21.06
2.24	17.5(0.8) $^{+0}_{-0.1}$	17.9(7)**	18.2	21.06

2+1 QCD:  
Taniguchi+ (WHOT-QCD),  
1711.02262

# SU(3) YM vs Maxwell

**SU(3) Yang-Mills**  
(quantum)


**Maxwell**  
(classical)



Propagation of the force is clearly different  
in YM and Maxwell theories!




# Preparing Static $Q\bar{Q}$





- APE smearing for spatial links
- Multi-hit for temporal links
- No gradient flow for  $W(R, T)$

$$V(R) = - \lim_{T \rightarrow \infty} \log \langle W(R, T) \rangle$$

$$\langle O(x) \rangle_{Q\bar{Q}} = \lim_{T \rightarrow \infty} \frac{\langle \delta O(x) \delta W(R, T) \rangle}{\langle W(R, T) \rangle}$$



# Ground State Saturation





$\beta=6.819$  ( $a=0.029$  fm),  $R=0.46$  fm

Appearance of plateau  
for  $t/a^2 < 4$ ,  $T/a > 15$



Grand state saturation  
under control

# Ground State Saturation



$\beta=6.819$  ( $a=0.029$  fm),  $R=0.46$  fm

Appearance of plateau  
for  $t/a^2 < 4$ ,  $T/a > 15$



Grand state saturation  
under control

# Abelian-Higgs Model

## Abelian-Higgs Model

$$\mathcal{L}_{\text{AH}} = -\frac{1}{4}F_{\mu\nu}^2 + |(\partial_\mu + igA_\mu)\phi|^2 - \lambda(\phi^2 - v^2)^2$$

**GL parameter:**  $\kappa = \sqrt{\lambda}/g$

- type-I:  $\kappa < 1/\sqrt{2}$
- type-II:  $\kappa > 1/\sqrt{2}$
- Bogomol'nyi bound:  
 $\kappa = 1/\sqrt{2}$

## Infinitely long tube

- degeneracy

$$T_{zz}(r) = T_{44}(r) \quad \text{Luscher, 1981}$$

- conservation law

$$\frac{d}{dr}(rT_{rr}) = T_{\theta\theta}$$