

# Finite-size scaling, intermittency and the QCD critical point

F. K. Diakonov in collaboration with:  
N.G. Antoniou, N. Davis, N. Kalntis, A. Kanargias,  
X.N. Maintas and C.E. Tsagkarakis




FACULTY OF PHYSICS, UNIVERSITY OF ATHENS, GREECE

**CPOD 2018**  
Corfu, September 24-28

- 1 Critical region ; scaling properties
- 2 Ising-QCD thermodynamics in the critical region
- 3 Size of the critical region
- 4 Locating the CEP
- 5 Conclusions

# Phase diagram of QCD

A sketch for finite system(s)



Objective: Detection  
(existence?) of the QCD  
Critical Point (CP)



**Equilibrium?** (experiment)  
**3d-Ising** universality class,  
scaling properties with known  
critical exponents

*from R. V. Gavai, Contemporary Physics 57, 350 (2016)*

# 3d-Ising effective action

**3d-Ising effective action** (dimensionless form) for the order parameter  $\phi$  in the critical region:

$$S_{eff} = \int_V d^3\hat{x} \left[ \frac{1}{2} |\hat{\nabla}\phi|^2 + U(\phi) - \hat{h}\phi \right] \quad \text{with}$$

$$U(\phi) = \frac{1}{2} \hat{m}^2 \phi^2 + \hat{m} g_4 \phi^4 + g_6 \phi^6 \quad ; \quad \phi = \beta_c^3 \lim_{\delta V \rightarrow 0} \frac{n_{\uparrow} - n_{\downarrow}}{\delta V}$$

$$\hat{x} = x\beta_c^{-1}, \quad \hat{m} = \beta_c m, \quad m = \xi^{-1}, \quad \hat{h} = h\beta_c^{-1} \quad \xi = \text{correlation length}$$

universal constants  $g_4 \approx 0.97, g_6 \approx 2.1$   $h = \text{ordering field}$

$$\text{Partition function } \mathcal{Z} = \sum_{\{\phi\}} \exp(-S_{eff}[\phi])$$

*M. M. Tsypin, Phys. Rev. Lett. 73, 2015 (1994)*

# Ising-QCD partition function

Constructing the **Ising-QCD** partition function in the critical region:

$$(n_{\uparrow}, n_{\downarrow}) \implies (n_B, n_{\bar{B}})$$

Scaling properties describable restricting to **protons!**

*Y. Hatta and M. A. Stephanov, PRL 91, 102003 (2003)*

Use **constant configurations** for the field  $\phi = \frac{N}{V}$  with  $N = N_p$ :

$$\mathcal{Z}_{IQCD} = \sum_{N=0}^{\Lambda} \zeta^N \exp \left[ -\frac{1}{2} \hat{m}^2 \frac{N^2}{\Lambda} - g_4 \hat{m} \frac{N^4}{\Lambda^3} - g_6 \frac{N^6}{\Lambda^5} \right]$$

with  $\zeta = \exp[(h - h_c)\beta_c]$  ( $h_c = 0$  for 3d-Ising) and  $\Lambda = \frac{V}{V_0}$   
( $V_0 =$  proton volume)

# Ising-QCD partition function (continued)

Thermodynamic quantities in  $\mathcal{Z}_{IQCD}$ :

$$\mathcal{Z}_{IQCD} = \sum_{N=0}^{\Lambda} \zeta^N \exp \left[ -\frac{1}{2} \hat{m}^2 \frac{N^2}{\Lambda} - g_4 \hat{m} \frac{N^4}{\Lambda^3} - g_6 \frac{N^6}{\Lambda^5} \right]$$

## Direct mapping

$h - h_c$  is mapped to  $\mu_B - \mu_c$  in  $\zeta$

$$T \text{ in } \hat{m} = \zeta^{-1} \beta_c$$

$$\zeta = \zeta_{0,\pm} \left| 1 - \frac{T}{T_c} \right|^{-\nu},$$

$\zeta_{0,\pm}$  non-universal with  $\frac{\zeta_{0,+}}{\zeta_{0,-}} = 2$

Volume in  $\Lambda$

## More general

$$h \rightarrow (\mu_B - \mu_c) - \tan \alpha (T - T_c)$$

$$\zeta \rightarrow \zeta_{0,\pm} \left| \frac{T}{T_c} - 1 + \tan \alpha \frac{(\mu_B - \mu_c)}{T_c} \right|^{-\nu}$$

*J.J. Rehr and N.D. Mermin, PRA 8, 472 (1973)*

Robustness for small  $\alpha$ !

$$\left( \nu = \frac{2}{3} \text{ for 3d-Ising} \right)$$

# Ising-QCD partition function (in action)

Use  $\mathcal{Z}_{IQCD}$  to:

- Calculate proton multiplicity moments  $\langle N^k \rangle$  ( $k = 1, 2, \dots$ )



Scaling laws? **critical exponents?**

**Size** of the **critical region?**

- Thermodynamic **response functions** (specific heat, susceptibilities,..)
- **Equation of state** (pressure) in the neighbourhood of the CEP

# Proton multiplicity moments and FSS

For  $\mu_B = \mu_c$ ,  $T = T_c$  we find:

$$\langle N^k \rangle \sim \Lambda^{kq}, \quad q = d_F/d, \quad k = 1, 2, \dots$$

⇓

**Finite size scaling (FSS) law** with  $d_F = \frac{5}{2}$  (and  $d = 3$ )

FSS exponent  $q$  is related to the isothermal critical exponent  $\delta$

$$q = \frac{d_F}{d} = \frac{\delta}{\delta + 1} \quad ; \quad \delta = 5 \text{ (3d - Ising)}$$

Measurement of  $q \Rightarrow$  measurement of  $\delta$

**Unrealistic task**, needs **systems of different sizes**  
**freezing out at the critical point!**



# Finite size scaling and intermittency

Making the unrealistic possible: in **FSS** regime the **local scaling**:

$$\langle n(\mathbf{x})n(\mathbf{x}') \rangle \sim |\mathbf{x} - \mathbf{x}'|^{-(3-d_F)}$$

is valid also **globally** ( $|\mathbf{x} - \mathbf{x}'| = O(V^{1/3})$ ) leading to:

$$\langle N \rangle \propto \Lambda^{\frac{d_F}{3}}$$

**large distance singular** behaviour of **density-density correlator**



Singularity for **small distances** in **proton transverse momentum** space:

$$\lim_{\mathbf{k} \rightarrow \mathbf{k}'} \langle n(\mathbf{k})n(\mathbf{k}') \rangle \sim |\mathbf{k} - \mathbf{k}'|^{-2q} \quad ; \quad q = \frac{d_F}{3} = \frac{5}{6}$$

**Intermittency** (critical opalescence)  $\Rightarrow q$  is observable!

*N.G. Antoniou, N. Davis, F.K. D., PRC 93, 014908 (2016)*

The non-Gaussian kurtosis:

$$\kappa_{nG} = \frac{C_4 - 3C_2^2}{C_2^2} \quad ; \quad C_k = \langle (N - \langle N \rangle)^k \rangle, \quad k = 2, 3, ..$$

becomes **negative** approaching the critical point

*M.A. Stephanov, PRL 107, 052301 (2011)*


Calculate cumulants  $C_k$  and  $\kappa_{nG}$  through  $\mathcal{Z}_{IQCD}$ :

$$\frac{\partial^2}{\partial (\ln \zeta)^2} \ln \mathcal{Z}_{IQCD} = C_2 \quad ; \quad \frac{\partial^4}{\partial (\ln \zeta)^4} \ln \mathcal{Z}_{IQCD} = C_4 - 3C_2^2$$

and explore their behaviour close to the critical point!

# Size of the critical region

Departing from the critical point  $\Rightarrow$  Gradual destruction of the FSS law:  
 $(\zeta = 1, t = 0)$   $\langle N \rangle \sim \Lambda^{\frac{5}{6}}$



Notation:  $\zeta = \exp[(\mu_B - \mu_c)\beta_c]$ ,  $t = \frac{T - T_c}{T_c}$  ( $\alpha = 0$ )

# Size of the critical region (continued)


In a region around  $\zeta = 1$ ,  $t = 0$  (CP) it holds:

$$\langle N \rangle \sim \Lambda^{\tilde{q}}$$

- $\tilde{q} = \frac{3}{4} \Rightarrow$  scaling ( $q$ ) in mean field theory
  - $\tilde{q} = 1 \Rightarrow$  trivial scaling

**Critical region:** region in  $(\ln \zeta, t)$ -plane for which  $\frac{3}{4} < \tilde{q} < 1$

# Size of the critical region (first result)



**Critical region**  $\Delta\mu_B$   
 $\approx 5 \text{ MeV}$

(for  $T_c \approx 160 \text{ MeV}$ )




Very **narrow** along  
 $\mu_B$ -axis

*N.G. Antoniou, F.K. D.,  
X.N. Maintas, C.E. Tsagarakis,  
PRD 97, 034015 (2018)*

# Finite size scaling region

FSS condition:  $\tilde{\zeta}_\infty > V^{1/3}$



Bounds along the  $t$  axis!

**System dependent!**

For **medium** ( $20 < A < 50$ ) size nuclei,

**FSS region:**

$$3 \text{ MeV} < \Delta T < 5 \text{ MeV}$$



(for  $T_c \approx 160 \text{ MeV}$ )



**Narrowness** along  $T$ -axis too

*N.G. Antoniou, F.K. D.,  
arXiv:1802.05857 [hep-ph]*

# Kurtosis within the critical region




**Negative sharp minimum**  
of  $\kappa_{nG}$  at the CP

*N.G. Antoniou, F.K. D., N. Kalntis, A. Kanargias*  
*arXiv:1711.10315 [nucl-th]*

Alternative(s) for the critical region size

*N.G. Antoniou, F.K. D., arXiv:1802.05857 [hep-ph]*

# Critical region size: $\kappa_{nG}$ vs. FSS varying $\alpha$




Critical region size along  $\mu_B$  is  $3 \text{ MeV} \leq \Delta\mu_B \leq 11 \text{ MeV}$  for all  $\alpha$ !

*N.G. Antoniou, F.K. D., arXiv:1802.05857 [hep-ph]*



# Intermittency in Si + Si collisions (NA49, SPS, CERN)



Si + Si central collisions at  $\sqrt{s} = 17.2$  GeV create a final state **within the critical region**:  $\tilde{q} \approx 0.96!$




Use this result to locate CEP!

Ignore **large** statistical errors...

Typical range for spatial correlations [1, 8] fm  
 $\Rightarrow$  tr. mom. space  $M^2 \in [400, 11000]$

# Locating Si + Si final state within the critical region



Line  $\tilde{q} = 0.96$  determines  $\mu_c$   
for known  $T_c$  (Lattice QCD)

recent result:  $T_c = 163 \text{ MeV}$

*S. Datta et al, PRD 95, 054512 (2017)*

Freeze-out of Si\*:

$$(\mu_{Si}, T_{Si}) = (260, 162.2) \text{ MeV}$$

$$T_c = 163 \text{ MeV}$$



$$\ln \zeta_{Si} = 0.0143 \Rightarrow$$

$$\mu_c = 257.7 \text{ MeV}$$

\*: *F. Becattini, J. Manninen and M. Gazdzicki, PRC 73, 044905 (2006)*

# Predictions for NA61/SHINE freeze-out states

Freeze-out conditions for Ar+Sc and Xe+La  $\Rightarrow$  use NA49 results

$$\sqrt{s} = 17.2 \text{ GeV}$$

Freeze-out of central Ar+Sc:


$$(\mu_{ArSc}, T_{ArSc}) = (258, 160.9) \text{ MeV}$$

Freeze-out of central Xe+La:


$$(\mu_{XeLa}, T_{XeLa}) = (251, 158.2) \text{ MeV}$$

*N.G. Antoniou, F.K. D., arXiv:1802.05857*

*[hep-ph]*





# Enriched sketch of the critical region



F. Becattini, et al, *PRC* 90, 054907 (2014); N.G. Antoniou, F.K. D., *arXiv:1802.05857 [hep-ph]*

# Enriched sketch of the critical region for $\alpha \neq 0$



- **Critical (FSS) region is very narrow**  $O(5 \text{ MeV})$  along the  $\mu_B$  and the  $T$  axis.
- Beam energy scan program at RHIC with  $\Delta\mu_B \approx 50 \text{ MeV}$  is very unlikely to approach the critical region.
- **Important NA49 result: freeze-out state of central Si+Si collisions at  $\sqrt{s} = 17.2 \text{ GeV}$  lies within the critical (FSS) region!** (needs accurate measurements to reduce statistical errors)



Can be used as a **guide** for detecting the QCD CEP.

- **Basic strategy: Accurate measurements of FSS exponent  $\tilde{q}$  (intermittency analysis) and corresponding freeze-out parameters  $(\mu_B, T)$  in A+A collisions with  $25 < A < 50$ .**

# Conclusions (continued)

- $\sqrt{s} \approx 17 \text{ GeV}$  seems to be the **appropriate beam energy** for approaching  $\mu_c$ . **Peripheral collisions** can be used for **fine changes in  $T$**  allowing the entrance into the FSS region.

For A+A collisions at  $\sqrt{s} = 17.2 \text{ GeV}$  we propose:

Accurate measurements of  $(\tilde{q}, \mu_B, T)$  in central collisions  
for  $25 < A < 32$ .

Accurate measurements of  $(\tilde{q}, \mu_B, T)$  in peripheral collisions  
for  $32 < A < 50$ .

**Prediction: Strong intermittency effect in peripheral Ar+Sc collisions** at  $\sqrt{s} \approx 17 \text{ GeV}$  (NA61/SHINE experiment).

(See *N.G. Antoniou, F.K. D., arXiv:1802.05857 [hep-ph]*)

Thank you!