



Contribution ID: 1083

Type: **Invited Oral Presentation**

M3Or1C-01 [Invited]: Direct observation of nanometer scale niobium hydride ($\text{Nb}_{1-x}\text{H}_x$) on SRF Nb cavity surface using cryogenic atomic force microscopy (AFM)

Wednesday 24 July 2019 09:30 (30 minutes)

Cryogenic atomic force microscopy (AFM) allowed us to directly observe segregation morphology of nanometer size hydride phases ($\text{Nb}_{1-x}\text{H}_x$), which are responsible for Q degradation of superconducting radio frequency (SRF) Nb cavity at high RF field regime. Although niobium hydride has been recognized with cryo-optical microscopy, electron diffraction or ion mass spectroscopy, identification of nm-scale Nb hydride segregation in terms of saturation temperature has never been accomplished due to difficulties on direct observation at cryogenic temperature. In this study, we address the first direct observation of nanometer size hydride segregation on the surface of superconducting RF niobium cavities as a function of cooling temperature, and quantitatively compare its segregation morphology with respect to state of art cavity treatments.

Author: Dr SUNG, Zuhawn (Fermi National Accelerator Laboratory)

Co-authors: Dr ROMANENKO, Alexander; Dr GRASSELLINO, Anna (Fermi National Accelerator Laboratory); BAFIA, Daniel (Illinois Institute of Technology)

Presenter: Dr SUNG, Zuhawn (Fermi National Accelerator Laboratory)

Session Classification: M3Or1C - Focus Series C: Superconducting RF Cavity Materials III