### 3<sup>rd</sup> Heavy Flavor Meet Indore 2019, March 18-20

#### (more or less) Recent topics in (open) heavy-flavor production

#### **Outline**

- Introduction and structuration
- « Understanding » the R<sub>AA</sub> v<sub>2</sub>
- Lessons from recent collective initiatives (bulk, hadronization,transport,...)
- New observables
- More lessons from recent collective initiatives
- The role of transport
- A word on small systems
- Conclusions


P.B. Gossiaux SUBATECH, UMR 6457


Université de Nantes, IMT Atlantique, IN2P3/CNRS

## Why open heavy flavors in A-A?

- 1.  $\Phi$ : Produced early, number conserved through time evolution (even at LHC)  $\Rightarrow$  signature (hard probes) of early (hot) phase
- 2.  $\Phi$ : Strongly affected by the QGP phase (up to factor 5 in the final spectrum)
- 3.  $\Phi$ : Weakly affected by late time evolution (heavy, colour transparency)
- 4. Interpretation: Allows physical picture based on physical scales ( $t_{relax} \alpha m_Q/T^2 \Rightarrow$  clear hierarchy for s, c and b)
- 5. Theory: Allows *some* pQCD calculations for the initial production and annihilation,...
- 6. Theory: ...

Usually advocated as an ideal probe of dense matter






$$-rac{d}{dt}\langle ec{p}
angle = ec{A}(ec{p},T) = \eta_D(ec{p},T) imes ec{p} \quad \eta_D[\mathrm{fm}^{-1}]: \;\; \mathrm{Relaxation\; rate}$$

$$\frac{d}{dt}\langle\vec{p}_{T,i}\vec{p}_{T,j}\rangle=\kappa_T(\vec{p},T)\delta_{i,j}$$
  $\kappa_T[\mathrm{GeV}^2\mathrm{fm}^{-1}]$ : Transverse diffusion coef. (p space);  $\hat{q}=2\kappa_T=4B_0$ 

Similar in longitudinal direction  $\kappa_L [{
m GeV}^2 {
m fm}^{-1}]$  : Longitudinal diffusion coef.

In general, no relation between these coefficients except  $\kappa_T = \kappa_L$  for p=0.

### Transport coefficients and inelastic processes




$$\Delta \langle \vec{p} \rangle = \vec{A}(\vec{p}, T) \times L + (\Delta \vec{p})_{\text{rad}}$$

- contribution from « radiated » part
- In most of existing schemes:  $(\Delta \vec{p})_{\rm rad} = \mathcal{F}(\eta_D, \kappa_T, \kappa_L, p, L)$

Seeked ransport coeff.





## Transport coefficients at low momentum p≈m<sub>o</sub>

Langevin regime => Einstein relation:  $\kappa = 2TE_{Q}\eta_{D}$ 


$$D_s = \left( = \frac{1}{6} \lim_{t \to \infty} \frac{\langle (\mathbf{x}(t) - \mathbf{x}(0))^2 \rangle}{t} \right)$$

 $T/T_c$ 

For historical reasons, physics displayed as a function of  $2\pi T$  x the spatial diffusion coefficient

$$(2\pi T)D_s = \frac{4\pi T^3}{\kappa} = \frac{2\pi T^2}{E_Q\eta_D} \quad \Rightarrow \quad \tau_{\rm relax} = \eta_D^{-1} = (2\pi T)D_s \times \frac{E_Q}{2\pi T^2}$$
 Gauge for the coupling strength 
$$\begin{array}{c} \text{lQCD results} \\ \text{The sole direct rigorous calculation of the transport coeff to my knowledge} \end{array}$$
 
$$\tau_{\rm relax}(T_c) \approx m_Q[{\rm GeV}] \times (3\pm 1.5) \, {\rm fm}$$
 
$$\begin{array}{c} \text{Topical model of the coupling strength} \\ \text{The sole direct rigorous calculation of the transport coeff to my knowledge} \\ \end{array}$$


## Lanscape of HF theory and modeling in URHIC



### **Today Stakes and Motivation**

- Very subttle interplay between fundamental theory and experimental data to build and calibrate / constrain the models in order to better understand HF – QGP interaction (including QGP constituents)
- Then maybe one day: probing the bulk!
- On a more phenomenological and data driven level:
   Are we able to constrain / extract the in medium transp. coefficients with a « reasonnable » precision ?
  - => What is the impact of various « extra » ingredients ?
  - Do we agree on what « reasonnable » means (x 2 ? 50% ?, 10% ?)
- Need to intercompare the approaches.


Big motivation of OHF community



Major goal for the future: from indication to real quantification!

### A bit of structure

HQ propagation in QM & URHIC...




## pQCD inspired models (f.i. Nantes)


#### **Colisional component**

- One-gluon exchange model: reduced IR regulator  $\lambda$  m<sup>2</sup><sub>D</sub> in the hard propagator, fixed on HTL Energy loss
- Running coupling  $\alpha_{\text{eff}}\left(\textbf{t}\right)$  and self consistent Debye mass

$$m_{Dself}^{2}(T) = (1+n_{f}/6) 4\pi\alpha_{eff}(m_{Dself}^{2})T^{2}$$



#### Radiative component



- Extention of Gunion-Bertsch approximation beyond mid-rapidity and to finite mass  $m_{O}$  ) distribution of induced gluon radiation per collision ( $\Delta E_{rad} \alpha E L$ ):
  - $P_g(x, \mathbf{k}_\perp, \mathbf{q}_\perp, m_Q) = \frac{3\alpha_s}{\pi^2} \frac{1 x}{x} \left( \frac{\mathbf{k}_\perp}{\mathbf{k}_\perp^2 + x m_Q^2} \frac{\mathbf{k}_\perp \mathbf{q}_\perp}{(\mathbf{k}_\perp \mathbf{q}_\perp)^2 + x m_Q^2} \right)^2$

LPM effect for moderate gluon energy


## Quasi particle models (f.i DQPM)

 Non perturbative effects near Tc are captured by α<sub>s</sub>(T), leading to thermal masses/widths, determined from fits to IQCD EoS.

A. Peshier et al. PLB 337 (1994), PRD 70 (2004); M. Bluhm et al. EPJC 49 (2007); W. Cassing et al. NPA 795 (2007)

- Relaxation rates larger then in pQCD for all T relevant for QGP, slightly smaller than the ones from TAMU
- H. Berrehrah et al, PHYSICAL REVIEW C 90, 064906 (2014)
- Implemented for HF dynamics in e.g. PHSD (full off-shell, off-equilirium transport).
- T. Song et al. PRC 92 (2015), PRC 93 (2016)

#### **But also CATANIA**



## Potential models (TAMU)

Thermodynamic T-matrix approach, T = V + VGT, given by a two-body driving kernel V, estimated from the IQCD internal/free energy for a static Q-Qbar pair; increase of coupling with QGP at small momentum

D. Cabrera, R. Rapp PRD 76 (2007); H. van Hees, M. Mannarelli, V. Greco, R. Rapp PRL 100 (2008)


Comprehensive sQGP approach for the EoS, light quark & gluon spectral functions, quarkonium correlators and HQ diffusion.

F. Riek, R. Rapp PRC 82 (2010); S. Liu, R. Rapp arxiv:1612.09138

Resonance correlations in the T-matrix naturally lead to recombination (resonance recombination model) near T<sub>c</sub> from the same underlying interactions!


M. He, R. Fries, R. Rapp PRC 82 (2010), PRC 86 (2012)

Implemented through Langevin dynamics in hydro evolution or in **URQMD** 




#### A bit of structure

No-Model approach (« data driven »)



### A bit of structure

HQ propagation in QM & URHIC...




## **Models & Effective Theories**

	elastic	Elastic + radiative	radiative	Other
Transport coefficient based (LV,)	TAMU POWLANG HTL Catania LV	Duke	ASW	ADS/CFT POWLANG IQCD  DABMOD (poster QM R. Katz)  S. Li et al, arXiv:1803.01508
Cross section (or   M   <sup>2</sup> ) based (Boltzmann,)	AMPT MC@sHQ el URQMD PHSD Catania BM	Djordjevic et al MC@sHQ el + rad BAMPS CUJET3 HYDJET++ Abir and Mustafa LBL-CCNU VNI/BMS LIDO (poster QM W. Ke)	SCET <sub>G,M</sub>	

Red: Transport models

## Basic Consequences of HQ interaction with QGP for the R<sub>AA</sub>

#### The pattern seen in the data



#### The acknowledged effects

Flow bump: due to

- (radial) flow of the medium and coupling at small p<sub>T</sub>
- recombination with light quarks

shadowing: due to initial state nuclear effects

Quenching & energy loss: due to


- elastic and inelastic scatterings
- opacity of the medium

Italic: extrinsinc to the HF coupling with QGP AKA « energy loss model»

- Dominated by elastic interactions
- m<sub>Q</sub> >> T => needs « many » collisions to equilibrate
- Physics close to « Langevin »

## Basic Consequences of HQ interaction with QGP for the R<sub>AA</sub>

#### The pattern seen in the data



#### The acknowledged effects

Flow bump: due to

- (radial) flow of the medium and coupling at small p<sub>T</sub>
- recombination with light quarks


shadowing: due to initial state nuclear effects

Quenching & energy loss: due to


- elastic and inelastic scatterings
- opacity of the medium

- Dominated by radiative energy loss (with important coherence effects: $\Delta E_{
  m rad} \propto C_A \hat{q} L^2$ )
- Eikonal regime (propagation along straight lines)
- 1 single transport coefficient dominates the whole physics:  $\hat{q} \propto \kappa_T$
- HQ do not equilibrate with the medium
- $m_Q$  becomes a subscale of the physics  $(m_Q << p_T)$

## Basic Consequences of HQ interaction with QGP for the R<sub>AA</sub>



- Interplay between elastic and radiative interactions...
- ... whose dominance depends on the path length
- Fluctuations need to be taken properly into account
- Elastic component: Not clear that Langevin regime still applies (harder and harder collisions)
- 3 transport coefficients in momentum space  $(\eta, \kappa_L, \kappa_T)$  are « only » constrained by Fluc. Dissip. Th.
- Radiative component acquires NLO in m<sub>Q</sub>/p and starts being sensitive to  $\hat{e}$  and  $\hat{e}_2$




S. Cao et al, Phys. Rev. C 94, 014909 (2016)

$$\begin{split} \frac{dN_g}{dy dl_{\perp}^2 d\tau} &= 2 \; \frac{\alpha}{\pi} \; P(y) \; \frac{1}{l_{\perp}^4} \left(\frac{1}{1+\chi}\right)^4 \; \sin^2\left(\frac{l_{\perp}^2}{4l^-(1-y)}(1+\chi) \; \tau\right) \\ &\times \left[\left\{\left(1-\frac{y}{2}\right)-\chi+\left(1-\frac{y}{2}\right)\chi^2\right\} \; \hat{q} + \frac{l_{\perp}^2}{l^-}\chi \left(1+\chi\right)^2 \; \hat{e} + \frac{l_{\perp}^2}{(l^-)^2}\chi \left(\frac{1}{2} - \frac{11}{4}\chi\right) \hat{e}_2\right] \end{split}$$

Abir and Majumder, Phys. Rev. C 94, 054902 (2016)

See as well Aichelin, Gossiaux & Gousset, PRD (2013)

### Basic Consequences of HQ interaction with QGP for the v<sub>2</sub>



Small pT: height of v2 at low  $p_T$  sensitive to:

- Bulk anisotropy, mostly at the late times
- The drag force acting locally on HF

high  $p_T$  non-0  $v_2$  is due to anisotropic Eloss (same ingredients as for the RAA + geometrical anisotropy of initial distribution of matter)

intermediate  $p_T$ : onset and offset of many competing effects.

!!! Alternative pointed out recently within transport model (AMPT & MPC) study: so-called « escape mechanim » characterized by a large  $v_2$  component stemming from  $N_{coll} \approx 1$ L. He et al, Physics Letters B753 (2016) 506




#### 2 Important remarks:

- Any energy loss model, even the roughest one, will generate these typical structures in the  $R_{AA}$  and the  $v_2$ . Getting a correct **quantitative** agreement is much more involved.
- Quantitative predictions also depends on those « extra ingredients »

## Models vs DATA at LHC (Sapore Gravis Report compilation)


Eur. Phys. J. C (2016) 76




Despite various prescriptions for Energy loss, a lot of models can cope with the data

## Models vs DATA at LHC (Sapore Gravis Report compilation)

Eur. Phys. J. C (2016) 76





### Tension between $R_{AA}$ and $v_2$ (at low $p_T$ ): the Catania Cocktail



$$\tau_{\rm relax} = \eta_D^{-1} = (2\pi T)D_s \times \frac{m_Q}{2\pi T^2}$$

S.K. Das et al, Physics Letters B747 (2015) 260






 $\eta_D \alpha T^2$ : pQCD (fixed  $\alpha_s$ ), AdS/CFT


 $\eta_{\rm D} \alpha$  T: pQCD (running  $\alpha_{\rm s}$ )

 $\eta_D \alpha T^0$ : QPM, DQPM, U potential (TAMU)

Tuned to reproduce  $R_{AA} => Larger$  coupling with the bulk near  $T_c$  (when the hydro  $v_2$  has fully developped)  $=> Larger v_2$ 



### Tension between $R_{AA}$ and $v_2$ (at low $p_T$ ): the Catania Cocktail




#### **Extra increase from LV => Boltzmann dynamics**

#### Should be seen as a *decrease* passing from Boltzmann =>LV


In models considering  $\gamma$   $\alpha$  T<sup>0</sup> like QPM, DQPM, TAMU: microscopic d $\sigma$ /d $\theta$  generate more diffusion at large angles

=> Encoding the physics into Langevin scheme, we do not describe properly the fluctuations at large momentum (as seen f.i. in the Tsallis like asymtotic distribution).




=> For dynamical evolution, one needs to cranck down the interaction and the FP coefficients in order to reproduced a  $\times$  given  $\times$  R<sub>AA</sub>

=> Smaller « extracted » coefficients ( $\approx$  -20%) & smaller  $v_2$ .



S.K. Das et al, Physics Letters B747 (2015) 260

### Tension between $R_{AA}$ and $v_2$ (at low $p_T$ ): the Catania Cocktail




Nice guideline but need:


- To consider extra ingredients (bulk, initial v<sub>2</sub>,...)
- To assess the uncertainties on « Coal » and « HR »
- ... before one can think of ruling out other trends for  $\eta_D$ .

- Nonperturbative effects near Tc are captured by  $\alpha_{\rm s}$  (T), determined from fits to IQCD pressure and interation measure and leading to pretty large thermal masses.
- Bulk of massive partons described by Boltzmann dynamics, with gain and loss term is tuned to a fixed  $\eta/s(T)$ M. Ruggieri et al., Phys. Rev. C 89, 054914 (2014).

$$g^{2}(T) = \frac{48\pi^{2}}{(11N_{C} - 2N_{f}) \ln \left[\lambda \left(\frac{T}{T_{C}} - \frac{T_{S}}{T_{C}}\right)\right]^{2}}.$$

- Hadronization performed through (local) coalescence + fragmentation
- Good common description of R<sub>AA</sub> and v<sub>2</sub>






- Nonperturbative effects near Tc are captured by  $\alpha_s$  (T), determined from fits to IQCD pressure and interation measure and leading to pretty large thermal masses.
- Bulk of massive partons described by Boltzmann dynamics, with gain and loss term is tuned to a fixed  $\eta/s(T)$  M. Ruggieri et al., Phys. Rev. C 89, 054914 (2014).

$$g^{2}(T) = \frac{48\pi^{2}}{(11N_{C} - 2N_{f}) \ln\left[\lambda \left(\frac{T}{T_{C}} - \frac{T_{S}}{T_{C}}\right)\right]^{2}}$$

- Hadronization performed through (local) coalescence + fragmentation
- Good description of R<sub>AA</sub> and v<sub>2</sub>
- ... Leading to an indirect extraction of the D<sub>s</sub> coefficient... in the bulk of IQCD calculations



...and slowly increasing -> LO pQCD

### Data driven extraction of transport coefficients

NEW

Y. Xu et al arXiv:1710.00807v1

Poster Y. Xu, QM 2018

except ghat)

« Minimal model approach »: Bayesian analysis by the Duke group

$$\frac{d\vec{p}}{dt} = -\eta_D(p)\vec{p} + \vec{\xi} + \vec{f}_g \leftarrow \frac{dN_g}{dxdk_\perp^2 dt} = \frac{2\alpha_s P(x)\hat{q}_g}{\pi k_\perp^4} \sin^2\left(\frac{t - t_i}{2\tau_f}\right) \left(\frac{k_\perp^2}{k_\perp^2 + x^2 M^2}\right)^4$$


Usual Langevin  $\langle \xi_i(t)\xi_j(t')
angle = \kappa \delta_{ij}\delta(t\!-\!t')$  with  $\kappa=\frac{2T^2}{D_s}$ 

+ coal / frag hadronization and hadronic rescattering

$$D_{s}(T,p) = \frac{1}{1+(\gamma^{2}p)^{2}} (D_{s}2\pi T)^{\text{lin}}(T;\alpha,\beta) + \frac{(\gamma^{2}p)^{2}}{1+(\gamma^{2}p)^{2}} (D_{s}2\pi T)^{\text{pQCD}}(T,p)$$

$$(D_s 2\pi T)^{\text{pQCD}} = 8\pi/(\hat{q}/T^3)$$
$$(D_s 2\pi T)^{\text{linear}} = \alpha \cdot (1 + \beta(T/T_c - 1))$$

Encodes possible Non Perturbative Effects around  $T_c$  through parameters  $\alpha$  (magnitude),  $\beta$  (slope) and  $\gamma$  (inverse momentum range of NP effects)




### Duke "Bayesian approach"


- Choice of 60 « prior » for which the physical observabes are calculated
- Gaussian emulator to build a fast surrogate of physics
- Random walk throughout parameter space, with acceptance and rejection according to likelihood (with all uncertainties assumed to be uncorrelated).





$$(D_s 2\pi T)^{\text{linear}} = \alpha \cdot (1 + \beta (T/T_c - 1))$$




Rather small value => strong coupling!











#### Duke "Bayesian approach" vs models




All together (IQCD, Bayesian analysis and most recent models) make a strong case for NP physics « around  $T_c$  » and at «low»  $p_T$  ... needs to be precised in the future

- Does not mean that all models inhold the same physics...
- D<sub>s</sub> (p=0) does not represent the full physics (different momentum dependences)
- D<sub>s</sub> (finite p) in Duke's el + rad approach should not be compared to the same quantity in purely elastic models (additional contribution to energy loss due to the rad. part)




#### Status of high p<sub>T</sub> HQ

Over the past years, steady development of several sophisticated pQCD-based radiative Energy loss schemes in order to cope with the radiation of energetic partons: BDMPSZ, AMY, higher twist, DGLV, SCET... some of them leading to successful comparison with the data in their numerical implementation...



### Status of high p<sub>T</sub> HQ

Over the past years, steady development of several sophisticated pQCD-based radiative Energy loss schemes in order to cope with the radiation of energetic partons: BDMPSZ, AMY, higher twist, DGLV, SCET... some of them leading to successful comparison with the data in their numerical implementation...



... Overall success of pQCD for describing the gluon radiation from a hot medium.


Beware :  $\hat{q}$  is « just » an indirect result in some of those formalisms

### Status of high p<sub>T</sub> HQ: prospects

... Overall success of pQCD for describing the gluon radiation from a hot medium.... However, in a regime where  $m/p_T$  is small => The genuine mass ordering has still to be quantified and scrutinized more precisely between various approaches

Clear case for b-quark physics

#### Comparing some callibration curve for main approaches:



R. Rapp et al, Nuclear Physics A 979 (2018) 21–86 arXiv: 1803.03824 EMMI Task Force


- Rather good agreement between Djordjevic ans SCET<sub>G</sub> (same  $\alpha_s$ =), and also with LBL-CCNU (although smaller value of  $\alpha_s$ .
- Trend reproduced by Nantes implementation of BDMPS for intermediate pT (for which it should apply)
- Much increased value for the CUJET3 stemming from the assumption of magnetic monopoles « around » Tc => If all other ingredients are under control, offers a unique opportunity to probe the QGP dof with high  $p_T$  partons


### More on v<sub>2</sub> and higher flow harmonics


Goal: better understand the coupling btwn HQ and QGP by:


- departing from the nearly equilibrated regime
- exploring the consequences of inertia due to large m<sub>o</sub>
- Our observation: reduction of  $v_n/\epsilon_n$  for a) larger n, b) larger mass and c) larger centrality...

Does the HQ flow follows the light sector ? => EbE analysis










#### More on v<sub>2</sub> and higher flow harmonics



# Impact of T-dependence of Ds on T on the correlation C





Confirms previous trends up to the 5th moment! No correlation expected for pQCD-like

Poster R. Katz, QM 2018

DABMOD: extension of the Eloss scenarios considered: same trend!

Seems promissing, but systematic study needs to be performed on an ensemble of bulks + initial fluctuations







Interplay between positive and negative contributions

Need to understand the  $v_2$ (time), especially in light of the recently advocated escape mechanism

R. Rapp J.Phys. G36 (2009) 064014

Steady increase of v<sub>2</sub> (all c quarks)





« charm quarks more hydrodynamic then light quarks » Ok, but need to think twice about it!!!

## Recent Collective actions beyond Sapore Gravis

- **Heavy Quark Working Group** (convener: X-N Wang); in the spirit of the Jet Collaboration, the goal is, in a first stage, to :
  - Collect and compare the transport coefficients from various models,
  - Measure and understand their consequences by first studying a simpler brick problem
  - Estimate some systematics + uncertainties
- LBL-CCNU (XN Wang, S. Cao)
- Duke (S. Bass , S. Cao, M. Nahrgang,
   Y. Xu)
- Catania (V. Greco, S. Das, S. Plumari,
   F. Scardina)
- TAMU (R. Rapp, M. He)
- Frankfurt pHSD (E. Bratkovskaya, T. Song, H. Berrehrah)
- Nantes (J. Aichelin, PB Gossiaux, M. Nahrgang)



S. Cao et al,

arXiv: 1809.07894


## **HQ Working Group**

S. Cao et al, arXiv: 1809.07894

Collect and compare the transport coefficients from various models:

c-quarks

What is used by various models to fit the data



- Obviously not satisfying: Larger dispersion than the predictions for concrete observables... WHY?
- Because of « extra ingredients », chosen differently in each model (partly) !!!
- More complex then for the case of jets (several FP coefficients)

## Recent Collective actions beyond Sapore Gravis

- EMMI Rapid Reaction Task Force (organizers: R. Rapp, PB Gossiaux, A. Andronic, R. Averbeck,, S. Maschiocchi):
  - Global strategy to extract the diffusion coefficient from the intercomparison between models and data
  - Collect and analyse all ingredients from various models
  - Identify constrains from IQCD
  - Initiate discussions to assess the limitations of some existing models.

R. Rapp\*1, P.B. Gossiaux\*2, A. Andronic\*3,4, R. Averbeck\*3, S. Masciocchi\*3, A. Beraudo<sup>5</sup>, E. Bratkovskaya<sup>3,6</sup>, P. Braun-Munzinger<sup>3,7</sup>, S. Cao<sup>8</sup>, A. Dainese<sup>9</sup>, S.K. Das<sup>10,11</sup>, M. Djordjevic<sup>12</sup>, V. Greco<sup>11,13</sup>, M. He<sup>14</sup>, H. van Hees<sup>6</sup>, G. Inghirami<sup>3,6,15,16</sup>, O. Kaczmarek<sup>17,18</sup> Y.-J. Lee<sup>19</sup>, J. Liao<sup>20</sup>, S.Y.F. Liu<sup>1</sup>, G. Moore<sup>21</sup>, M. Nahrgang<sup>2</sup>, J. Pawlowski<sup>22</sup>, P. Petreczky<sup>23</sup> S. Plumari<sup>11</sup>, F. Prino<sup>5</sup>, S. Shi<sup>20</sup>, T. Song<sup>24</sup>, J. Stachel<sup>7</sup>, I. Vitev<sup>25</sup>, and X.-N. Wang<sup>26,18</sup>

Goal to attack the problem with a broad view right from the beginning...

R. Rapp et al,

Nuclear Physics A 979 (2018) 21–86

arXiv: 1803.03824 EMMI Task Force

(20 monthes since first meeting)

#### **EMMI RRTF**

#### Topics:

- Initial spectra and shadowing
- Bulk evolution and consequence on HF observables
- Transport implementation
- Hadronization
- Microscopic models for HF energy loss and constrains from QCD at low and high momentum
- Future observables

#### EMMI RRTF: Consequences from the **bulk choice** (and partly transport)


<u>Question</u>: What is the role of the different medium evolution models, and how do different predictions for the bulk cooling and expansion temperature in the current models manifest themselves in HF observables?

Method: adopt a common  $\alpha_s$ =0.4-pQCD x 5 cross section for thermal light partons acting on c-quarks (or associated FP coefficients for models based on FP) in all frameworks.


One Interaction for all of them; not aimed at

R. Rapp et al, arXiv: 1803.03824


Protons from the bulk at FO



c quarks at FO



reproducing the data !!!



This allows to probe the effect of the bulk with a mechanism that has a D<sub>s</sub> roughly similar to the one extracted from IQCD

For most bulks:

$$R_{AA}(c, 10 \, \text{GeV}) \approx 0.3 - 0.4$$


For 30%-50%:


$$R_{AA}(c, 10 \, \text{GeV}) \approx 0.4 - 0.6$$

No feed down! Some correlation between dN(p) and  $R_{AA}(c)$  but not systematic


#### EMMI RRTF: Consequences from the **bulk choice**

#### Protons from the bulk at FO





#### c quarks at FO



N.B.: LBL-CCNU could not implement scattering on thermal- massive partons

For most bulks:

$$v_2(c, p_T = 4 \, \text{GeV}) \approx 0.4 - 0.6$$

Max v<sub>2</sub> reached between 2 and 4 GeV/c

Some correlation between  $v_2(p)$  and  $v_2(c)$  but not systematic

- Some correlation between  $R_{AA}(c)$  and  $v_2(c)$  from various bulks, but rather large residuals => Non « scalable » bulkS
- Adopting a (limited number of) common bulk(s) would permit to shrink the residuals in the « extraction » of the optimal transport coefficients.


#### HQ-Hadronization (would deserve a full talk)

#### Acknowledged:

towards the end of QGP, hadronization of (of equilibrium) HQ can proceed through a dual mechanism:

#### Low $p_T$ :

- The quark partner(s) are already present in the hot cooling medium
- New specific recombination mechanism; no obvious calibration
- The footprint of reconfinment (?!)
- Crucial to explain the flow bump in R<sub>AA</sub>(D) and sizable v<sub>2</sub>(D) => large impact.



#### High $p_T$ :

- The quark partner(s) needed to create the HF-hadron have to be generated from the vacuum
- « usual » fragmentation calibrated on p+p and e++e- data (Petersen,...)

#### <u>Uncertain (and not disputed enough):</u>

Genuine physical recombination process!

# Regained interest in view of the recent $\Lambda_c$ measurement

#### **HQ** - Recombination

#### <u>Instantaneous coalescence:</u>

Greco, Ko & Levai. Phys. Rev. C 68 (2003) 034904 V. Greco, C. Ko, and R. Rapp, Phys. Lett. B 595 (2004) 202 Y. Oh et al, Phys. Rev. C79 (2009) 044905 R. J. Fries, V. Greco, and P. Sorensen, Ann. Rev. Nucl. Part. Sci. 58 (2008) 177



Poster V. Minissale, QM 2018

time

Wigner density of hadronic states
x-p correlation (6D) or « just » in
p space (3D/1D)

x-p correlation (6D p space (3D/1D)

Known issues with energy momentum conservation; small effects at intermediate  $p_T$ ? space


Also: one free parameter per resonances

#### Latest Catania's coalesence model:

- Full 6D coalescence
- New normalization to impose  $P_{coal} \rightarrow 1$  for  $p_T \rightarrow 0$
- Resonance decay
- Mini jet contribution
- Inclusion of  $\Lambda_c$  baryonic states
  - => reduction of R<sub>AA</sub>(D) at small p<sub>T</sub>
  - => increase of  $\Lambda_c/D^0$  wrt pp and pPb.

// S. Plumari, QM 2018

41



#### **HQ** - Recombination


#### **Resonance Recombination Model:**

$$\left(\frac{\partial}{\partial t} + \vec{v} \cdot \vec{\nabla}\right) F_M(t, \vec{x}, \vec{p}) = -\frac{\Gamma}{\gamma_p} F_M(t, \vec{x}, \vec{p}) + \beta(\vec{x}, \vec{p})$$

time


Ravagli and d Rapp, Phys. Lett. B

655 (2007) 126{131,



- A possible way to solve energy-momentum conservation
- Process governed by the interaction of HQ with QGP around  $T_c => natural link with the energy loss model.$

In EMMI RRTF, comparison between Instant. Coal. & RRM



starting from the same bulk and from the same c spectrum

Significant differences found both for D meson p<sub>⊤</sub> spectrum and  $v_2$ .

R. Rapp et al, arXiv: 1803.03824


#### **HQ** - Recombination

#### Acknowledged:

towards the end of QGP, hadronization of (of equilibrium) HQ can proceed through a dual mechanism:

#### Low $p_T$ :

- The quark partner(s) are already present in the hot cooling medium
- New specific recombination mechanism; no obvious calibration
- The footprint of reconfinment (?!)
- Crucial to explain the flow bump in R<sub>AA</sub>(D) and sizable v<sub>2</sub>(D) => large impact.



#### High $p_T$ :

- The quark partner(s) needed to create the HF-hadron have to be generated from the vacuum
- « usual » fragmentation calibrated on p+p and e++e- data (Petersen,...)

But also energy density dependent (PHSD) !!!

#### <u>Uncertain</u> (and not disputed enough):

- Genuine physical recombination process:
  - Instantaneous Parton Coalescence with local (x,p) correlations (Greco, Ko & Levai 2003), Xor in momentum space (Oh et al 2009): known violation of energy-momentum conservation, advocated to have small effects at finite  $p_{\tau}$
  - Resonance Recombination Model (Ravagli and Rapp, 2009): kinetic c+qbar -> D; spirit of dynamical recombination around  $T_c$  ( $P_{recomb} = \Delta \tau \times \Gamma_{res}(p)$ ; a way to solve the energy-momentum conservation issue
  - In medium Fragmentation (Beraudo et al., 2015): string from HQ + thermal light
- Differences in the « technical implementations » , e.g. normalisation

#### EMMI RRTF: Consequences from various Hadronization Mechanisms


We define and display the H<sub>AA</sub> quantity

$$H_{AA} = rac{rac{dN_D}{dp_T}}{rac{dN_{c ext{ final}}}{dp_T}}$$

...which exhibits at best the specific effects of hadronization :

Significant uncertainties!

=> Yes, one can for sure put more constrains with  $D_s$  and  $\Lambda_c$ , but probably one has also to converge on more robust schemes for « basic » D mesons



#### Consequences from the choice for Bulk, Hadronization and init. Spectrum

#### But without shadowing and hadronic rescatterings

Pb-Pb,  $\sqrt{s_{NN}}$ =2.76 TeV, D meson

0-10% | Model-average |
Data (ALICE) |

MODEL average +  $\sigma$  |

10


12

14

16


 $p_{T}$  (GeV)

R. Rapp et al, arXiv: 1803.03824




- You asked for it, you got it... « Theory error band »
- At least as large as the experimental one!

## Tension between $R_{AA}$ and $v_2$ (at low $p_T$ ): the Catania Cocktail completed



- Probably one of the reasons why some models like EPOS2+MC@sHQ with NOT (const.  $\eta_D$ ) can cope both with  $R_{AA}$  and  $v_2$ .
- What should we do next?




## **HQ Working Group**

S. Cao et al, arXiv: 1809.07894

Collect and compare the transport coefficients from various models:

c-quarks

What is used by various models to fit the data




- Obviously not satisfying: Larger dispersion than the predictions for concrete observables... WHY?
- Because of « extra ingredients », chosen differently in each model (partly) !!!
- More complex then for the case of jets (several FP coefficients)

## **HQ-Working Group (convener: X-N Wang)**

S. Cao et al, arXiv: 1809.07894

- The goal is to: Collect and compare the transport coefficients from various models,
  - Measure and understand their consequences by first studying a simpler brick problem
  - Estimate some systematics + uncertainties

Best controlled QGP ever: uniform fixed temperature for all models (with same initial condition FONLL-like @ RHIC)



<u>Main result</u>: Nice structuration of the transport coefficients in different classes. For each class, the work illustrates the maximal accuracy reachable once all other ingredients are either fixed or chosen commonly

## **HQ-Working Group (convener: X-N Wang)**

S. Cao et al, arXiv: 1809.07894

49

- The goal is to: Collect and compare the transport coefficients from various models,
  - Measure and understand their consequences by first studying a simpler brick problem
  - Estimate some systematics + uncertainties


Best controlled QGP ever: uniform fixed temperature for all models (with same initial condition FONLL-like @ RHIC)

1) Rescale the coefficients to match  $R_{AA}$ =0.3 at p=15 GeV & « final time » 3 fm/c 2) Compare them! Duke T=250 MeV Duke LBL-CCNU LBL-CCNU LBL-CCNU T = 250 MeV t = 3 fmCatania QPM Catania QPM Catania QPM A (GeV / fm) [elastic part] 1.2 Catania pQCD Catania pQCD Catania pQCD (tune 2) TAMUFrankfurt PHSD Frankfurt PHSD Frankfurt PHSD 0.9 Nantes col.+rad. Nantes col.+rad. Larger discrepancies Nantes col.+rad. Nantes col. Nantes col. Nantes col pQCD & 0.6 0.3 El + rad. 0.0 20 p<sub>T</sub> (GeV) 10 10 20 p (GeV) p (GeV)


<u>Going beyond</u>: A) radiative energy loss for everyone B) thermal mass of light partons for everyone C) consider alternate observables affording to probe FP coefficients from different perspective D) adopt common ingredients (like common bulk) and (weaker) exchange ingredients between models.


## Alternate observable: HF-HF correlations

- Back to back D/Dbar or B/Bbar: As compared to γ-D/B: "trigger" itself is affected but symmetry between both particles limitates the various effects.
- ► Elastic Eloss vs radiative Eloss: The purely collisional scatterings lead to a larger average <p\_²> then the radiative "corrections" (need for large scattering to be efficient)... although both types can give correct agreement with the data at intermediate p<sub>T</sub>.



Expected consequences for azimuthal correlations (probe of B<sub>T</sub>: good: complimentary to usual RAA and v2)







Tuned to reproduce the R<sub>AA</sub>

## Alternate observable: HF-HF correlations

Assumption of back 2 back emission of initial QQbar (naïve LO...)

M Nahrgang et al, PRC90 (2014), 1305.3823







- $\triangleright$  Indeed, rather large differences found for both b and c, and all kind of p<sub>T</sub> cuts (... but good to see there is an effect though,...)
- For the smallest  $p_T$  bin and elastic energy loss, we even find an inversion of the correlation ("hot partonic wind" push;  $v_0$  bulk =>  $v_1$  correl; underlying event)


## Alternate observable: azimuthal correlations


...but higher orders can have a significant impact:





> LO; (a): back to back peak







➤ NLO;



(d): "flavor excitation": no strong azimuthal correlation expected






(e): gluon splitting: strong peak around  $\Delta \phi = 0$ 

(f): higher order FE; both Q and Qbar in the "remnant" region


### Alternate observable: azimuthal correlations


- NLO effect simulated with MC@NLO + HERWIG (parton shower)
- ► Gluon splitting processes lead to an initial enhancement of the correlations around  $\Delta \phi$ =0; Strong broadening of the  $\Delta \phi$ = $\pi$  peak ("vacuum" radiation is dominant)
- ➤ For intermediate p<sub>T</sub>: increase of the variances due to Eloss from 0.43 (initial NLO) to 0.51 (+20%) for the purely elastic mechanisms and to 0.47 (+10%) for the interaction including radiative corrections.
- Correlations at large p<sub>T</sub> seem to be dominated by the initial correlations. Nothing will be learned on the Eloss mechanisms in this region
- Different NLO+parton shower approaches agree on bottom quark production, differences remain for charm quark production
- Confirmation by other groups (Duke, CCNU-LBL,...)



### Alternate observable: azimuthal correlations

- NLO effect simulated with MC@NLO + HERWIG (parton shower)
- ► Gluon splitting processes lead to an initial enhancement of the correlations around  $\Delta \phi$ =0; Strong broadening of the  $\Delta \phi$ = $\pi$  peak ("vacuum" radiation is dominant)
- For intermediate p<sub>T</sub>: increase of the variances due to Eloss from 0.43 (initial NLO) to 0.51 (+20%) for the pt mechanisms and to 0.47 (+10%) for the interact radiative corrections.
- Correlations at large  $p_T$  seem to be dominated I 0.100 correlations. Nothing will be learned on the Elomechanisms in this region
- Different NLO+parton shower approaches agree quark production, differences remain for charm production
- Confirmation by other groups (Duke, CCNU-LBL







## Alternate observable: momentum imbalance

 $> \gamma - D/B/c$  jet /b jet:

In QGP: Longitudinal fluctuations of the HQ energy loss crucially depend on the precise mechanism and cannot be measured easily in usual observables like  $R_{AA}$  or  $v_2$ 



No E loss => perfect probe of initial  $\vec{P}_{HQ}$ 




differential probability to loose energy  $\boldsymbol{\omega}$  per unit time


➤ Of course: NLO effect in the production mechanisms makes it not so trivial (not to speak about exp. Issues... RUN3 ? RUN4 ?)

## Alternate observable: momentum imbalance

- Back to back D/Dbar or B/Bbar: As compared to γ-D/B: "triggering" itself is affected but symmetry between both particles could limitate the various effects:
- Large number of c-cbar from various NN collisions => large uncorrelated background
- Competing effects due to energy loss: ...







- decorrelation due to various path lengths + fluctuations: reduction
- Of course, needs for significant correlations in pp, which is still not quite settled

L. Vermunt et al. arXiv:1710.09639

## New Observables are coming

## Short term, mid-term, long term,...


What	Good for ?	Caviat
Event shape engineering	Strength and T dependence of the interaction	Might be sensitive to the bulk and initial stage => play collective
Heavy light - correlations	b/c-jet substruture, nature of the interaction	Might be sensitive to various HF creation in pp, to be calibrated
$\Lambda_c$ , $D_s$ , $B_s$ ,	Understanding hadronization esp. Recombination (if generic enough not to require 1 new free parameter per state) or limits of statistical models	Dynamical treatment of confinement ? Inputs from IQCD probably needed
v <sub>1</sub> (y)	Constrain (E,B), vorticity, initial tilt of matter initial distribution of HQ in transverse plane	Isn't it a bitt too much for this poor observable?

#### ...but still a need to better understand old ones!

The nature of the interaction (Elastic vs Elastic + Radiative vs AdS/CFT) in the intermediate p<sub>T</sub> range

One should exploit both

- Analysis of the path length dependence
- The larger "collinearity" found in radiative collisions, which could be seen in some structure of the v<sub>n</sub> (p<sub>T</sub>), to be better understood and studied jointly for B and D




## Adopting common baseline or exchanging some ingredients

Pioneering work done done by Nantes and Tamu (P. B. Gossiaux et al., arXiv:1102.1114)

Recent study made by EMMI RRTF using VISHNU 2D+1 viscous hydro as a common background

and LANGEVIN dynamics for HQ

R. Rapp et al, arXiv: 1803.03824





## Adopting common baseline or exchanging some ingredients

 Pioneering work done done by Nantes and Tamu / Recent study made by EMMI RRTF...

Y. Xu et al, Physical Review C99 (2018) arXiv: 1803. 10734 Catania, Duke, Frankfurt & Nantes

- Even more recent: 6 HQ Eloss models in the same VISHNU viscous 2D+1 hydro through Langevin dynamics ...Confirms the dispersion of results, now after full implementation in bulk evolution
- Others "goodies": impact of initial conditions
   & impact of Einstein relation.
- Nice work, BUT....





## Adopting common baseline or exchanging some ingredients

#### .... Some methodological issues:


- Those comparison can "only" be done for global models relying on external T and velocity fields
  (provided f.i. by fluid dynamics) while there are a lot of valuable transport models dealing with HF
  + soft evolution, like PHSD, CATANIA,... (they can always provide such fields by averaging on
  several events, but not clear it is completely relevant)
- For the time all these comparisons where performed using Langevin forces as a substitute to Boltzmann dynamics (computationally less demanding), although Y. Xu et al emit strong doubts whether it is a good way to proceed.
- In some case: comparison of transport coefficient used both in pure elastic processes and in elastic + radiative processes: can lead to some confusion.

## Need to think deeper

## Various approaches to transport


#### Bottom-up schemes (microscopic -> mesoscopic):

- Assume (effective) degrees of freedom and (effective) interactions
- Take insights and constrains from the fundamental QCD theory, but often inholds some free parameter
- Rely on more or less sophisticated realizations of the transport theory




#### <u>Langevin from Einstein relation (Fluctuation Dissipation Theorem) view point:</u>

- In general, not possible to accommodate all fluctuations as well as the asymptotic distribution (at a fixed T)
- Requires to impose some Einstein –
  relation / fluctuation dissipation
  theorem between the drag
  coefficient and the diffusion
  coefficients. Several prescriptions
  have been tested and lead to various
  results:
- Lesson 1: avoid resorting to Langevin just to spare some tedious programming: Going for the microscopic process is more faithfull to physics.
- My personal opinion (from comparison to Boltzmann benchmark): The most conservative way to implement FDT is to preserve  $\eta_{\text{D}}$  and  $\kappa_{\text{T}}$  and to adapt  $\kappa_{\text{L}}.$



#### Langevin from Boltzmann view point:


• Lesson 2: For coarse grained observables like the  $R_{AA}$  and the  $v_2$ , the agreement between the 2 transport schemes essentially depends on the isotropization strength of the cross section (i.e., the Debye mass of the gluon propagator)



For  $m_D = g T \approx 2 T$  found f.i. in the Quasi Particle Model, extra coupling is found for the  $R_{AA}$  using LV, which can be suppressed by reducing the FP coefficients by  $\approx 30 \%$ 

#### Langevin from Boltzman view point:


 For « exclusive process », momentum distributions differ significantly, even after imposing Einstein relation (ER):



These differences should me seen in observables like g-HQ correlations


#### **Boltzmann from Langevin view point:**

- There are a lot of situations where Langevin dynamics applies, but not Boltzmann, thanks to the large mass of the particles.
- It is even a result proven for dynamical systems (conditions on the velocity applies as well)
- In a dense strongly coupled system, this is likely to be the case!




Gluon spectral functions (Liu & Rapp 2016)


## Visual summary






## HF in small systems: initial, final (collectivity) or both?





 $CP(\Delta\Phi)$ 



Finite v<sub>2</sub> seen in p-Pb data but no significant difference in the spectrum?


It seems to me that we get similar modifications from both the 0<sup>th</sup> and 2<sup>nd</sup> order harmonic (to be checked)
 (b) D: √s = 5.02 TeV, -1.0 < y<sub>h</sub>, f<sub>i</sub> < 0.0</li>

Physical origin ?

• Both coupling to hydro and CGC lead to azimuthal correl. for HF in p-Pb.4


Duke: Y. Xu, Nucl.Part.Phys.Proc. 276-278 (2016) 225 POWLANG: A. Beraudo et al: JHEP 1603 (2016) 123

H. Fujii, K. Watanabe, Nucl.Phys. A920 (2013) 78



both  $v_2 \approx 4\%$ 

## Recent study using EPOS-HQ (Hard Probes)




#### Sensitivity

Saturation	Hadronisation	E Loss
YES	YES	YES

... and chemistry! Complicated!!!

Our implementation of the coalescence has consequences up to 10 GeV

#### **Conclusions**



- Existing models offer the possibility to describe most of the OHF experimental AA data while being compatible with existing theory constrains...
- ... however with unequal precision and no consensus on the physical NP content
- Improvements and quantitative understanding is on their way, but it will still take some time and a lot of efforts => need for ressources, bright (young) people and collective work.
- Open Heavy Flavors are maybe not an ideal probe of QGP yet, but they are quite fascinating and offer bright future for the field, with multiple interconnections.

## Back UP

**IQCD** 

# IQCD Calculation of D<sub>s</sub>

- Lattice QCD at finite T is performed in Euclidean space notoriously difficult to calculate dynamical quantities.
- Up to 2014, Ds was evaluated directly through the (narrow) diffusion peak of the spectral function evaluated from current – current correlator (hard)
- From 2014: Use of the field field correlator in order to obtain a better shaped spectral function:

$$G_{\rm E}(\tau) \equiv -\frac{1}{3} \sum_{i=1}^{3} \frac{\left\langle \operatorname{Re} \operatorname{Tr} \left[ U(\beta; \tau) g E_{i}(\tau, \mathbf{0}) U(\tau; 0) g E_{i}(0, \mathbf{0}) \right] \right\rangle}{\left\langle \operatorname{Re} \operatorname{Tr} \left[ U(\beta; 0) \right] \right\rangle}$$


• Then obtain the variance  $\kappa$  of stochastic forces (a transport coefficient;  $\kappa$  = 2 x B) from the slope of spectral function  $\rho_E$  at  $\omega$  = 0:

$$\kappa \equiv \lim_{\omega \to 0} \frac{2T \rho_{\rm E}(\omega)}{\omega} \quad \text{with } \rho_{\rm E} \, \text{extracted from} \quad G_{\rm E}(\tau) = \int_0^\infty \! \frac{\mathrm{d}\omega}{\pi} \, \rho_{\rm E}(\omega) \, \frac{\cosh[\omega(\frac{\beta}{2} - \tau)]}{\sinh[\frac{\omega\beta}{2}]}$$

• Main result :  $\kappa/T^3=1.8\dots 3.4$  then convert to D<sub>s</sub>

# IQCD Calculation of D<sub>s</sub>

• This leads to  $2\pi TD_s=3.7\dots6.9$ 



- Still drastic approximations/limitations: quenched QCD, heavy quark vs. charm quark, (no) continuum extrapolation,...
- Relaxation time  $\eta^{-1}=rac{m_Q D_s}{T}=rac{0.59\dots 1.1\,m_Q}{T^2}pprox 2.5\dots 5\,{
  m fm/c}$

High pT

### Status of high p<sub>T</sub> HQ: prospects

#### Other challenges:

- Better understanding of heavy mass effect and medium properties in the radiation (especially on the coherence effects)
- Embedding in a realistic medium (for some models)
- In a « jetty » picture: Combination of induced Eloss affecting the « initial » DGLAP evolution and the final « on shell » HQ


"Drag Induced Radiation and Multi-Stage Effects in Heavy-Flavor Energy Loss"

S. Cao et al, arXiv:1711.09053

S. Cao, QM 2018


Also poster G-Y Qin

#### Hard scale Q=p<sub>T</sub>



#### Low scale $Q_0 = m_0$

« Usual » time ordered LBL-CCNU evol. (Higher twist with  $\hat{e}$  and  $\hat{e}_2$  corrections )



Good agreement with CMS data for D and B, some influence of higher order term at intermediate  $p_T$ 

## **HQ-WG**

### **Heavy-Quark Working Group**



After 3 meetings, footprints of the physics start to emerge... but no firm conclusion yet



For step 1: Compare HQ spectra from different models in static medium with common initial condition

Models	note	basic	tune 1
LBL-CCNU ★	fix $\alpha_s$	$\alpha_s = 0.3$	$\alpha_s = 0.26$
Duke	fix $\alpha_s$	$\alpha_s = 0.3$	$\alpha_s = 0.23$
Catania QPM	$\alpha_s(T)$	K = 1	K = 2
Catania pQCD	$\alpha_s(T)$	K = 1	K = 3.4
TAMU	U-potential	no tuning	no tuning
Frankfurt PHSD	$\alpha_s(T)$	no tuning	no tuning
Nantes col. + rad.	$lpha_s(q^2)$	K = 1	K = 0.8
Nantes col. only	$lpha_s(q^2)$	K = 1	K = 1.5

\*: Radiative included

Basic: original model

Tune 1: favorite tuning of each group in order to describe *D* meson data with their own ingredients (background, hadronization,...); K = rate multiplyer


## **HQ-Working Group (convener: X-N Wang)**


- The goal is to: Collect and compare the transport coefficients from various models,
  - Measure and understand their consequences by first studying a simpler brick problem
  - Estimate some systematics + uncertainties

Best controlled QGP ever: uniform fixed temperature for all models (with same initial condition FONLL-like @ RHIC)

1) Rescale the coefficients to match  $R_{AA}$ =0.3 at p=15 GeV & « final time » 3 fm/c

2) Compare them!





<u>Main result</u>: Nice structuration of the transport coefficients in different classes. For each class, the work illustrates the maximal accuracy reachable for each class once all other ingredients are either fixed or chosen commonly

### **EMMI - RRTF**

### **EMMI RRTF**

#### Thanks to the generosity of EMMI!!!

1<sup>st</sup> meeting: 18-22 July 2016; GSI (Germany) 2<sup>nd</sup> meeting: 12-14 Dec 2016; GSI (Germany)

#### Researchers involved:

- Organizers
- HQ-WG members
- Other key players in HF QGP tomography (A. BERAUDO, M. DJORDJEVIC, C. GREINER, G. INGHIRAMI, H. VAN HEES, I. VITEV, CUEJET...)
- IQCD experts (O. KACZMAREK, P. PETRECZKY)
- QCD and EFT experts (G. MOORE , J. PAWLOWSKI)
- Selected experimentalists (J. BIELCIK, P. BRAUN MUNZINGER, E. BRUNA, Z CONESSA, A. DAINESE, YJ LEE, F. PRINO, J. STACHEL)

### **EMMI RRTF**


Goal to attack the problem with a broad view right from the beginning...


#### Topics:

- Initial spectra and shadowing
- Bulk evolution and consequence on HF observables
- Transport implementation
- Hadronization
- Microscopic models for HF energy loss and constrains from QCD at low and high momentum
- Future observables

Selected topics presented here

Collection of models vs data by Yen-Jie Lee:






Some « outliers »... Is it acceptable or does it has measurable consequence either on the « extraction » of the transport coefficient (for « tunable » models) or on the agreement with experiment.

« We all do FONLL / GM - VFNS » ... yes, but with slightly different parameters !

Right now: data better than uncertainty band in theory:


3 best fits extrated from members of the ALICE collaboration, with a fair wish to explore various hardness (BASELINE)



Consequence of the baseline on the RAA of D mesons in the Nantes Model (thermalisation at small  $p_T =>$  the initial profile has a large impact in the ratio)

Consequence for the collective chase: ideally, all models should adopt the baseline; minimal action: check that the  $\chi^2$  with their own production model is at least as small as the one found in the baseline (rejection if  $\chi^2/NDF > 2$ ).

- RAA for c quarks with MNR+EPS09NLO
- Proposal: multiply the input c-quark  $p_T$  spectrum from FONLL by this RAA and use the band to define a band on final  $R_{AA}$  and  $v_2$




#### Consequence for the collective chase:


- ideally, all models should adopt the same prescription for shadowing, and the uncertainties on the shadowing should then be recast in a global systematic error on the D<sub>s</sub>, common to all models
- If some models have an intrinsic theory to evaluate shadowing and are unwilling to modify this for the sake of consistency (f.i. EPOS3), the minimal « quality control » should be to implement the common prescription for shadowing and display the consequences on the observables and on their extraction of the transport parameters in order to document the origin of possible differences.
- Perspective (apart from hoping on better control on shadowing):
  - Go for B or to v<sub>2</sub>, less sensitive to shadowing
  - Uncertainties on the shadowing may partly factor out if the ratio of R<sub>AA</sub> at different centralities is considered

### **EMMI RRTF: the bulkS**


Model	$dN_{\pi^+}/dy \ (dS/d\eta)$		$dN_p/dy$	
	0-10%	30-50%	0-10%	30-50%
UrQMD	495	152	34	11
TAMU	682 (12400)	170 (3080)	58	15
Nantes	478	129	38	10
Catania	(14000)	(3700)		
LBL-CCNU/Duke	653 (12600)	160 (3080)		
CUJET	610 (10820)	142 (2610)	45	11
POWLANG	(9100)	(1450)		
PHSD	722	148	31	6
exp.	$670 \pm 68$	$163 {\pm} 15$	31±4	8±1

Table 1: Inclusive  $\pi^+$  and proton numbers (*i.e.*, including strong and electromagnetic feeddown) per unit momentum-space rapidity in Pb-Pb(2.76 TeV) collisions in the various bulk evolution models. Also shown in parentheses are the values for the total entropy per unit space-time rapidity at the end of the QGP phase (as available). As a reference the last row shows experimental values from Ref. [51].






### EMMI RRTF: the average vs the various models



### EMMI RRTF: TAMU pQCD x 10




## Hadronization

### **HQ** - Recombination


### **In-Medium recombination:**

A Beraudo et al., Eur. Phys. J. C75 no. 3, (2015) 121



- String formed at Tc from HQ and light quark sampled out of thermal bath
- Novel mechanism which allows a natural transition from the low p<sub>T</sub> to the high p<sub>T</sub>
- Also leads to large flow bumps and extra v<sub>2</sub> contribution

# Hadronisation (Ds)




ALICE data for D and Ds mesons (JHEP 1603 (2016) 082) compared with TAMU-model predictions (M- He et al., PLB 735 (2014) 445)

## Other constrains

# Model + Exp. Constrains on D<sub>s</sub>

F.i., recent STAR study:

#### **STAR Collab. PRL 118 (2017)**



compare with	$2\pi T D_s$	$\chi^2/\text{NDF}$	<i>p</i> -value
SUBATECH [11]	2-4	15.2 / 8	0.06
TAMU c quark diff. [13]	5 - 12	10.0 / 8	0.26
TAMU no c quark diff. [13]	-	29.5 / 8	$2 \times 10^{-4}$
Duke [14]	7	35.7 / 8	$2 \times 10^{-5}$
LBT [15]	3-6	11.1 / 8	0.19
PHSD [10]	5 - 12	8.7 / 7	0.28
3D viscous hydro [42]	-	3.6 / 6	0.73

TABLE I.  $D^0$   $v_2$  in 0–80% centrality Au+Au collisions compared with model calculations, quantified by  $\chi^2/\text{NDF}$  and the p-value (the probability of observing a  $\chi^2$  that exceeds the current measured  $\chi^2$  by chance).  $2\pi T D_s$  values quoted are in the range of  $T_c$  to  $2T_c$ .  $\chi^2/\text{NDF}$  is calculated in the  $p_T$  range wherever the model calculation is available.


- Not really conclusive;  $\chi^2/NDF$  is not a smooth function of  $D_s$ : large residuals
- Beware: Models are essentially validated at finite p<sub>(T)</sub>; extrapolation at 0 momentum might contain further uncertainties
- Urgent need for collective actions to better control the « residuals »

## Momentum Imbalance


## $\gamma$ - b/c jet: Best HF Correlation ever ?

 $> \gamma - D/B/c$  jet /b jet:

In QGP: Longitudinal and transverse (qhat) fluctuations of the HQ, which crucially depend on the Eloss mechanism and cannot be measured in usual observables like  $R_{AA}$  or  $v_2$ 



No E loss => perfect probe of initial  $ec{P}_{HQ}$ 





differential probability to loose energy  $\omega$  per unit time

➤ Of course: NLO effect in the production mechanisms makes it not so trivial (not to speak about exp. Issues... RUN3 ? RUN4 ?)

# Next best thing: HF-HF correlations

- $\triangleright$  Back to back D/Dbar or B/Bbar: As compared to  $\gamma$ -D/B: "triggering" itself is affected but symmetry between both particles could limitate the various effects:
- Large number of c-cbar from various NN collisions => large uncorrelated background
- Competing effects due to energy loss: ...






 $\triangleright$  Goal of the study: investigate whether  $p_T$ - $p'_T$  correlations survive NLO effects

L. Vermunt et al. arXiv:1710.09639

Method for "systematics": use 2 event generators: PYTHIA (6.4) & EPOS3




 In pythia, those topologies are generated by coupling LO processes (implying 0,1 or 2 HQ) and ISR + FSR ... This will be referred to as « LO + NLO ccbar » (strictly speaking, no NLO!)

 $\triangleright$  Goal of the study: investigate whether  $p_T$ - $p'_T$  correlations survive NLO effects

L. Vermunt et al. arXiv:1710.09639

Method for "systematics": use 2 event generators: PYTHIA (6.4) & EPOS3




- In pythia, those topologies are generated by coupling LO processes (implying 0,1 or 2 HQ) and ISR + FSR ... This will be referred to as « LO + NLO ccbar » (strictly speaking, no NLO!)
- Same « strategy » in EPOS3, with « semi-hard pomeron » approach (with some soft evolution included), with various LO Born processes.

 $\triangleright$  Goal of the study: investigate whether  $p_T$ - $p'_T$  correlations survive NLO effects


L. Vermunt et al. arXiv:1710.09639

Method for "systematics": use 2 event generators: PYTHIA (6.4) & EPOS3




- In pythia, those topologies are generated by coupling LO processes (implying 0,1 or 2 HQ) and ISR + FSR ... This will be referred to as « LO + NLO ccbar » (strictly speaking, no NLO!)
- Same « strategy » in EPOS3, with « semi-hard pomeron » approach (with some soft evolution included), with various LO Born processes.
- In pythia, possibility to restrict to LO ccbar production processes with massive elements (MSEL=1 -> MSEL=4 flag), still switching on the ...
   ISR + FSR ... This will be referred to as « LO ccbar »

Including NLO effects in the charm production (N.B. :beauty would be better for our purpose, but very low statistics)



due to flavor excitation like process

 $\underline{\text{N.B.}}$ : Pythia MSEL=4: at least 1 ccbar pair un each event => Normalized according to high-p<sub>T</sub> LO charm creation in Pythia MSEL=1



- Different p<sub>T</sub> imbalance for 3 production models in pp
- 2 of them show that NLO effects does not completely destroy the perfect correlation found in LO production
- Similar results for DDbar


#### Abolute correlation:

$$C(p_T, p_T') := \frac{1}{N} \frac{d^2 N(p_T, p_T')}{dp_T dp_T'} - \frac{1}{N} \frac{dN(p_T)}{dp_T} \times \frac{1}{N} \frac{dN(p_T')}{dp_T'}$$

- Vanishes if  $d^2N$  factorizes  $(d^2N(p,p') = dN(p) \times dN(p'))$
- Satisfies  $\int C(p_T, p_T') dp_T d_T' = 0$

Relative correlation: 
$$C_R(p_T,p_T'):=rac{C(p_T,p_T')}{rac{1}{N}rac{dN(p_T)}{dp_T} imesrac{1}{N}rac{dN(p_T')}{dp_T'}}$$

■ Reveals correlation at finite p<sub>T</sub>



# Guidelines & strategies

### Guide lines

### Collectivity of the « HQ in URHIC » community

- As theorists, we should maintain and amplify the recent collective efforts in order to
  - Provide more systematic systematical errors from our models as well as "figures of merit" like the Catania cocktail
  - Adopt a reasonable base line for the "extra ingredients" which will make the role of the "core ingredients" more transparent" => possible new "structures" emerging as in the HQ-WG brick study

Good exemple in EMMI RRTF: baseline for c-quark initial spectrum provided by some members of ALICE

 We should as well rely more intensively and systematically on modern methods like Bayesian analysis, which allow a better quantitative contact with the experimental community

### Mid-term strategies


### But also, from the theory view point:

• Get more insights and constrains from IQCD (transport coefficient at finite p?) See discussions f.i. in

G. Aarts et al, Eur. Phys. J.A 53, no. 5, 93 (2017)

(proc. Lorentz Workshop)

 and make contact with these in the models. In this respect, charm correlations and charm meson correlators -- which indicate the presence of mesonic dof around Tc -- should be evaluated in the existing and future models and compared with the ICD results



S. Y.F. Liu and R. Rapp, Phys. Rev. C 97, 034918 (2018)

S. Y.F. Liu, QM2018

 Develop models which span the low p – intermediate p – large p & low T – large T range and contain the proper physics in each region

### Mid-term strategies

### But also, from the theory view point:

- Develop transport theories beyond the classical approximation (and don't be afraid by numerical work, even if redious)
- Assess the role of QGP non-equilibrium effects on HF (including in the initial stage).
- Achieve a rigorous implementation of radiative energy loss in realistic bulks.
- Strutinize the consequences of other effects usually discarded
  - Evolution in turbulent plasma Stanislaw Mrowczynski, Eur.Phys.J. A54 (2018) no.3, 43
  - Fluctuation of chromoelectric field

QM Poster A. I. Sheikh


• ...


Any suggestion welcome!

# Small systems




# HF in small systems: initial, final (collectivity) or both?





 $CP(\Delta\Phi)$ 



Finite v<sub>2</sub> seen in p-Pb data but no significant difference in the spectrum?

It seems to me that we get similar modifications from both the 0th and 2nd order harmonic (to be (b) D:  $\sqrt{s} = 5.02 \text{ TeV}, -1.0 < y_{h}$   $f_b < 0.0$ 


checked)

Physical origin?

Both coupling to hydro and CGC lead to azimuthal correl. for HF in p-Pb.4

Duke: Y. Xu, Nucl.Part.Phys.Proc. 276-278 (2016) 225 POWLANG: A. Beraudo et al: JHEP 1603 (2016) 123

H. Fujii, K. Watanabe, Nucl. Phys. A920 (2013) 78



both  $v_2 \approx 4\%$ 

## HF in small systems: initial, final (collectivity) or both?

### Some thoughts:


- Very difficult topic, even in the light sector
- What does m<sub>Q</sub> bring to the problem?

favor CGC	disfavor CGC	Pro hydro	disfavor hydro
Effect deeply rooted in the production mechanism	?	Other indications of collectivity in p-Pb	Theoretical status of hydro in p- Pb unclear (see M. Strickland's talk)
K Watanabe: « From the CGC view point, heavy mesons should be more correlated than light hadrons" (reinteraction less efficient).			Recombination in small systems not under control (large differences due to technical implementation)
			Absolute value found up to now too low: $v2(B) < v2(D) < v2(\pi)$

interesting.

M. Greif et al, Phys. Rev. D 96, 091504 (2017)


## HF: from p-Pb -> Pb-Pb



Gradual transition from CGC -> geometry?


# p-Pb (Duke)

 arXiv:1510.07520v2: viscous hydrodynamics model (vHLLE) + improved Langevin + mixed hadronization




$$rac{dec{p}}{dt} = -\eta_D(p)ec{p} + ec{\xi} + ec{f}_g \qquad D = 4rac{T^2}{\hat{q}}rac{C_A}{C_F}$$
 D  $\hat{q}$ 

- $D(2\pi T) = 5$  tuned from central PbPb
- Differential pT (in Hard probes talk) for D=5 saturates at 2%: incomplete coupling with QGP (short lifetime)




# p-Pb (Beraudo et al)

Arxiv 1512.05186v2: essential role of initial flutuations (main source of anosotropy):
 MC Glauber + av. init medium cond after psi\_2 rotation + PYTHIA + EPS09
 shadowing + kt broadening + viscous hydrodynamics (ECHO) + POWLANG + string fragmentation with the light partner taken from finite T medium



Langevin + 2 different Eloss models: HTL (pQCD) & IQCD

Major contribution to v2(D): hadronisation



Some similarity with ALICE mass ordering, (no substraction in Beraudo et al as single part. analysis)

