Chromoelectric Field Fluctuations and Heavy Quarks in Hot and Dense QCD Medium

MUNSHI G. MUSTAFA



Saha Institute of Nuclear Physics, Kolkata

Content:

- Why Heavy Quark?
- Heavy Quark Production
- Heavy Quark Propagation
 - Collisional Energy Loss
 - Radiative Energy Loss
 - **▶** Heavy-flavoured hadron spectra (Jet Quenching)
- Chromoelectric Field Fluctuations
 - Collisional Energy gain
 - Heavy-flavoured Nuclear Suppression factor
 - \rightarrow J/ψ Nuclear Suppression factor
- Conclusion


Why Heavy Quarks?

- Heavy Quarks ➤ Charm (c) & Bottom (b)
 - **Charm:** $M_c = 1.27^{+0.07}_{-0.09}$ **GeV**
 - **Bottom:** $M_b = 4.19^{+0.18}_{-0.06}$ **GeV**
- **■** Production time: $\tau_Q = 1/2M_Q \le 0.1 \text{ fm}/c$
- **■** Thermalisation time (τ_{th}) :
 - Successive equlibrium: $au_{th}^g < au_{th}^{u,d} < au_{th}^c < au_{th}^b < au_{th}^{life}$ [J. Alam, S. Raha & B. Sinha, PRL 73 (1994) 1895]
 - Therma. time for HQs $au_{th}^Q \sim rac{M_Q}{T} imes au_{th}^{u,d}$ [G. Moore & D. Teaney, PRC 2005]
 - $extstyle au_{th}^{u,d} < au_{th}^Q$: no production at QGP and hadronic phase
 - Produced at very early time interactions in Hard Scattering of partons in Nucleons. Initial distribution of HQs ➤ frozen
 - $lue{ }$ Since $g,\,u,\,d$ thermalise early and provide an expanding thermal background

Why Heavy Quarks?

- Heavy Quarks propagation:
 - propagates in foreground with gluons & light quarks as an expanding thermal background for the non-equilibrated HQs ➤ expansion dynamics
 - interacts with equilibrated degrees of freedom > energy-loss of HQ
 - ightharpoonup requires dynamics ightharpoonup final distrⁿ of HQs (Transport Eq.)
- Uniqueness of Heavy Quarks:
 - distinguishable down to lowest momenta > medium coupling (Hydro)
 - lacktriangle cleaner energy-loss probe lacktriangle reflected in leading particle p_\perp spectra, flow
 - tests understanding of mass dependence
 - Quarkonia: QGP thermometer

Heavy Quark Production in Heavy-Ion Collisions:




- Heavy-lon Coll.: ALICE: 0809.1062[nucl-ex]
 - **Charm:** LHC $\sim 10 \times$ RHIC **Bottom:** LHC $\sim 100 \times$ RHIC
- lacksquare Produced in pairs (Qar Q) at early times
- QCD provides a framework to compute production x-section and distribution

HQs Production and Initial Distribution

Factorisations allows that an observable can be expressed as a convolution of short distance (hard scattering of partons) and the long distance contribution describing the initial hadrons


$$\frac{d\sigma}{dy_1 \, dy_2 \, dp_{\perp}} = x_1 x_2 p_{\perp} \sum_{ij} f_i^{(1)}(x_1, \mu_F^2) f_j^{(2)}(x_2, \mu_F^2) \hat{\sigma}_{ij}(x_i, x_j, \alpha_s(\mu_F^2), \mu_F^2)$$




- \bullet i, j are partons in hadrons
- $lue{} \sigma_{ij}$ are partonic x-sections $lue{}$ pQCD
- f_i are parton distrⁿ fn. (PDFs) in hadron which are nonperturbative
- $lue{}$ μ_F is the factorization scale between hard processes and the non-perturbative PDFs
- $lue{}$ μ_F dependence should cancel in order by order in pQCD calculation

HQs Production and Initial Distribution

- Factorisations:
 - $igoplus gg o Qar Q; \quad qar q o Qar Q; \quad gg o Qar Qg; \quad qar q o Qar Qg \,\,\, ext{[Leading Order (LO), NLO \& FONLL]}$





- The short distance contribution describing the matrix elements of hard processes are free from mass singularities and calculable in pQCD
- PDFs obey DGLAP evolution equation ➤ knowing PDFS at one scale can be calculated at any other scale
- In elimentary [pp and $p\bar{p}$] collisions an observable is well controlled through factorization

Heavy-Ion Collisions & Uncertainties

- Does factorisation work in A+A? May or may not! But assumed!
- PDFs in nuclei (nPDFs) are different from those in hadrons: geometry, wavefunction are different, nPDFs require appropriate modifications
- Several nPDFS that prametrise the nuclear dependence of incoming parton distribution functions
- "uncertainity in intial state interactions"

- The frgmentation function, used to obtain p_{\perp} spectra, should also be afftected by the presence of hot and dense QCD medium!
- "uncertainity in final state interactions"

Heavy Quark Propagation and Final distribution

Produced at early time $(\tau_Q < \tau_{th})$; No production at later time

- Total no. of HQ gets frozen very early in the history of collisions
- Immediately upon their production they will propagate through QGP
- One is left with the task of determining the HQ distribution
- Details of the distribution may reflect the characteristics and development of QGP

Fokker-Planck Equation

Dynamics of a heavy quark propagation as a foreground particle in a QGP can be studied using the simplified Boltzmann approach when the background bulk medium is in thermal equilibrium




Extensive Body of Literatures

- Chakraborty & Syam, Lett. Nuvo Cim.'84
- ► Svetitsky, PRD'88
- Alam, Raha & Sinha, PRL'94
- ▶ Mustafa, Pal & Srivastava, PRC'97
- ► Mustafa PRC'05
- ▶ Moore & Teany PRC'05; Rapp &Hees'05

Fokker-Planck Equation


- Fokker-Planck Eq. ➤ Simplified Boltzmann Eq.
 - Boltzmann Eq ➤ No external force and isotropic in space
 - Soft Scatt. ➤ Taylor expan. of Coll. Integral ➤ Landau Eq.
 - Landau Eq. ➤ an integro-diff^l Eq. involving transport coeffs. ➤ describes collision processes of two particles
 - Landau Eq. ➤ FP Eq. when distribution of the background particles (QGP) is thermal whereas foreground particle (HQs) is non-thermal




Propagation of Heavy Quark in QGP

- Energy-Loss
 - Collisional Loss


Radiative Loss






Collisional Processes


Feynman Diagram



Heavy Quark Transport Coefficients from FP Equation

lacksquare Drag $\gamma(p)$ and Diffusions $\mathcal{B}_0(p)$






- Mustafa, Pal & Srivastava, PRC'97
- Mustafa PRC'05


Differential Collisional Energy Loss

$$ho$$
 Drag Coeff.: $\gamma(p) = -\frac{1}{p} \frac{\mathsf{dE}}{\mathsf{dx}}$

Tilder Differnetial Collisional Energy-Loss: $\frac{dE}{dx}$


[Mustafa, Pal, Srivastava, PRC57 (1998) 889]





HQ Energy-Loss Distribution [Mustafa, PRC72 (2005) 014905]

Collisional Energy-Loss > Transport-coeffs. > Fokker-Planck Eq. > Energy-Loss Distribution





- Mom. diffusion begins
 - Peak Shifts to lower p >
 Energy Loss
 - Peak broadens > Momentum
 Dispersion


Heavy Quarks Dynamics

Substantial Collisional E-Loss

Mustafa, PRC72 (2005) 014905








- Coll. E-Loss is important
- lacksquare Predicted R_{AA} but no data then

- lacksquare For light quarks R_{AA} :
- Mustafa et al., PRL100 (2008) 072301; Acta Phys.Hung. A22 (2005)

Radiative Energy Loss

- HQ encounters inelastic scattering as it traveses a medium
 - Emits a gluon



- Differential Loss: dE/dx
- Very simple but first estimation:
 - Mustafa, Pal, Srivastava, Thoma, PLB 428 (1998) 234
 - Improvements in formalism

Radiative Energy Loss

- Singnificant Points:
 - The nature of the medium through which the energetic parton propagates ➤ thermally equilibrated perturbative medium with a collection of static scatt. centres with specified density
 - Kinematic approximations for interation between medium partons (thermal background) and the projectile parton that is propagating in the medium
 - The virtuality and branching/splitting of hard parton to reduce its off-shellness ➤ multiple splitting may occure in the medium ➤ multipluon final state should include interference of emitted gluons

HQ Radiative Energy Loss

Suppression due to Mass and Dead Cone

General notion: heavy quark radiates less than light quark

$$\begin{array}{|c|c|c|c|c|}\hline qq' \to qq' & |\mathcal{M}_{qq' \to qq'}|^2 & = \frac{8}{9}g^4\frac{s^2}{t^2} \\ \hline Qq \to Qq & |\mathcal{M}_{Qq \to Qq}|^2 & = \frac{8}{9}g^4\frac{s^2}{t^2} \left(1 - \frac{M^2}{s}\right)^2 \\ \hline qq' \to qq'g & |\mathcal{M}_{qq' \to qq'g}|^2 & = 12g^2\frac{8}{9}g^4\frac{s^2}{t^2}\frac{1}{k_\perp^2} \\ \hline Qq \to Qqg & |\mathcal{M}_{Qq \to Qqg}|^2 & = 12g^2\frac{8}{9}g^4\frac{s^2}{t^2}\frac{1}{k_\perp^2} \end{array}$$


2001 Dokshitzer and Kharzeev proposed (Phys. Lett. B 519, 199 (2001)) "dead cone" effect => heavy quark small energy loss.

$$\left(1 + \frac{\theta_0^2}{\theta^2}\right)^{-2} \qquad \theta_0 = M/E$$

HQ Radiative Energy Loss

Generalised Dead Cone

ullet RHIC data (PHENIX) $lackbrack R_{AA}^{HQ} \sim R_{AA}^{LQ} \,lacksquare$ (Heavy quark puzzle)



Hierarchy employed in this study
$$\sqrt{s}, E \gg \sqrt{|t|} \sim q_{\perp} \gg \omega > k_{\perp} \gg m_D$$

Emission angle Range
$$-\pi < \theta < +\pi$$

Mass Range

$$\begin{split} \left| \mathcal{M}_{Qq \to Qqg} \right|^2 &= 12g^2 \left| \mathcal{M}_{Qq \to Qq} \right|^2 \frac{1}{k_\perp^2} \left(1 + \frac{M^2}{s \tan^2(\frac{\theta}{2})} \right)^{-2} \\ &= 12g^2 \left| \mathcal{M}_{Qq \to Qq} \right|^2 \frac{1}{k_\perp^2} \left(1 + \frac{M^2}{s} e^{2\eta} \right)^{-2} \\ \mathcal{D} = \left(1 + \frac{M^2}{s \tan^2(\frac{\theta}{2})} \right)^{-2} \end{split}$$

Martinez,
Mustafa,
Uphoff, PRD85
(2012) 054012

0 < M/E < 1

HQ Radiative Energy Loss

Radiation Spectrum [Abir, Greiner, Martinez, Mustafa, Uphoff, PRD85 (2012)054012]

$$lacksquare Qq o Qqg$$
 and $Qar q o Qar qg$; Replace $Q o ar Q$

$$\left. \frac{dn_g}{d\eta \ dk_\perp^2} \right|_{Qq \to Qqg} = \left. \frac{C_A \alpha_s}{\pi} \frac{1}{k_\perp^2} \mathcal{D} \right.$$

$$\left. \frac{dn_g}{d\eta \ dk_\perp^2} \right|_{Qg \to Qgg} = \frac{C_A}{C_F} \left. \frac{dn_g}{d\eta \ dk_\perp^2} \right|_{Qg \to Qgg}; \ \frac{C_A}{C_F} = \frac{9}{4}$$

Radiative E-Loss [Abir, Jamil, Mustafa, Srivastava, PLB715 (2012) 183]


Differential Radiative E-Loss

$$-\frac{dE}{dx} = \frac{\langle \omega \rangle}{\langle \lambda \rangle}$$

 $\langle \omega
angle =$ Mean energy of the emitted gluon


 $\langle \lambda
angle =$ Mean free path of the heavy quark


Radiative E-Loss [Abir, Jamil, Mustafa, Sivastava, PLB715 (2012) 183]






Radiative E-Loss [Abir, Jamil, Mustafa, Sivastava, PLB715 (2012) 183]






Comparison with Parton Cascade Model [Younus et al, Phys.Rev. C91 (2015) 024912]





Jet Quenching



- Jets are produced back to back
 - Awayside will pass through medium
 - Interact with the medium
 - Lose energy and quenched
 - Results in suppression of hadronic yields in AA than NN
- Nuclear Suppression Factor: $R_{AA} = \frac{\left(\mathsf{Yield}\right)^{AA}}{N_{\mathsf{coll}} \ \left(\mathsf{Yield}\right)^{NN}}$
- Quenching depends on amount of energy-loss suffered in the medium


D-Meson @ LHC 2.76ATeV






- DATA: CMS Collaboration, CMS-PAS-HIN-16-011.
- Theory: Abir, Jamil, Mustafa, Srivastava, PLB715 (2012) 183
- Saraswat, Shukla, Singh, NPA961(2017)169 M G Mustafa (SINP)

D-Meson @ LHC 5.02ATeV





- Data: CMS Collaboration, CMS-PAS-HIN-16-001.
- Theory: Abir, Jamil, Mustafa, Srivastava, PLB715 (2012) 183
- Saraswat, Shukla, Singh, NPA961(2017)169

M G Mustafa (SINP) 3rd HF Meet ⋅ ⋅ ⋅ IIT, Indore: 19/3/19 29 / 41

Collisional Energy

- Assumed that collisional energy lost by the particle per unit time is small compared to the energy of the particle itself
- the change in the velocity of the particle during the motion may be neglected, i.e, the particle moves in a straight line trajectory.
- The energy loss of a particle is determined by the work done per unit time by the retarding chromo electric forces acting on the particle in the plasma while moving:

$$\left. rac{dE}{dt}
ight|_{\mathsf{coll}} = \left. Q^a ec{v} \cdot ec{\mathcal{E}}^a
ight|_{ec{r} = ec{v}t}$$

- The collisional energy loss does not take into account :
- the field fluctuation in the plasma
- the particle recoil in collisions
- One needs to take in to account both : How??

Collisional Energy Gain [Chakraborty, Mustafa, Thoma, PRC75 (2007) 064908]

The work done per unit time reads as

$$\left. rac{dE}{dt}
ight|_{\mathsf{fluc}} = \left\langle Q^a ec{v}(t) \cdot ec{\mathcal{E}}_t^a(ec{r}(t),t)
ight
angle$$

- The electric field \mathcal{E}^a_t consists of the induced field and a spontaneously generated microscopic field, the latter being a random function of position and time
- The stochastic time dependence comes because of explicit fluctuations of field in time due to change of the motion of the particle from one field point to another
- $lue{r}$ Both $\vec{r}(t)$ and $\vec{v}(t)$ can be obtained from EOM:

$$\frac{d\vec{p}}{dt} = Q^a \left[\mathcal{E}_t^a(\vec{r}(t), t) + \vec{v} \times \mathcal{B}_t^a(\vec{r}(t), t) \right]$$

The magnetic field \mathcal{B}^a_t consists of an induced and a microscopic part as in the case of \mathcal{E}^a_t

Collisional Energy Gain [Chakraborty, Mustafa, Thoma, PRC75 (2007) 064908]

The spectrum of the chromo-electromagnetic fields follows from the fluctuation-dissipation theorem

$$\langle \mathcal{E}_i^a \mathcal{E}_j^b \rangle_{\beta;\omega,k} = \delta^{ab} \langle \mathcal{E}_i \mathcal{E}_j \rangle_{\beta;\omega,k}$$


$$\langle \mathcal{E}_{i}\mathcal{E}_{j}\rangle = \frac{8\pi}{e^{\beta\omega} - 1} \left[\mathcal{P}_{ij}^{L} \frac{\operatorname{Im} \, \varepsilon_{L}}{\left|\varepsilon_{L}\right|^{2}} + \mathcal{P}_{ij}^{T} \frac{\operatorname{Im} \, \varepsilon_{T}}{\left|\varepsilon_{T} - \frac{\omega^{2}}{k^{2}}\right|^{2}} \right]$$


- This is completely determined by the dielectric functions of the medium
- This leads to energy gain as

$$\begin{aligned} \frac{dE}{dx} \Big|_{fluc} &= \frac{C_F \alpha_s}{16\pi^3 E} \int d^3k \left[\frac{\partial}{\partial \omega} \langle w \vec{\mathcal{E}}_L^2 \rangle + \langle \vec{\mathcal{E}}_T^2 \rangle \right]_{\omega = \vec{k} \cdot \vec{v}} \\ &= 2\pi C_F \alpha_s^2 \left(1 + \frac{N_f}{6} \right) \frac{T^3}{E v^2} \ln \frac{1+v}{1-v} \ln \frac{k_{\text{max}}}{k_{\text{min}}} \end{aligned}$$

Collisional Energy Gain

The field fluctuations lead to energy gain:





33 / 41


Chakraborty, Mustafa, Thoma, PRC75 (2007) 064908

M G Mustafa (SINP) 3rd HF Meet · · · IIT, Indore: 19/3/19

Fractional Energy Gain of Charm Quark

9 Coll. Gain/Energy= $\frac{\Delta E}{E}$


A Ikbal, Z Ahamed, M G Mustafa, : arXiv:1902.06989 Ikbal, Ahamed, Shukla, Mustafa, Phys .Rev. C98 (2018) 034915



Diffusion Coefficient of Charm Quark

Langevin Dynamics

A Ikbal, Z Ahamed, M G Mustafa, : arXiv:1902.06989 Ikbal, Ahamed, Shukla, Mustafa, PRC98 (2018) 034915



- lacktriangledown Drag : $\gamma = -rac{1}{p}rac{dE}{dx}$
- FD Theorem:

Mom. Diff.: $D=2\gamma ET$


Einstein Relation:


$$D_s \stackrel{=}{\underset{p \to 0}{=}} \frac{T}{\gamma M} = \frac{2T^2}{D}$$

Banerjee, Datta, Gavai and Majumdar, PRD 85 014510 (2012)

Open Charm and Field Fluctuation

Nuclear Modification factor (R_{AA}) for D Meson at 2.76 TeV Ikbal, Ahamed, Shukla, Mustafa, Phys .Rev. C98 (2018) 034915







- DATA:CMS Collaboration , CMS-PAS-HIN-16-011
- ALICE Collaboration, JHEP 1603 (2016) 081, 2016

Open Charm/Beauty and Field Fluctuation

Nuclear Modification factor (R_{AA}) for D & B-Meson at 5.02 TeV lkbal, Ahamed, Shukla, Mustafa, Phys .Rev. C98 (2018) 034915






37 / 41


DATA: CMS Collaboration , CMS-PAS-HIN-16-011

M G Mustafa (SINP) 3rd HF Meet · · · IIT, Indore: 19/3/19

Hidden Charm and Field Fluctuation

Nuclear Modification factor (R_{AA}) for J/ψ at 2.76 TeV A Ikbal, Z Ahamed, M G Mustafa, : arXiv:1902.06989





- ALICE Collaboration, J. High Energy Phys. 07, 051 (2015)
- CMS Collaboration, Eur. Phys. J. C 77, 252 (2017)

Conclusion:

- Discussed why are heavy quarks important
- Heavy quark production in high energy HIC: X-section in pQCD
- Collisional E-Loss & E-Gain
 - Relation to Transport Coeffs: Drag and Diffusion
 - Coll. E-Loss Distr. & importance of Coll. E-Loss
- Radiative E-Loss
 - Generalised Dead Cone and radiative E-Loss
 - Light and heavy quark lose energy in a similar fashion
- lacksquare Importance Chromoelectric Field Fluctuations & Energy Gain
- lacksquare Nuclear suppression R_{AA} for various heavy quarked meson
 - $lue{r}$ Both D and B Meson R_{AA} @ LHC for 2.76ATeV and 5.02ATeV in Pb+Pb
 - $= J/\Psi R_{AA}$ © LHC for 2.76ATeV in Pb+Pb without regeneration

Collaborators:

- Raktim Abir (AMU, Aligarh)
- Purnendu Chakraborty (Basirhat College, W. Bengal)
- Umme Jamil (D.R. College, Golaghata, Assam)
- Dipali Pal (USA)
- Asik Ikbal (VECC, Kolkata)
- D. K. Srivastava (VECC, Kolkata)
- Zubayer Ahmed (VECC, Kolkata)
- Prashant Shukla (BARC, Mumbai)
- Markus H. Thoma (Giessen Univ., Germany)
- Carsten Greiner (Goethe Univ. Frankfurt, Germany)
- Jan Uphoff (Goethe Univ., Frankfurt, Germany)
- M. Martinez (FIAS, Frankfurt, Germany)

THANK YOU