
Next-to-leading power corrections for Drell-Yan and prompt photon production

PARTICLEFACE meeting
Coimbra, 26-28 Feb.

Eric Laenen



UNIVERSITEIT VAN AMSTERDAM




Universiteit Utrecht

Outline

- ✦ Next-to-leading power threshold logarithms
 - ▶ Regions, factorization at NNLO
- ✦ NLP logs at NLO, simple formula
 - ▶ extension to prompt photon
- ✦ LL resummation at NLP
 - ▶ using NLP webs

Threshold logarithms



Log of “energy excess above
production threshold”

$$S \geq s \geq Q^2$$

$$L^2 = \ln \left(1 - \frac{Q^2}{s} \right) \equiv \ln^2(1 - z)$$

NLP threshold behavior

- ♦ For Drell-Yan, DIS, Higgs, singular behavior in perturbation theory when $z \rightarrow 1$

$$\delta(1-z) \left[\frac{\ln^i(1-z)}{1-z} \right]_+ \ln^i(1-z)$$

- ▶ plus distributions have been organized to all orders (=“resummation”), also possible for $\ln(1-z)$?
- ♦ “Zurich” method of threshold expansion allows computation (for NNNLO Higgs production)

$$(1-z)^p \ln^q(1-z)$$


Anastasiou, Duhr, Dulat, Furlan,
Gehrmann, Herzog, Mistlberger

– done to $p=37..$

Larkoski, Neill, Stewart, Moult, Kolodrubetz, Rothen,
Zhu, Tackmann, Vita, Feige ;
Beneke, Campanario, Mannel, Peckja

- ♦ Much development in SCET
- ♦ Useful also for improving NNLO slicing (N-jettiness) methods
- ♦ Alternative terminology to “NLP”
 - ▶ Next-to-soft
 - ▶ Next-to-eikonal

PARTICLEFACE workshop on NLP, Nov. 2018



Next-to-eikonal Feynman rules

- ♦ Keep 1 term more in k expansion beyond eikonal approximation

$$\begin{aligned} \text{scalar : } & \frac{2p^\mu + k^\mu}{2p \cdot k + k^2} \rightarrow \frac{2p^\mu}{2p \cdot k} + \frac{k^\mu}{2p \cdot k} - k^2 \frac{2p^\mu}{(2p \cdot k)^2} \\ \text{fermion : } & \frac{\not{p} + \not{k}}{2p \cdot k + k^2} \gamma^\mu u(p) \rightarrow \left[\frac{2p^\mu}{2p \cdot k} + \frac{\not{k} \gamma^\mu}{2p \cdot k} - k^2 \frac{2p^\mu}{(2p \cdot k)^2} \right] u(p) \end{aligned}$$

- Becomes emitter-spin dependent, recoil now included
- Is there predictive power for the next-to-eikonal terms?

Eikonal term

Classic NLP result: Low's theorem

- These rules are good for emissions from external lines. At NLP order, also 1 “internal” emission contributes

$$\Gamma^\mu =$$

- Low's theorem (scalars, generalization to spinors by Burnett-Kroll, to massless particles by Del Duca): **LBKD theorem**

✓ Work to order k , and use Ward identity

$$\Gamma^\mu = \left[\frac{(2p_1 - k)^\mu}{-2p_1 \cdot k} + \frac{(2p_2 + k)^\mu}{2p_2 \cdot k} \right] \Gamma + \left[\frac{p_1^\mu (k \cdot p_2 - k \cdot p_1)}{p_1 \cdot k} + \frac{p_2^\mu (k \cdot p_1 - k \cdot p_2)}{p_2 \cdot k} \right] \boxed{\frac{\partial \Gamma}{\partial p_1 \cdot p_2}}$$

- Elastic amplitude still determines the emission to NLP accuracy,
 - note the derivative
 - detailed knowledge of “internal part” not needed

NLP logarithms for Drell-Yan


- ♦ Goal: combine (N)LP matrix elements with (N)LP phase space to **predict** $\ln^i(1-z)$ for NNLO Drell-Yan

$$\frac{1}{\sigma^{(0)}} \frac{d\hat{\sigma}}{dz} \sim \int d\Phi_{\text{LP}} |\mathcal{M}|_{\text{LP}}^2 + \int d\Phi_{\text{LP}} |\mathcal{M}|_{\text{NLP}}^2 + \int d\Phi_{\text{NLP}} |\mathcal{M}|_{\text{LP}}^2 + \dots$$

- ▶ We pursue two methods:
 - ✓ 1. Method of regions
 - ✓ 2. Factorization
- ▶ NLO is “easy”, real test at NNLO

NLP logs in Drell-Yan at NNLO

- Check NLP Feynman rules for NNLO Drell-Yan *double real* emission




- Result at NLP level, agrees with equivalent exact result. C_F^2 terms e.g.

$$K_{\text{NE}}^{(2)}(z) = \left(\frac{\alpha_s}{4\pi} C_F\right)^2 \left[-\frac{32}{\epsilon^3} \mathcal{D}_0(z) + \frac{128}{\epsilon^2} \mathcal{D}_1(z) - \frac{128}{\epsilon^2} \log(1-z) \right. \\ \left. - \frac{256}{\epsilon} \mathcal{D}_2(z) + \frac{256}{\epsilon} \log^2(1-z) - \frac{320}{\epsilon} \log(1-z) \right. \\ \left. + \frac{1024}{3} \mathcal{D}_3(z) - \frac{1024}{3} \log^3(1-z) + 640 \log^2(1-z) \right],$$

$$\mathcal{D}_i = \left[\frac{\log^i(1-z)}{1-z} \right]_+$$

- Next, 1 Real- 1 Virtual




Diagnosis: method of regions

Beneke, Smirnov

♦ How does it work?

- ▶ Divide up k_1 (=loop-momentum) integral into hard, 2 collinear and a soft region, by appropriate scaling

Hard : $k_1 \sim \sqrt{\hat{s}} (1, 1, 1)$; Soft : $k_1 \sim \sqrt{\hat{s}} (\lambda^2, \lambda^2, \lambda^2)$;
 Collinear : $k_1 \sim \sqrt{\hat{s}} (1, \lambda, \lambda^2)$; Anticollinear : $k_1 \sim \sqrt{\hat{s}} (\lambda^2, \lambda, 1)$.



- ▶ expand integrand in λ , to leading and next-to-leading order
- ▶ but then integrate over *all* k_1 anyway!
- ▶ Treat emitted momentum as soft and incoming momenta as hard

$$k_2^\mu = (\lambda^2, \lambda^2, \lambda^2)$$

Method of region result

Bonocore, EL, Magnea, Vernazza, White

- ◆ Results

- ▶ Hard region (expansion in λ^2): *LP + some NLP*
- ▶ Soft region (expansion in λ^2): *ZERO*
- ▶ (anti-)collinear regions (expansion in λ): *NLP only*

- ◆ Result:

- ▶ the full $K^{(1)}_{1r,1v}$ is reproduced, including constants

- ◆ For **predictive power**, need factorization

A factorization approach from Low's theorem

Bonocore, EL, Magnea, Melville, Vernazza, White



♦ Can we *predict* the $\ln(1-z)$ logarithms from lower orders?

► Factorize the cross section,

✓ H: the hard and the soft function

✓ J: incoming-jet functions

♦ Next, add one extra soft emission. Let every blob radiate!



Del Duca

► Compute each new “blob + radiation”, and put it together. New: **radiative jet function**

$$J_\mu(p, n, k, \alpha_s(\mu^2), \epsilon) u(p) = \int d^d y \, e^{-i(p-k) \cdot y} \langle 0 | \Phi_n(y, \infty) \psi(y) j_\mu(0) | p \rangle$$

Factorization approach to NLP logarithms

- ✦ Upshot: a factorization formula for the emission amplitude

$$\mathcal{A}_{\mu,a}(p_j, k) = \sum_{i=1}^2 \left(\frac{1}{2} \tilde{\mathcal{S}}_{\mu,a}(p_j, k) + g \mathbf{T}_{i,a} G_{i,\mu}^\nu \frac{\partial}{\partial p_i^\nu} + J_{\mu,a}(p_i, n_i, k) \right) \mathcal{A}(p_j) - \mathcal{A}_{\mu,a}^{\tilde{\mathcal{J}}}(p_j, k)$$

Soft function

Orbital term

Jet function

Overlap

- ▶ J_μ is needed at one-loop level

Predicted NLP threshold logs vs exact result

- Compute blobs, one-loop radiative jet function, contract with cc amplitude and integrate over phase space. Exact calculation gives

$$\begin{aligned}
 K_{\text{rv}}^{(2)}(z) = & \left(\frac{\alpha_s}{4\pi} \right)^2 \left\{ C_F^2 \left[\frac{32\mathcal{D}_0(z) - 32}{\epsilon^3} + \frac{-64\mathcal{D}_1(z) + 48\mathcal{D}_0(z) + 64L(z) - 96}{\epsilon^2} \right. \right. \\
 & + \frac{64\mathcal{D}_2(z) - 96\mathcal{D}_1(z) + 128\mathcal{D}_0(z) - 64L^2(z) + 208L(z) - 196}{\epsilon} - \frac{128}{3}\mathcal{D}_3(z) \\
 & \left. \left. + 96\mathcal{D}_2(z) - 256\mathcal{D}_1(z) + 256\mathcal{D}_0(z) + \frac{128}{3}L^3(z) - 232L^2(z) + 412L(z) - 408 \right] \right. \\
 & + C_A C_F \left[\frac{8\mathcal{D}_0(z) - 8}{\epsilon^3} + \frac{-32\mathcal{D}_1(z) + 32L(z) - 16}{\epsilon^2} + \frac{64\mathcal{D}_2(z) - 64L^2(z) + 64L(z) + 20}{\epsilon} \right. \\
 & \left. \left. - \frac{256}{3}\mathcal{D}_3(z) + \frac{256}{3}L^3(z) - 128L^2(z) - 60L(z) + 8 \right] \right\}, \tag{4.6}
 \end{aligned}$$

$$L(z) = \ln(1 - z)$$

- Result:** *perfect agreement* for 4 powers of the next-to-eikonal/soft logarithms at NNLO

$$\ln^3(1 - z), \quad \ln^2(1 - z), \quad \ln^1(1 - z), \quad \ln^0(1 - z),$$

Colour-singlet final states

- ◆ Generalize NLP factorization (the LBKD theorem) beyond Drell-Yan, to arbitrary colour-singlet final states
 - ▶ look at NLO only, i.e. predict

$$D_1 = \left[\frac{\ln(1-z)}{1-z} \right]_+ \quad D_0 = \left[\frac{1}{1-z} \right]_+ \quad L_1 = \ln(1-z) \quad L_0 = \ln^0(1-z)$$

- ✓ where “1-z” can take different forms for 2 -> 2,3 etc scattering
- ▶ apply to Drell-Yan, (multi-)Higgs, (vector boson pairs)
- ▶ for inclusive and fully differential cross sections

NLP terms in colorless final states @NLO

Bonocore, Del Duca, EL, Magnea, Vernazza, White
1706.04018

Previous factorization at NLO

$$\mathcal{A}_{\mu,a}^{(1)}(\{p_i\}, k) = \sum_{l=1}^2 \left[g_s \mathbf{T}_{l,a} G_{l,\mu}^\nu \frac{\partial}{\partial p_l^\nu} + J_{\mu,a}^{(1)}(p_l, n_l, k) \right] \mathcal{A}^{(0)}(\{p_i\})$$

✓ G is a projector, \mathbf{T} a color matrix

- ▶ initial quarks: $J_\mu^a(p, n, k) = g_s \mathbf{T}^a \left[\frac{(2p - k)_\mu}{2p \cdot k} + \frac{ik^\beta}{p \cdot k} S_{\beta\mu} \right] \quad S_{\beta\mu} = \frac{i}{4} [\gamma_\beta, \gamma_\mu]$
- ▶ initial gluons: $J_{\mu,\rho\sigma}^a(p, n, k) = g_s \mathbf{T}^a \left[\frac{(2p - k)_\mu}{2p \cdot k} \eta_{\rho\sigma} - \frac{ik^\beta}{p \cdot k} M_{\beta\mu,\rho\sigma} \right] \quad M_{\beta\mu,\rho\sigma} = i(\eta_{\beta\rho}\eta_{\mu\sigma} - \eta_{\beta\sigma}\eta_{\mu\rho})$
- ▶ notice the spin-dependent Lorentz generator (“next-to-soft theorem”)
- ▶ notice derivative term (Low’s theorem)

Lorentz generator

- ♦ The derivative term can be written as the **orbital part** of Lorentz generator

$$G_{l,\mu}^\nu \frac{\partial}{\partial p_l^\nu} = \frac{k^\nu}{p_l \cdot k} \left[p_{l,\nu} \frac{\partial}{\partial p_l^\mu} - p_{l,\mu} \frac{\partial}{\partial p_l^\nu} \right] = -\frac{ik^\nu L_{\nu\mu}^{(l)}}{p_l \cdot k}$$

- ♦ so that


$$\begin{aligned} \mathcal{A}_{\mu,a}^{(1)}(\{p_i\}, k) &= \sum_{l=1}^2 g_s \mathbf{T}_{l,a} \left[\frac{(2p_l - k)_\mu}{2p_l \cdot k} - \frac{ik^\nu}{p_l \cdot k} \underbrace{\left(L_{\nu\mu}^{(l)} + \Sigma_{\nu\mu}^{(l)} \right)}_{J_{\nu\mu}^{(l)}} \right] \mathcal{A}^{(0)}(\{p_i\}) \\ &= \sum_{l=1}^2 g_s \mathbf{T}_{l,a} \left[\frac{p_{l,\mu}}{p_l \cdot k} - \frac{ik^\nu J_{\nu\mu}^{(l)}}{p_l \cdot k} \right] \mathcal{A}^{(0)}(\{p_i\}) \end{aligned}$$

- leads to Scalar + Orbital + Spin part of the NLP amplitude

Colour singlet production in gg channel

◆ Square amplitude

$$|\mathcal{A}_{\text{NLP}}|^2 = \sum_{\text{colours}} \left(\mathcal{A}_{\text{scal.}}^{\sigma_1, \mu_1 \nu_1} + \mathcal{A}_{\text{spin}}^{\sigma_1, \mu_1 \nu_1} + \mathcal{A}_{\text{orb.}}^{\sigma_1, \mu_1 \nu_1} \right)^* \mathcal{P}_{\mu_1 \mu_2}(p_1, l_1) \mathcal{P}_{\nu_1 \nu_2}(p_2, l_2) \mathcal{P}_{\sigma_1 \sigma_2}(k, l_3) \\ \times \left(\mathcal{A}_{\text{scal.}}^{\sigma_2, \mu_2 \nu_2} + \mathcal{A}_{\text{spin}}^{\sigma_2, \mu_2 \nu_2} + \mathcal{A}_{\text{orb.}}^{\sigma_2, \mu_2 \nu_2} \right), \quad (3.7)$$



✓ where $\mathcal{P}_{\alpha\beta}(p, l) \equiv \sum_{\lambda} \epsilon_{\alpha}^{(\lambda)}(p) \epsilon_{\beta}^{(\lambda)*}(p) = -\eta_{\alpha\beta} + \frac{p_{\alpha} l_{\beta} + p_{\beta} l_{\alpha}}{p \cdot l}$

- ▶ Can be done using $-\eta_{\alpha\beta}$ only (external ghosts are beyond NLP)
- ▶ Truncate to NLP, leads to

$$|\mathcal{A}_{\text{NLP}}|^2 = \sum_{\text{colours}} \left\{ |\mathcal{A}_{\text{scal.}}^{\sigma, \mu\nu}|^2 + 2\text{Re} \left[\left(\mathcal{A}_{\text{spin}}^{\sigma, \mu\nu} + \mathcal{A}_{\text{orb.}}^{\sigma, \mu\nu} \right)^* \mathcal{A}_{\text{scal.}}^{\sigma, \mu\nu} \right] \right\}$$

- ▶ Easy part: scalar (eikonal) part

$$\sum_{\text{colours}} |\mathcal{A}_{\text{scal.}}^{\sigma, \mu\nu}|^2 = 2g_s^2 N_c (N_c^2 - 1) \frac{p_1 \cdot p_2}{p_1 \cdot k p_2 \cdot k} |\mathcal{A}_{\mu\nu}|^2$$

CS production in gg channel

- ✦ Spin * scalar vanishes (anti-symmetry in $\mu\nu$)
- ✦ Orbital part leads to shifts in momentum dependence

$$\sum_{\text{colours}} 2 \operatorname{Re} [\mathcal{A}_{\text{orb.}}^{\sigma, \mu\nu} \mathcal{A}_{\text{scal.}}^{\sigma, \mu\nu}] = \frac{2g_s^2 N_c (N_c^2 - 1) p_1 \cdot p_2}{p_1 \cdot k p_2 \cdot k} \left[\delta p_1^\alpha \frac{\partial}{\partial p_1^\alpha} + \delta p_2^\alpha \frac{\partial}{\partial p_2^\alpha} \right] |\mathcal{A}_{\mu\nu}|^2$$


✓ where

$$\delta p_1^\alpha = -\frac{1}{2} \left(\frac{p_2 \cdot k}{p_1 \cdot p_2} p_1^\alpha - \frac{p_1 \cdot k}{p_1 \cdot p_2} p_2^\alpha + k^\alpha \right), \quad \delta p_2^\alpha = -\frac{1}{2} \left(\frac{p_1 \cdot k}{p_1 \cdot p_2} p_2^\alpha - \frac{p_2 \cdot k}{p_1 \cdot p_2} p_1^\alpha + k^\alpha \right)$$

- ✦ Result is simple: dipole times shifted squared LO amplitude

$$|\mathcal{A}_{\text{NLP}}|^2 = \frac{2g_s^2 N_c (N_c^2 - 1) p_1 \cdot p_2}{p_1 \cdot k p_2 \cdot k} |\mathcal{A}_{\mu\nu}(p_1 + \delta p_1, p_2 + \delta p_2)|^2$$

CS production in $q\bar{q}$ channel



- ♦ Scalar plus orbital part very similar to gg case

$$|A_{\text{NLP}}^\sigma|_{\text{scal.}+\text{orb.}}^2 = \frac{g_s^2 C_F}{\underset{\text{blue}}{z}} \frac{s}{p_1 \cdot k p_2 \cdot k} |A(p_1 + \delta p_1, p_2 + \delta p_2)|^2$$

- ▶ except for the $1/z$, which is due to the kinematic shift

$$s \rightarrow (p_1 + p_2 + \delta p_1 + \delta p_2)^2 = s + 2(\delta p_1 + \delta p_2) \cdot (p_1 + p_2)$$

- ▶ which is the same as

$$s \rightarrow z s$$

- ♦ But the spin part now does not cancel:

$$\sum_{\text{colours}} 2 \text{Re} \left[A_{\text{scal.}}^\dagger A_{\text{spin}} \right]_{\text{NLP}} = -g_s^2 N_c C_F \frac{2p_1 \cdot p_2}{p_1 \cdot k p_2 \cdot k} \underbrace{\frac{k \cdot (p_1 + p_2)}{p_1 \cdot p_2}}_{\text{blue}} |\mathcal{A}(p_1, p_2)|^2$$

- ▶ precisely compensates $1/z \cong 1 + (1-z)!!$

$$\underbrace{\frac{k \cdot (p_1 + p_2)}{p_1 \cdot p_2}}_{\text{blue}} = -(1-z)$$

Squared amplitudes and cross sections

✦ In summary

✦ **gluons** $|\mathcal{A}_{\text{NLP}}|^2 = \frac{2g_s^2 N_c (N_c^2 - 1) p_1 \cdot p_2}{p_1 \cdot k p_2 \cdot k} |\mathcal{A}_{\mu\nu}(p_1 + \delta p_1, p_2 + \delta p_2)|^2$

✦ **quarks** $|A_{\text{NLP}}|^2 = g_s^2 C_F \frac{s}{p_1 \cdot k p_2 \cdot k} |A(p_1 + \delta p_1, p_2 + \delta p_2)|^2$

✦ Up to colour factors the same:


✦ **eikonal (dipole) factor times shifted Born cross section**

✦ **Born can be loop-induced, have complex parts etc.**

✦ Combine carefully with phase space for general inclusive formula

$$\frac{d\hat{\sigma}_{\text{NLP}}^{(gg)}}{dz} = C_A K_{\text{NLP}}(z, \epsilon) \hat{\sigma}_{\text{Born}}^{(gg)}(zs) \quad K_{\text{NLP}}(z, \epsilon) = \frac{\alpha_s}{\pi} \left(\frac{4\pi\mu^2}{s} \right)^\epsilon z(1-z)^{-1-2\epsilon} \frac{\Gamma^2(-\epsilon)}{\Gamma(-2\epsilon)\Gamma(1-\epsilon)}$$

Single Higgs production



Infinite top mass limit not needed
extra operators = shift in kinematics



♦ Single Higgs production

$$\frac{d\sigma_{\text{NLP}}^h}{dz} = \frac{\alpha_s^3 C_A}{288\pi^2 v^2} F(z\tau, \epsilon) \left(\frac{2 - \mathcal{D}_0(z)}{\epsilon} + 2\mathcal{D}_1(z) - \mathcal{D}_0(z) - 4\log(1 - z) + 2 \right)$$

- ▶ with F the well-known Born function. \mathcal{D} 's and \mathcal{L} 's agree with exact calculation, but also with full top mass dependence!

Dawson; Spira, Djouadi,
Graudenz, Zerwas

Di-Higgs production



Borowka, Greiner, Heinrich, Jones,
Kerner, Schlenk, Zirke

◆ Double Higgs production at NLO-NLP

$$z \frac{d\sigma_{\text{NLP}}^{hh}}{dz} = \frac{\alpha_s}{3\pi} C_A \left(\frac{\bar{\mu}^2}{s} \right)^\epsilon \left[\frac{12 - 6\mathcal{D}_0(z)}{\epsilon} + 12\mathcal{D}_1(z) - 24 \log(1 - z) \right] \sigma_{\text{Born}}^{hh}(zs)$$

▶ where

$$\frac{d\hat{\sigma}_{\text{Born}}^{hh}}{dt} = \frac{\alpha_s^2}{8\pi^3} \frac{1}{512 v^4} \left[|C_\Delta F_\Delta + C_\square F_\square|^2 + |C_\square G_\square|^2 \right]$$

✓ with triangle and box graphs, again for full top mass dependence

▶ Should be useful for numerical evaluations, and seeing new patterns

◆ Similar result for triple-Higgs production



De Florian, Mazzitelli

Final state partons: Prompt photon production

Beenakker, van Beekveld, EL, White
to appear

With FS partons: prompt photon

- ♦ Two LO channels: $q\bar{q}$ and qg




- ♦ With extra radiation, different ways to define threshold. We shall use “w” \rightarrow 1

$$\begin{aligned}u_1 &= (p_1 - p_\gamma)^2 \equiv -svw \\t_1 &= (p_2 - p_\gamma)^2 \equiv s(v - 1) \\s_4 &= s + t_1 + u_1 = sv(1 - w)\end{aligned}$$

- ♦ Two issues to deal with
 - shifting kinematics in $2 \rightarrow 2$ kinematics
 - soft fermion emission

Gluon emission

- ♦ For $q\bar{q}$ channel



- ♦ Can in fact write down general formula

$$\begin{aligned}\mathcal{A}_{\text{NLP}} &= \mathcal{A}_{\text{scal}} + \mathcal{A}_{\text{spin}} + \mathcal{A}_{\text{orb}} \\ &= \sum_{j=1}^{n+2} \frac{g_s \mathbf{T}_j}{2p_j \cdot k} \left(\mathcal{O}_{\text{scal},j}^{\sigma} + \mathcal{O}_{\text{spin},j}^{\sigma} + \mathcal{O}_{\text{orb},j}^{\sigma} \right) \otimes i\mathcal{M}_{\text{H}}(p_1, \dots, p_i, \dots, p_{n+2}) \epsilon_{\sigma}^*(k),\end{aligned}$$

- ▶ color charge and spin generator depends on emitting IS or FS particle
- ▶ orbital part on IS or FS particle

Squared amplitude at NLP

- ♦ Result: again dipoles plus momentum shift
- ♦ Important to implement $2 \rightarrow 3$ momentum conservation in $2 \rightarrow 2$ matrix element
 - ▶ used Catani-Seymour dipoles (FKS is also possible) Gervais

$$\begin{aligned}
 |\mathcal{A}_{\text{NLP}, q\bar{q} \rightarrow \gamma gg}|^2 = & \frac{C_F}{C_A} \left[C_F \frac{2p_1 \cdot p_2}{(p_1 \cdot k)(p_2 \cdot k)} |\mathcal{M}_{q\bar{q} \rightarrow \gamma g}(p_1 + \delta p_{1;2}, p_2 + \delta p_{2;1})|^2 \right. \\
 & + \frac{1}{2} C_A \frac{2p_1 \cdot p_R}{(p_1 \cdot k)(p_R \cdot k)} |\mathcal{M}_{q\bar{q} \rightarrow \gamma g}(p_1 + \delta p_{1;R}, p_R - \delta p_{R;1})|^2 \\
 & + \frac{1}{2} C_A \frac{2p_2 \cdot p_R}{(p_2 \cdot k)(p_R \cdot k)} |\mathcal{M}_{q\bar{q} \rightarrow \gamma g}(p_2 + \delta p_{2;R}, p_R - \delta p_{R;2})|^2 \\
 & \left. - \frac{1}{2} C_A \frac{2p_1 \cdot p_2}{(p_1 \cdot k)(p_2 \cdot k)} |\mathcal{M}_{q\bar{q} \rightarrow \gamma g}(p_1 + \delta p_{1;2}, p_2 + \delta p_{2;1})|^2 \right].
 \end{aligned}$$


Note sign change for final state emitter

- ♦ Integrate over NLO phase, agrees with NLO calculation including $\ln(1-w)$ terms

Gordon, Vogelsang

Soft fermions

- At NLP (not LP) one can have soft fermion emission



- Effective feynman rule for left diagram (note that “u(k)” is of order \sqrt{k})

$$i\mathcal{M}_{\text{NLP},1,g} = \frac{g_s T_{c_m c_j}^a}{(p_1 - k)^2 + i\varepsilon} \epsilon^\mu(p_1) \bar{u}(k) \gamma_\mu \not{p}_1 \mathcal{M}_{c_j}(p_1, p_2, \dots, p_{n+2})$$

- Right diagram

$$i\mathcal{M}_{\text{NLP},1,g} = \frac{g_s T_{c_m c_i}^b}{(p_1 - k)^2 + i\varepsilon} \bar{u}(k) \gamma_\rho u(p_1) \mathcal{M}_{\rho, b}(p_1, p_2, \dots, p_{n+2}).$$

- Squaring amplitude and integration over phase space gives agreement with exact NLO
 - Must keep careful track of singular regions

LL resummation of NLP logarithms

Bahjat-Abbas, Bonocore, EL, Magnea, Sinninghe Damsté, Vernazza, White
to appear

LL resummation of NLP logarithms

- ♦ We have organized NLP threshold logs at NLO and NNLO for Drell-Yan. Can one resum them?
- ♦ First resummation conjecture: just change kernel in regular resummation formula

Kraemer, EL, Spira; 1998
EL, Magnea, Stavenga

$$\frac{1+z^2}{1-z} \longrightarrow \frac{2}{1-z} - 2$$

- ▶ reproduced NNLO NLP logs of van Neerven et al
- ♦ Physical kernel approach for inclusive quantities
 - ▶ using single log behaviour of kernel
- ♦ Recent LL resummation using SCET

Soar, Moch, Vermaseren, Vogt;
Moch, Vogt; Mattizelli, de Florian


Beneke, Broggio, Garny, Jaskiewicz,
Szafron, LV, Wang, 2018

$$\Delta_{\text{NLP}}^{\text{LL}}(z, \mu) = \exp \left[4S^{\text{LL}}(\mu_h, \mu) - 4S^{\text{LL}}(\mu_s, \mu) \right] \times \frac{-8C_F}{\beta_0} \ln \frac{\alpha_s(\mu)}{\alpha_s(\mu_s)} \theta(1-z).$$

Eikonal exponentiation


- After eikonal approximation, we suddenly see interesting patterns.

One loop vertex correction, in eikonal approximation




$$\mathcal{A}_0 \int d^n k \frac{1}{k^2} \frac{p \cdot \bar{p}}{(p \cdot k)(\bar{p} \cdot k)}$$

Two loop vertex correction, in eikonal approximation



$$\mathcal{A}_0 \frac{1}{2} \left(\int d^n k \frac{1}{k^2} \frac{p \cdot \bar{p}}{(p \cdot k)(\bar{p} \cdot k)} \right)^2$$


Exponential series



Yennie, Frautschi, Suura '61

QCD exponentiation: webs

- Not immediately generalizable to QCD, seemingly
 - Vertices terms have color charges, which don't commute
 - Still, an exponentiation theorem holds



$$\sum_D \mathcal{F}_D C_D = \exp \left[\sum_i \bar{C}_i w_i \right]$$

Webs

Exp

[

$$C_F \left(\text{triangle with vertical gluon} \right) + \left(-\frac{1}{2} C_A C_F \right) \left(\text{triangle with crossed gluons} \right) + \left(-\frac{1}{2} C_A C_F \right) \left(\text{triangle with Y-junction gluons} \right) + \dots$$

]

Gatheral; Frenkel, Taylor; Sterman

EL, Stavenga, White

Eikonal approximation from QM path integrals

EL, Stavenga, White


Use textbook result

$$\text{Sum of all diagrams} = \exp \left(\text{Connected diagrams} \right)$$




Write scattering amplitude as first-quantized path integral

$$M(p_1, p_2, \{k\}) = \int \mathcal{D}A_s \mathcal{D}x(t) H[x] f_1[A_s, x(t)] f_2[A_s, x(t)] e^{iS[A_s]}$$

Eikonal vertices are sources for gauge bosons along line



Disconnected



Connected

Can be generalized to non-abelian case (using replica trick)

NLP amplitude exponentiation via path integral


EL, Magnea, Stavenga, White

- Fluctuations around classical path are NE corrections

- All NLP corrections from external lines exponentiate
- Keep track via scaling variable λ

$$p^\mu = \lambda n^\mu$$

$$f(\infty) = \int_{x(0)=0} \mathcal{D}x \exp \left[i \int_0^\infty dt \left(\frac{\lambda}{2} \dot{x}^2 + (n + \dot{x}) \cdot A(x_i + nt + x) + \frac{i}{2\lambda} \partial \cdot A(x_i + p_f t + x) \right) \right]$$



- Exponentiation then in terms of NLP webs

$$\sum C(D) \mathcal{F}(D) = \exp [\bar{C}(D) W_E(D) + \bar{C}'(D) W_{NE}(D)]$$

Bonocore, EL, Magnea, Melville, Vernazza, White

LL resummation for cross section at NLP

- ✦ Can show that phase space NLP effects behave as

$$\varepsilon (1 - z)$$

- ▶ i.e. softness suppression comes with singularity suppression
- ✦ Can show that there are no LL enhancements from purely collinear regions (single log)
- ✦ LL effects come then only from NLP soft function

Exponentiating NLP soft function

- ✦ Moments of cross section

$$\int_0^1 d\tau \tau^{N-1} \frac{d\sigma_{\text{DY}}}{d\tau} \Big|_{\text{LL, NLP}} = \sigma_0(Q^2) q_N(Q^2) \bar{q}_N(Q^2) \bar{\mathcal{S}}_{\text{NLP}}(N, Q^2, \epsilon),$$

- ✦ with NLP soft function (f's are NLP Wilson lines)

$$\tilde{\mathcal{S}} = \frac{1}{N_c} \sum_n \text{Tr} \left[\langle 0 | f_2^\dagger f_1 | n \rangle \langle n | f_1^\dagger f_2 | 0 \rangle \right] \delta \left(z - \frac{Q^2}{\hat{s}} \right).$$

- ✦ Exponentiation then gives

$$\int_0^1 d\tau \tau^{N-1} \frac{d\sigma_{\text{DY}}}{d\tau} \Big|_{\text{LL, NLP}} = \sigma_0(Q^2) q_{\text{LL, NLP}}(N, Q^2) \bar{q}_{\text{LL, NLP}}(N, Q^2) \\ \times \exp \left[\frac{\alpha_s C_F}{\pi} \left(2 \log^2(N) + \frac{4 \log(N)}{N} \right) \right].$$

- ▶ agrees with old conjecture

Summary

- ✦ NLP factorization (LBDK theorem) leads to strong prediction for NLP threshold logs
 - ▶ Drell-Yan at NNLO
- ✦ Explicit formulae at NLO for any colour singlet final state, and now also coloured final state
- ✦ LL resummation at NLP for Drell-Yan done
 - ▶ NLL seems much harder..