

Signature of supersymmetry and $L_\mu - L_\tau$ gauge boson at Belle-II

Tuesday 5 February 2019 10:30 (30 minutes)

In this talk we will discuss a proposal that the γ + missing energy signal at the Belle-II detector will be a smoking gun for supersymmetry (SUSY) in the presence of a gauged $L_\mu - L_\tau$ symmetry. A striking consequence of breaking the enhanced symmetry appearing in the limit of degenerate (s)leptons is the non-decoupling of the radiative contribution of heavy charged sleptons

to the $\gamma - Z'$ kinetic mixing. The signal process, $e^+e^- \rightarrow \gamma Z' \rightarrow \gamma$ + missing energy, is an outcome of this ubiquitous feature. We take into account the severe constraints laid down on gauged $L_\mu - L_\tau$ models by several low-energy observables and show that any significant excess in all but the highest photon energy bin would be an undeniable signature of such heavy scalar fields in SUSY coupling to Z' . The number of signal events depends crucially on the logarithm of the ratio of stau to smuon mass in the presence of SUSY. In addition, the number is also inversely proportional to the $e^+ - e^-$ collision energy, making a low-energy, high-luminosity collider like Belle-II an ideal testing ground for this channel. This process can probe large swathes of the slepton mass ratio vs the additional gauge coupling (g_X) parameter space. More importantly, it can explore the narrow slice of $M_{Z'} - g_X$ parameter space still allowed in gauged $L_\mu - L_\tau$ models for superheavy sparticles.

Author: Prof. ROY, Sourov (Indian Association for the Cultivation of Science, Kolkata)

Co-author: Mr BANERJEE, Heerak (IACS)

Presenter: Prof. ROY, Sourov (Indian Association for the Cultivation of Science, Kolkata)

Session Classification: Plenary Session VII