

Charged Lepton Flavour Violation using Intense Muon Beams at Future Facilities

A. Baldini, D. Glenzinski, F. Kapusta, Y. Kuno, M. Lancaster,
J. Miller, S. Miscetti, T. Mori, A. Papa, A. Schöning, Y. Uchida

A submission to the 2020 update of the European Strategy for Particle Physics on behalf of the COMET, MEG, Mu2e and Mu3e collaborations.

Abstract

Charged-lepton flavour-violating (cLFV) processes offer deep probes for new physics with discovery sensitivity to a broad array of new physics models — SUSY, Higgs Doublets, Extra Dimensions, and, particularly, models explaining the neutrino mass hierarchy and the matter-antimatter asymmetry of the universe via leptogenesis. The most sensitive probes of cLFV utilize high-intensity muon beams to search for $\mu \rightarrow e$ transitions.

We summarize the status of muon-cLFV experiments currently under construction at PSI, Fermilab, and J-PARC. These experiments offer sensitivity to effective new physics mass scales approaching $\mathcal{O}(10^4)$ TeV/ c^2 . Further improvements are possible and next-generation experiments, using upgraded accelerator facilities at PSI, Fermilab, and J-PARC, could begin data taking within the next decade. In the case of discoveries at the LHC, they could distinguish among alternative models; even in the absence of direct discoveries, they could establish new physics. These experiments both complement and extend the searches at the LHC.

Contact: André Schöning [schoning@physi.uni-heidelberg.de]

Executive Summary

- Charged-lepton flavour-violating (cLFV) processes provide an unique discovery potential for physics beyond the Standard Model (BSM). These cLFV processes explore new physics parameter space in a manner complementary to the collider, dark matter, dark energy, and neutrino physics programmes.
- The global programme includes searches for $\mu \rightarrow e$, $\tau \rightarrow e$, and $\tau \rightarrow \mu$ transitions at experiments hosted in Europe, the US, and Asia. The relative rates among the various transitions are model dependent and comparisons among these transitions offer powerful model discrimination. A full exploration of cLFV parameter space requires the pursuit of all available $\mu \rightarrow e$ and $\tau \rightarrow e, \mu$ transitions.
- The most sensitive exploration of cLFV is provided by experiments that utilize high-intensity muon beams to search for cLFV $\mu \rightarrow e$ transitions: a muon decaying into an electron and a photon, $\mu^+ \rightarrow e^+ \gamma$ (MEG experiment at PSI); a muon decaying into three electrons $\mu^+ \rightarrow e^+ e^- e^+$ (Mu3e experiment at PSI); and the coherent neutrinoless conversion of a muon into an electron in the field of a nucleus, $\mu^- N \rightarrow e^- N$ (Mu2e experiment at Fermilab and COMET experiment at J-PARC).
- These “golden” search channels provide complementary sensitivity to new sources of cLFV since the relative rates depend on the details of the underlying new physics model. Thus, it is important to pursue a programme with experiments exploring all three $\mu \rightarrow e$ cLFV transitions to maximize discovery potential, and, in the event of discovery, to help differentiate the various BSM models through a comparison of the rates.
- Current limits for cLFV $\mu \rightarrow e$ transitions are in the $10^{-12} - 10^{-13}$ range and probe effective new physics mass scales above $10^3 \text{ TeV}/c^2$. Next-generation experiments at MEG, Mu3e, Mu2e, and COMET expect to improve these sensitivities by as much as four orders of magnitude on the timescale of the mid-2020s. This dramatic improvement in sensitivity offers genuine discovery possibilities in a wide range of new physics models with SUSY, Extra Dimensions, an extended Higgs sector, lepto-quarks, or those arising from GUT models.
- European contributions are vital to the success of all four of these experiments. Europe hosts two of them (MEG, Mu3e) and provides significant detector components for the others (Mu2e, COMET).
- Beginning in the latter half of the next decade, upgrades to the beamlines at PSI, Fermilab, and J-PARC offer the possibility to further explore this parameter space. Improvements in sensitivity by an additional factor of 10–100 are possible with: a High intensity Muon Beamline (HiMB) at PSI to enable an upgraded Mu3e (Phase-II); the PIP-II linac at Fermilab to enable an upgraded Mu2e (Mu2e-II); an increased intensity at J-PARC to enable an upgraded COMET (Phase-II). A next-generation MEG experiment is also being explored. Like their predecessors, significant European participation in the design, construction, data taking, and analysis will be important to the success of these future endeavors and represents a prudent investment complementary to searches at colliders.
- We urge the committee to strongly support the continued participation of European institutions in experiments searching for cLFV $\mu \rightarrow e$ transitions using high-intensity muon beams at facilities in Europe, the US, and Asia, including possible upgraded experiments at next-generation facilities available in the latter half of the next decade at PSI, Fermilab, and J-PARC.

Objectives

Historically, flavour-changing neutral currents have played a significant role in revealing details of the underlying symmetries at the foundation of the SM. In the SM there is no known global symmetry that conserves lepton flavour. The discoveries of quark mixing and neutrino mixing, each awarded Nobel Prizes, provided profound insights to the underlying physics. Motivated by these past successes, there exists a global programme to explore cLFV processes providing deep, broad probes of BSM physics.

The objective of our programme is to search for evidence of new physics beyond the SM using cLFV processes in the muon sector. These processes offer powerful probes of BSM physics and are sensitive to effective new physics mass scales of $10^3 - 10^4$ TeV/ c^2 , well beyond what can be directly probed at colliders. Over the next five years, currently planned experiments in Europe, the US, and Asia will begin taking data and will extend the sensitivity to cLFV interactions by orders of magnitude. The current experiments each benefit from significant contributions by European institutions. Further improvements are possible and new or upgraded experiments are being considered that would utilize upgraded accelerator facilities at PSI, Fermilab, and J-PARC and could begin taking data in the 2025–2030 timeframe. Strong European participation will be important for the success of these next-generation muon cLFV experiments.

Scientific Context

Flavour violation has been observed in quarks and neutrinos, so it is natural to expect flavour violating effects among the charged leptons as well. In fact, once neutrino mass is introduced, the SM provides a mechanism for cLFV via lepton mixing in loops. However, the rate is suppressed by factors of $(\Delta m_{ij}^2/M_W^2)^2$, where Δm_{ij}^2 is the mass-squared difference between the i^{th} and j^{th} neutrino mass eigenstates, and is estimated to be extremely small, for example $BF(\mu \rightarrow e\gamma) \sim 10^{-54}$ [1]. Many extensions to the standard model predict large cLFV effects that could be observed as new experiments begin data taking over the next five years. Significant improvements are expected across a wide variety of cLFV processes (e.g. $\tau \rightarrow \mu\mu\mu$, $\mu\gamma$, or $e\gamma$; $\mu \rightarrow e\gamma$, eee ; $\mu N \rightarrow eN$; Z or $H^0 \rightarrow e\mu$, $e\tau$, or $\mu\tau$; $K_L \rightarrow e\mu$). The largest improvements are expected in experiments that search for cLFV transitions using muons.

Experimentally, there are three primary muon-to-electron transitions used to search for cLFV¹: a muon decaying into an electron plus a photon, $\mu^+ \rightarrow e^+\gamma$; a muon decaying into three electrons, $\mu^+ \rightarrow e^+e^-e^+$; and direct muon-to-electron conversion via an interaction with a nucleus, $\mu^-N \rightarrow e^-N$. These three $\mu \rightarrow e$ transitions provide complementary sensitivity to new sources of cLFV since the observed rates will depend on the details of the underlying new physics model. For example, for models in which cLFV rates are dominated by γ -penguin diagrams, the $\mu \rightarrow e\gamma$ transition rate is expected to be $\sim 10^2$ times larger than the $\mu \rightarrow eee$ and $\mu N \rightarrow eN$ rates. On the other hand, if the cLFV rates are dominated by Z - or H -penguin diagrams, or if tree level contributions are allowed (e.g. as in some lepto-quark or Z' models), then the $\mu \rightarrow e\gamma$ rate is suppressed and $\mu \rightarrow eee$ and $\mu N \rightarrow eN$ rates can instead be largest. Thus, a programme with experiments exploring all three muon cLFV transitions maximizes the discovery potential and offers the possibility of differentiating among various BSM models by comparing the rates of the three transitions. This is discussed extensively in the literature, see for example references [3] and [4].

Searches for $\mu \rightarrow e$ transitions have been pursued since 1947 when Pontecorvo first searched for the $\mu \rightarrow e\gamma$ process. Since then, the sensitivity has improved by eleven orders of magnitude via a series of increasingly challenging experiments. The current best limits for the three $\mu \rightarrow e$

¹Muonium oscillations, $\mu^+e^- \rightarrow \mu^-e^+$, in which the muon and electron form a bound state, can also be used to set limits on cLFV interactions [2] but are not discussed here.

transitions are $BF(\mu^+ \rightarrow e^+\gamma) < 4.2 \times 10^{-13}$ [5], $BF(\mu^+ \rightarrow e^+e^-e^+) < 1 \times 10^{-12}$ [6], $R_{\mu e}(\text{Au}) < 7 \times 10^{-13}$ [7] at 90% CL, where $R_{\mu e}$ is the $\mu \rightarrow e$ conversion rate normalized to the rate of ordinary muon nuclear capture. Currently planned experiments in Europe, the US, and Asia will provide sensitivities well beyond these existing limits. The MEG experiment at PSI has recently completed an upgrade and expects to extend the $\mu^+ \rightarrow e^+\gamma$ sensitivity by about an order of magnitude with physics data taking beginning in 2019. Further improvements will require a new approach and/or advances in instrumentation. The first phase of the **Mu3e** experiment is under construction at PSI and with about 300 days of data taking is expected to improve the $\mu^+ \rightarrow e^+e^-e^+$ sensitivity by over two orders of magnitude. A second-phase experiment with additional instrumentation could offer a further one order of magnitude improvement with an upgraded muon beam providing $> 2 \times 10^9$ stop- μ^+/s (e.g. a high-intensity muon beam, HiMB, at PSI; or a dedicated μ^+ beamline from PIP-II at Fermilab). The **COMET** experiment under construction at J-PARC will extend the sensitivity to $\mu^-N \rightarrow e^-N$ by about two orders of magnitude by the early-2020s, while the **Mu2e** experiment under construction at Fermilab will extend the sensitivity by about four orders of magnitude by the mid-2020s. On a longer timescale, upgrades in proton intensity offer the possibility of additional improvements. An upgrade to **Mu2e** that extends the sensitivity by another factor of ten or more, **Mu2e-II**, is proposed and would utilize about 100 kW of 0.8 GeV protons from the Fermilab PIP-II linac. An upgrade to **COMET**, **COMET Phase-II**, is proposed and would utilize about 56 kW of 8 GeV protons to reach a comparable sensitivity. The status of the currently planned experiments and their potential for further improvement is discussed in more detail in the next sections.

The outstanding sensitivities that can be achieved by the muon cLFV experiments provide access to new physics mass scales in the $10^3 - 10^4$ TeV/ c^2 range, well beyond what can be directly probed at colliders. In general, these experiments explore the BSM parameter space in a manner complementary to the rest of the HEP experimental programme.

The search for muon-cLFV explicitly probes for flavour-violation in either CP-conserving or CP-violating BSM interactions; in contrast, for instance, to muon g-2 which is sensitive to flavour conserving (and chirality-flipping) interactions. If the Fermilab **Muon g-2** experiment confirms the BNL measurement [8] and hence an a_μ value at odds with the SM beyond 5σ , it will establish the presence of a BSM muon interaction which has obvious ramifications for muon-cLFV, since, in many BSM scenarios, the two are closely related [9]. If the a_μ anomaly disappears, the muon-cLFV experiments are still compelling since they probe effective mass scales well beyond the TeV scale probed by **Muon g-2**.

As the charged counterpart of neutrino oscillations, cLFV plays a significant role in most of the BSM models seeking to explain the neutrino mass hierarchy and the universe's matter anti-matter asymmetry generated through leptogenesis. The cLFV measurements thus have considerable synergy with the neutrinoless double beta decay and neutrino oscillation research programmes. For example, there is a large class of models (see e.g. [10]) proposed to explain the smallness of the neutrino mass. These typically involve extensions to the Higgs sector and the existence of heavier neutrino partners, the properties of which — sterile or non-sterile, Dirac or Majorana, and the mass-scale of the neutrino partners — depend on the model. These heavy neutrino partners typically also play a role in generating a matter anti-matter asymmetry. The majority of these models predict large cLFV effects, and the comparison of cLFV and neutrino measurements together becomes a strong constraint on the model type and its parameters. Indeed, in the most natural models, where the neutrino partners are extremely massive, these measurements are one of the few portals into GUT-scale physics. In the Inverse Seesaw models [11], right-handed neutrinos with masses in the TeV-scale are produced that are potentially observable at the LHC. The present LHC limits are below 1 TeV whereas **Mu2e**, **COMET**, and **Mu3e** will extend this sensitivity to 2 TeV. More generally **Mu2e**, **COMET**, and **Mu3e** still have a sensitivity for RH neutrinos up to masses of a few PeV, well beyond the direct

detection limit of the LHC.

The $\mu \rightarrow e$ experiments also provide complementary information regarding the Majorana nature of neutrinos via the $\mu^- \rightarrow e^+$ transition: $\mu^- N(Z, A) \rightarrow e^+ N(Z - 2, A)$. This transition violates both lepton number and lepton flavour and can only proceed if neutrinos are Majorana. This search channel comes for “free” in the **Mu2e** and **COMET Phase-I** experiments. The **Mu2e** and **COMET** sensitivity to Majorana neutrinos will significantly extend beyond the current best limit [12] with a $\langle m_{e\mu} \rangle$ effective Majorana neutrino mass scale sensitivity down to the MeV region surpassing the $\langle m_{\mu\mu} \rangle$ sensitivity in the kaon sector which is limited to the GeV region [13].

The anomalies in B decays reported by the B-factories and **LHCb** and the **E821** a_μ anomaly have promoted a renewed interest in leptoquarks [14] and Z 's [15]. These models can generate large cLFV effects via tree-level contributions. Direct searches for leptoquarks at **ATLAS** and **CMS** place limits in the 400–800 GeV/ c^2 range which will ultimately increase to approximately 1 TeV/ c^2 with HL-LHC. While there are model-dependencies, the limits from muon cLFV experiments [16] are much stronger with sensitivities up to masses of 300 TeV/ c^2 beyond the present limit (120 TeV/ c^2) established from lepton-flavour violating B-decays [17].

Many experiments will search for the cLFV $\tau \rightarrow \mu$ and $\tau \rightarrow e$ transitions including **ATLAS**, **Belle-II**, **BES-II**, **CMS**, and **LHCb**. In general the existing limits will be extended by about an order of magnitude to the $10^{-9} - 10^{-10}$ range. The proposed **tauFV** experiment [18] may offer another order of magnitude improvement. Thus, the ultimate sensitivity offered by the tau-cLFV searches is several orders of magnitude below the sensitivity offered by the muon cLFV experiments. The relationship between tau-cLFV and muon-cLFV processes is model dependent. The large, close-to maximal, mixing in the neutrino sector favours scenarios in which the rates of cLFV are similar in the two sectors, but other scenarios are also possible in which tau-cLFV rates are significantly enhanced. A comparison of all the transitions: $\mu \rightarrow e$, $\tau \rightarrow e$ and $\tau \rightarrow \mu$ is a very important probe of flavour models. All measurements should be pursued.

In summary, experiments sensitive to violations of lepton flavour, lepton number, and lepton universality play a significant role in the search for BSM physics. It will be necessary to make as broad an array of measurements as possible in order to maximally probe the available parameter space. The muon-cLFV experiments explore cLFV transition rates that are many orders of magnitude beyond what is explored by other experiments and offer sensitivity to new phenomena with mass scales in the few PeV/ c^2 region. Over the next several years, the **MEG**, **Mu3e**, **Mu2e**, and **COMET** experiments have the best reach in their respective channels. Future upgrades could extend the sensitivity another one to two orders of magnitude by utilizing improved accelerator beamlines, and could begin data taking in the 2025–2030 timescale. These future experiments (e.g. **Mu3e Phase-II**, **Mu2e-II**, **COMET Phase-II**, **PRISM**) would offer the most sensitive probes of cLFV for the foreseeable future.

Beam Facilities

The muon-cLFV experiments rely on facilities with high-power proton beams capable of delivering high-intensity muon beams. The experimental infrastructure costs range from €5–50M with additional substantial facility costs. For example the **Mu2e** experiment has a total project cost of \$274M. Several facilities exist with proton beams and transport channels capable of providing muon beams at high intensities. The **PSI** laboratory utilizes 1.4 MW of 590 MeV protons to provide high-intensity beams of secondary particles, including the most intense low-energy muon beams in the world. The $\pi E5$ channel serves the particle physics community and provides positively charged muons with a momentum of 28 MeV/ c , a momentum bite of 5 – 7% FWHM, and rates up to 10^8 stop- μ^+ / s . At **Fermilab** 700 kW of 8.9 GeV/ c protons are available for various experiments, of which about 8 kW will be utilized to produce about

10^{10} stop- μ^-/s for Mu2e. Similarly, at J-PARC about 500 kW of up to 30 GeV protons are available for various experiments, of which about 3 kW (at 8 GeV/c) will be utilized to produce 10^9 stop- μ^-/s for COMET Phase-I. The Fermilab and J-PARC muon beamlines are expected to become operational in the next few years.

Future facilities, capable of providing stopped muon rates a factor of 10–100 larger, are being planned and could become available as early as 2025. These future facilities would enable next-generation muon-cLFV experiments with improved sensitivities. At PSI, strong requests from both the particle physics and material science communities have motivated studies to upgrade the existing muon beamlines (HiMB study). By optimizing the existing M target station and improving the transport efficiency, a new beamline could deliver over 10^{10} stop- μ^+/s for Mu3e Phase-II or a future extension of the MEG experiment. At Fermilab, the long-baseline neutrino programme motivates the need for a significant upgrade of the proton beam intensity. The PIP-II linac will be CW capable and will use superconducting RF technology to provide 1.6 MW of 0.8 GeV protons available for a variety of experiments [19]. Conceptual designs exist to provide about 100 kW of protons to an upgraded Mu2e experiment, Mu2e-II, with over 10^{11} stop- μ^-/s . At J-PARC, plans exist to provide 56 kW to produce 2×10^{11} stop- μ^-/s for COMET Phase-II. Further future, in conjunction with a 1.3 MW upgrade to the Main Ring at J-PARC, the PRISM project would utilize a fixed-field alternating gradient (FFAG) muon storage ring to produce a very intense, very high purity, monochromatic muon beam with the potential to make a whole programme of muon-based measurements at world-class sensitivities.

Methodology

The same basic experimental methodology is employed in searches for all three cLFV $\mu \rightarrow e$ processes. The experiment beamline begins by colliding protons onto a production target to produce low momentum pions. The resulting pions are either transported through a decay volume or directly stopped inside the target, and their decay muons are collected. These experiments require low momentum muons, typically with momenta less than 50 MeV/c, in order to stop them in thin targets at the center of the experimental apparatus. At these low momenta, muons stop in a few mm or less of material. To reach the target sensitivities requires high-intensity muon beams, $> 10^8$ stop- μ/s . The detector apparatus is designed to precisely determine the energy, momentum, and timing of particles originating from the muon stopping target. Because these experiments aim for such extreme sensitivities, their apparatus are customized to the final state of interest.

The $\mu^+ \rightarrow e^+\gamma$ process

The $\mu^+ \rightarrow e^+\gamma$ process is sensitive to new physics mass scales around 10^3 TeV/c² and primarily tests cLFV dipole couplings where new physics appears in loops. The most stringent limit on this process was established by the MEG experiment using data collected from 2009–2013, $BF(\mu^+ \rightarrow e^+\gamma) < 4.2 \times 10^{-13}$ at 90% CL [5]. The MEG II experiment has recently completed construction and aims to improve the sensitivity by an order of magnitude.

The experimental signature of a $\mu^+ \rightarrow e^+\gamma$ decay at rest is given by a back-to-back, photon-positron pair coincident in time and each with an energy of half the mass of the muon. Each event can be described by four observables: the photon and positron energies (E_γ , E_e), their relative direction ($\Theta_{e\gamma}$), and their relative emission time ($t_{e\gamma}$).

The background has two components: an intrinsic physics background coming from the radiative muon decay (RMD), $\mu \rightarrow e\nu\bar{\nu}\gamma$, when the neutrinos carry a small fraction of the available energy; and an accidental background that arises when an energetic positron from a standard muon decay overlaps with an energetic photon from RMD, e^+e^- annihilation-in-flight, or bremsstrahlung. The effective branching fraction for the accidental background is a strong

function of the muon beam intensity, I_μ , and the detector resolutions associated with the four observables, ΔE_γ , ΔE_e , $\Delta\Theta_{e\gamma}$, and $\Delta t_{e\gamma}$:

$$BF_{\text{eff}} \propto I_\mu^2 \times (\Delta E_\gamma)^2 \times \Delta E_e \times (\Delta\Theta_{e\gamma})^2 \times \Delta t_{e\gamma}. \quad (1)$$

In the MEG experiment, which used $I_\mu \sim 3 \times 10^7$ stop- μ^+/s , the accidental background accounted for over 90% of the events near the signal region ($E_\gamma > 48$ MeV). To achieve an improved sensitivity, MEG II will utilize a higher muon beam intensity. Mitigating the accidental background requires MEG II to upgrade its detector components to achieve the required, improved resolutions.

Status and Plans of the MEG II experiment

The MEG II experiment [20] is depicted in Fig. 1 of the Addendum. The main features are an e^+ spectrometer formed by a new cylindrical drift chamber plus precision pixelated timing counters, located inside a superconducting solenoid with a graded magnetic field along the beam axis, and a γ detector, located outside the solenoid, made up of a homogeneous volume of 900 liters of liquid xenon readout in the central region by silicon photomultipliers and in the forward and background region by photomultiplier tubes. The finer granularity of the silicon photomultipliers provides improved γ angular and energy resolution. Additional systems are used to further reduce RMD background, and to monitor the beam quality and stopping target *in situ*. The detector construction is complete and commissioning has begun. Physics data taking is expected to begin in 2019 and to last for a few years. The upgraded detector is expected to provide resolutions roughly a factor of two better than MEG, thus allowing MEG II to utilize the full muon beam intensity available at PSI, $I_\mu \sim 10^8$ stop- μ^+/s , to achieve a factor of ten improvement in expected sensitivity.

Search for $\mu^+ \rightarrow e^+\gamma$ at future facilities

Improvements in sensitivity to the $\mu^+ \rightarrow e^+\gamma$ process beyond the MEG II projection may be possible by utilizing the increased muon intensities that could be available from future facilities at PSI, Fermilab, and J-PARC. However, future experiments would have to devise ways to reduce the accidental background below the 10^{-15} level in order to fully exploit the discovery potential offered by the increased muon intensities.

The use of an active or segmented target could allow a determination of the muon decay vertex, which, in principle, should lead to a strong suppression of the accidental background. Initial studies [21] made for the MEG II project indicated that this additional suppression would be required but this idea could be more effective if different schemes (see below) are adopted. Improvements to ΔE_γ and $\Delta\Theta_{e\gamma}$ resolutions should be the most effective given their quadratic effect on the accidental background. Feasibility studies have been performed for two very different experimental concepts. One is based on a calorimetric detection of the photon, like MEG II, but with improved energy and timing resolutions [22]. The other is based on converting the photon and precisely measuring the trajectories of the resulting e^+e^- pair with a tracking spectrometer [22, 23]. The photon conversion concept is also being studied by Mu3e as a potential extension to its physics programme. These studies promise sensitivities around 10^{-15} at 90% CL after a few years running, but additional studies are required to verify the efficacy of these concepts.

The $\mu^+ \rightarrow e^+e^-e^+$ process

The $\mu^+ \rightarrow e^+e^-e^+$ process is sensitive to new physics at mass scales beyond 10^3 TeV/ c^2 and probes cLFV couplings that arise from dipole interactions where new physics appears in loop diagrams and from μeee contact interactions. The most stringent limit on this process was established by the SINDRUM experiment using data collected from 1983–1986, $BF(\mu^+ \rightarrow e^+e^-e^+) < 1 \times 10^{-12}$ at 90% CL [6]. Any improvement in the sensitivity beyond

this has a significant impact on models predicting cLFV, especially the associated four-fermion couplings. The Mu3e collaboration aims for a sensitivity of $BF(\mu^+ \rightarrow e^+e^-e^+) < 5 \times 10^{-15}$ at 90% CL in a first phase, Mu3e Phase-I, using the existing π E5 beamline at PSI. A further improvement is possible in a second phase, Mu3e Phase-II, with upgrades to the muon beam (HiMB project) and detector to reach a sensitivity that is four orders of magnitude better than the current experimental limit.

The experimental signature of a $\mu^+ \rightarrow e^+e^-e^+$ decay at rest is given by three charged particle tracks, corresponding to the $e^+e^-e^+$ decay products, coincident in time, originating from a common vertex, and with a total energy consistent with the mass of the muon. Since this is a three-body decay, the energy of the decay products ranges from < 1 MeV up to a maximum of about half the mass of the muon.

The background has two main components: an intrinsic physics background coming from radiative muon decays (RMD) when the photon converts to an e^+e^- pair; and an accidental background from the random combination of electrons and positrons from separate decays. The RMD background can be kept sufficiently small if the resolution on the $e^+e^-e^+$ energy sum can be kept below about 1 MeV, while the suppression of the accidental background additionally relies on excellent timing and vertex resolution.

Status and Plans of the Mu3e experiment

The Mu3e Phase-I experiment [24] is depicted in Fig. 2 of the Addendum. The main features include precision particle tracking with ultra-light, monolithic, silicon pixel tracking layers based on the HV-MAPS technology [25] cooled in an innovative manner using helium gas, plus a system of scintillating fibers and tiles to provide a time resolution below the sub-nanosecond level, all located inside a superconducting solenoid providing a constant 1 T field along the beam axis. A farm of GPUs will use a highly parallel track-fitting algorithm to perform full online tracking of all data as it streams continuously from the detector front-ends. The detector is capable of handling very high rates and the Mu3e Phase-I sensitivity will be limited by the muon rate that PSI can deliver. A new Compact Muon Beamline has been installed for Mu3e and will deliver a continuous high-purity beam of 28 MeV/c muons at a rate of about 10^8 stop- μ/s . The detector design is advanced and prototypes of all the main components have been built and successfully tested. Construction of the Mu3e Phase-I experiment will begin in 2019 and a first commissioning run is expected in 2020. After three years of operation the projected Phase-I sensitivity is $BF(\mu^+ \rightarrow e^+e^-e^+) < 5 \times 10^{-15}$ at 90% CL. However, the Mu3e experimental concept allows for further significant improvements in sensitivity.

Search for $\mu^+ \rightarrow e^+e^-e^+$ at future facilities

The sensitivity of Mu3e Phase-I is largely limited by the muon rate. The sensitivity to $\mu^+ \rightarrow e^+e^-e^+$ could be improved by more than an order of magnitude with a modest extension of the detector and access to a higher intensity muon beam. Roughly a factor two improvement in experimental acceptance is expected by extending the instrumentation in the forward and backward directions using the same pixel and scintillator technologies planned for Phase-I. At PSI, concepts for a new High Intensity Muon Beamline, HiMB, have been investigated. Recent studies indicate that by refurbishing target M, and by installing a new capture solenoid the muon rate could be increase up to 1.3×10^{10} stop- μ^+/s [26], which is more than sufficient for Phase-II. Future facilities at Fermilab and J-PARC could in principle provide similar muon intensities. If approved, the earliest HiMB could be installed is 2024. After three years of data taking at the increased muon intensity the projected Phase-II sensitivity is $BF(\mu^+ \rightarrow e^+e^-e^+) < 1 \times 10^{-16}$ at 90% CL.

Detector R&D should also continue to further improve the time resolution and to reduce the amount of detector material, both being important requirements for suppressing accidental

backgrounds at higher rates. Silicon pixel detectors with picosecond timing represent a very promising technology for future Mu3e upgrades.

The $\mu^- N \rightarrow e^- N$ process

The $\mu^- N \rightarrow e^- N$ process is sensitive to new physics at mass scales up to 10^4 TeV/ c^2 and probes cLFV couplings that arise from interactions where new physics appears in loop diagrams as well as from μeqq contact interactions. The most stringent limit on this process was established by the SINDRUM-II experiment using data collected in 2000, $R_{\mu e}(\text{Au}) < 7 \times 10^{-13}$ at 90% CL [7], where, by convention, $R_{\mu e}$ is the rate of the $\mu^- N \rightarrow e^- N$ conversion process normalized to the normal muon nuclear capture process, $\mu^- N(A, Z) \rightarrow \nu N^*(A, Z - 1)$. Significant improvements in sensitivity offer genuine discovery possibilities across many new physics models, including several to which it is uniquely sensitive. The COMET Phase-I experiment is currently under construction and aims to improve sensitivity by a factor of 100 starting in the next few years. The Mu2e experiment is under construction and aims to improve the sensitivity by four orders of magnitude by the mid 2020s, while COMET Phase-II is a proposed upgrade to COMET Phase-I that would achieve a similar sensitivity or better. Further improvements are possible for both experiments with upgrades to the beamline and detectors.

The direct conversion process, $\mu^- N \rightarrow e^- N$, is dominated by coherent interactions with the nucleus to provide a two-body final state yielding a clean experimental signature: an outgoing electron with an energy near the muon mass (the recoil nucleus is not directly observed). The time distribution of signal electrons should be consistent with the characteristic lifetime of the muonic atoms formed as the initial μ^- beam comes to rest in a stopping target.

Significant sources of background events can arise from muons that decay-in-orbit (DIO) — that is, decay while captured in the atomic orbit around a nucleus in the stopping target, from pions that survive to the stopping target, and from cosmic ray muons that decay-in-flight or interact in material to produce an electron with an energy near the muon mass. If the energy of the initial proton beam is above the anti-proton production threshold, then annihilations of the anti-protons can contribute an additional source of background. The steeply falling DIO background can be mitigated with excellent momentum resolution; the pion background can be suppressed by using a pulsed proton beam and employing a delayed live gate; the cosmic ray background can be removed using a high-efficiency cosmic veto system; and anti-proton backgrounds can be kept small by using absorbers to range out the anti-protons far away from the stopping target.

Both the COMET and the Mu2e experiments are based on a very clever idea first proposed by Lobashov and Djilkabaev in 1989 [27]. A system of three solenoids — a pion capture solenoid, a muon transport solenoid, and a detector solenoid — with graded magnetic fields which provide a significantly improved muon beam intensity to enable dramatic improvements in sensitivity relative to SINDRUM-II. The feasibility of this method has been demonstrated at low intensity at the MuSIC facility at the Research Center for Nuclear Physics in Osaka [28].

Status and Plans of the COMET experiment

The COMET Phase-I experiment [29] is depicted in Fig. 4 of the Addendum. The apparatus begins with a pion-production target made of graphite located inside the pion capture solenoid, which provides a graded magnetic field to collect low-momentum pions by reflecting them backwards with respect to the incoming proton beam. The muon transport solenoid is a curved 90-degree magnet that, together with a set of dipole coils, serves as a transport and charge- and momentum-selection channel for $\pi^- \rightarrow \mu^- \nu$ decays. The muon transport solenoid delivers a high-intensity μ^- beam to the detector solenoid, which houses an aluminum stopping target and active detector elements, including a cylindrical drift chamber (CDC) for the $\mu^- N \rightarrow e^- N$ search and low-mass straw chambers and a fast LYSO crystal calorimeter for

beam measurements. An active cosmic-ray veto system shadows the detector and stopping target regions outside the solenoid volume. Additional instrumentation monitors the proton and muon beams. The construction of the entire apparatus is at an advanced stage, with two of the magnets and the CDC complete, including significant European contributions to the cosmic-veto, muon stopping target and beam monitoring systems, the trigger and DAQ, and computing and software. Beam commissioning is expected to begin in 2020. The COMET Phase-I experiment will utilize about 3 kW of 8 GeV protons from the J-PARC Main Ring, delivered in pulses spaced by $1.17 \mu\text{s}$, to first make important measurements of the muon yield and determine rates for various background processes before concentrating on a search for cLFV. After 150 days of operation the projected COMET Phase-I sensitivity is $R_{\mu e} < 7 \times 10^{-15}$ at 90% CL [29]. This sensitivity can be significantly improved using a higher intensity proton beam and extending the COMET Phase-I apparatus.

Status and Plans of the Mu2e experiment

The Mu2e experiment [30] is depicted in Fig. 6 of the Addendum. The graded high-field pion production solenoid collects and focuses low-momentum pions towards the muon transport solenoid, which is an “S”-shaped magnet with a total path length of about 13 meters. The muon transport solenoid includes a set of collimators for momentum and charge-selection to provide $\sim 10^{10}$ stop- μ^-/s using 8 kW of 8 GeV protons from the Fermilab Booster delivered in pulses spaced $1.7 \mu\text{s}$ apart. The detector solenoid provides a graded magnetic field in the upstream region, which houses the stopping target, and a near constant magnetic field in the downstream region, which houses the active detector elements. A low-mass tracking system consisting of approximately 21k thin aluminized-mylar straws [31] and a calorimeter consisting of two annular disks of pure CsI crystals [32] precisely measure the timing, energy, and momenta of particles originating from the stopping target. The apparatus is shadowed on the outside by a large, scintillator-based cosmic-veto system. Ancillary systems are used to monitor the quality and intensity of the proton and muon beams. Construction of the solenoids and all the detector sub-systems has begun, with significant European contributions to the muon transport solenoid, the calorimeter, and the muon beam monitoring system. Commissioning is expected to begin in 2022. After 690 days of operation the projected sensitivity is $R_{\mu e} < 8 \times 10^{-17}$ at 90% CL [30]. This sensitivity can be improved by at least a factor of 10 using a higher intensity proton beam and upgrading the Mu2e apparatus.

Search for $\mu^- N \rightarrow e^- N$ at future facilities

The COMET Phase-I experiment can be extended, as depicted in Fig. 5 of the Addendum, and utilize 56 kW of 8 GeV protons from the J-PARC Main Ring, delivered in pulses spaced by $1.17 \mu\text{s}$, to reach a sensitivity of $R_{\mu e} < 2.6 \times 10^{-17}$ at 90% CL with about 230 days of operation. Further improvements by one order of magnitude from refinements to the experimental design and operation are being considered, within the beam power and the beam time as originally assumed [33]. These improvements include dipole steering fields in the curved muon transport and electron spectrometer sections to allow a more fine-tuned momentum selection which is important to optimize the acceptance and background rejection. The detailed measurements from Phase-I will provide important input to the final Phase-II design and construction. Data taking could begin in the mid-2020s.

The Mu2e experiment can be upgraded, Mu2e-II, to take advantage of the increased proton beam intensity available from the PIP-II project, currently in the design phase at Fermilab. The PIP-II linac is expected to become operational in the latter half of the 2020s and will provide 1.6 MW of 0.8 GeV protons with a programmable time structure. An Expression of Interest for Mu2e-II [34] was recently submitted to the Fermilab Physics Advisory Committee, which concluded that the science case was compelling and recommended that funding for high-priority R&D be identified. The Expression of Interest included signatures from 130 scientists

from 36 institutions in six countries, including Italy, Germany, and the UK. Using 100 kW of protons from PIP-II, the Mu2e-II projected sensitivity is a factor ten or more better than the Mu2e sensitivity. Data taking could begin in the late 2020s.

The COMET collaboration is also heavily involved in R&D towards the PRISM project, which combines COMET Phase-II with an FFAG muon storage ring to potentially provide muon beam intensities of $> 10^{12}$ stop- μ/s with a narrow momentum bite allowing the use of very thin stopping targets, and significantly reduced pion contamination owing to the increased transport path length. In conjunction with an upgrade to the J-PARC proton source to achieve 1.3 MW and to the detector systems to accommodate the higher rates, PRISM offers the potential to achieve sensitivities to $\mu^- N \rightarrow e^- N$ of the order of 10^{-19} . The monochromatic, pion-suppressed, high-intensity muon beam provided by PRISM will allow the use of stopping targets comprised of heavy elements, such as gold or lead, that can be important in understanding the underlying new physics operators in the event of a discovery [33].

Summary

The MEG, Mu3e, Mu2e, and COMET experiments use intense muon beams to provide the broadest, deepest, most sensitive probes of charged-lepton flavour violating interactions and to explore the BSM parameter space with sensitivity to new physics mass scales of $10^3 - 10^4$ TeV/ c^2 , well beyond what can be directly probed at colliders. Over the next five years, currently planned experiments in Europe, the US, and Asia will begin taking data and will extend the sensitivity to $\mu \rightarrow e$ charged-lepton flavour violating transitions by orders of magnitude. Further improvements are possible and new or upgraded experiments are being considered that would utilize upgraded accelerator facilities at PSI, Fermilab, and J-PARC. The schedule of planned and proposed experiments is summarized in the figure below. Strong European participation in the design, construction, data taking, and analysis will be important for the success of these future endeavors and represents a prudent investment complementary to searches at colliders.

We urge the committee to strongly support the continued participation of European institutions in experiments searching for charged-lepton flavour violating $\mu \rightarrow e$ transitions using high-intensity beams at facilities in Europe, the US, and Asia, including possible upgraded experiments at next-generation facilities available the latter half of the next decade at PSI, Fermilab, and J-PARC.




Figure 1: Planned data taking schedules for current experiments that search for charged-lepton flavor violating $\mu \rightarrow e$ transitions. Also shown are possible schedules for future proposed upgrades to these experiments. The current best limits for each process are shown on the left in parentheses, while expected future sensitivities are indicated by order of magnitude along the bottom of each row.

References

- [1] S.T. Petcov, Sov. J. Nucl. Phys. 25 (1977) 340.
- [2] L. Willmann, *et al.*, Phys.Rev.Lett. 82 (1999) 49.
- [3] L. Calibbi and G. Signorelli, Riv. Nuovo. Cimento, 41 (2018) 71.
- [4] V. Cirigliano, *et al.*, Phys. Rev. D80 (2009) 013002.
- [5] A. Baldini, *et al.* (MEG Collaboration), Eur. Phys. J. C76 (2016) 434.
- [6] U. Bellgardt, *et al.* (SINDRUM Collaboration), Nucl. Phys. B299 (1988) 1.
- [7] W. Bertl, *et al.* (SINDRUM-II Collaboration), Eur. Phys. J. C47 (2006) 337.
- [8] G.W. Bennett, *et al.* (E821 Collaboration), Phys. Rev. Lett. 92 (2004) 161802.
- [9] G. F. Giudice, P. Paradisi and M. Passera, JHEP11 (2012) 113.
- [10] T. Hambye, Proc. Nucl. Phys. B248 (2014) 13.
- [11] A. Abada, *et al.*, JHEP 11 (2014) 048.
- [12] J. Kaulard, *et al.* (SINDRUM-II Collaboration), Phys. Lett. B422 (1998) 334.
- [13] B. Yeo, Y. Kuno, M. Lee and K. Zuber, Phys.Rev. D96, no. 7 (2017) 075027.
- [14] B. Gripaios, *et al.*, JHEP (2015) 6; B. Dumont, *et al.*, arXiv:1603.05248 (2016); M. Bauer and M. Neubert, Phys. Rev.Lett. 116 (2016) 141802; S. Baek and K. Nishiwaki, Phys. Rev. D93 (2016) 015002.
- [15] A. Crivellin, *et al.*, Phys. Rev. D92 (2015) 050413.
- [16] J. Arnold, *et al.*, Phys. Rev. D88 (2013) 035009.
- [17] R. Aaij, *et al.* (LHCb Collaboration), Phys.Rev.Lett. 111 (2013) 141801.
- [18] G. Wilkinson *et al.*, <https://indico.cern.ch/event/706741/contributions/3017537>
- [19] M.Ball, *et al.* (PIP-II Accelerator Facility),
<http://pip2-docdb.fnal.gov/cgi-bin/ShowDocument?docid=113> (2018).
- [20] A. Baldini, *et al.* (MEG II Collaboration), Eur. Phys. J. C78 (2018) 380.
- [21] A. Papa, *et al.*, Nucl. Phys. Proc. Suppl. 248 (2014) 121.
- [22] G. Cavoto, *et al.*, Eur.Phys.J. C78 (2018) 37.
- [23] C. Cheng, B. Echenard, D.G. Hitlin, arXiv:1309.7679 (2013).
- [24] A. Blondel, *et al.* (Mu3e Collaboration), arXiv:1301.6113 (2013).
- [25] I. Peric, Nucl. Instrum. Meth. A582, 876 (2007).
- [26] A. Papa, NuFact 2018, Blacksburg, Virginia USA,
<https://indico.phys.vt.edu/event/34/contributions/701>
- [27] R.M. Dzhilkibaev and V.M. Lobashev, Sov.J.Nucl.Phys. 49(2), (1989) 384.
- [28] S. Cook, *et al.*, Phys. Rev. Accel. Beams 20(3), (2017) 030101.
- [29] R. Abramishvili, *et al.* (COMET Collaboration),
http://comet.kek.jp/Documents_files/PAC-TDR-2016/COMET-TDR-2016_v2.pdf
- [30] L. Bartoszek, *et al.* (Mu2e Collaboration), arXiv:1501.05241 (2015).
- [31] M. Lee (on behalf of Mu2e Collaboration), Nucl. Part. Phys. Proc., 273 (2016) 2530.
- [32] N. Atanov, *et al.*, IEEE Trans. Nucl. Sci., Vol 65, N. 8, (2018) 2073.
- [33] COMET submission to the European Strategy for Particle Physics 2020, and references therein.
- [34] F. Abusalma, *et al.* (Mu2e-II Experiment), arXiv:1802.02599 (2018).