


# SpecMAT

## Spectroscopy of exotic nuclei in a Magnetic Active Target


Riccardo Raabe  
KU Leuven, Instituut voor Kern- en Stralingsfysica



# The active-target method

## Time-Projection Chamber (TPC) + gas is the target

- Full 3D track reconstruction
- High luminosity, keeping energy resolution
- Energy of stopped particles  
Particle identification
- Versatile:  
different gases, pressures, configurations  
ancillary detectors



LOI I-119

**Letter of Intent to the  
ISOLDE and Neutron Time-of-Flight Experiments Committee  
for experiments with HIE-ISOLDE**

**Direct and resonant reactions using an Active Target**



# The active-target method

## Time-Projection Chamber (TPC) + gas is the target

- Full 3D track reconstruction
- High luminosity, keeping energy resolution
- Energy of stopped particles  
Particle identification
- Versatile:  
different gases, pressures, configurations  
ancillary detectors

### SpecMAT:

- Magnetic field parallel to the beam direction
- Drift field parallel to magnetic field




# The active-target method

## $\gamma$ -ray detection

- CeBr<sub>3</sub> scintillators:  
good efficiency and resolution
- Si photomultipliers

J. Diriken et al., PLB 736, 533 (2014)

J. Diriken et al., PRC 91, 054321 (2015)




# Physics goals

- What are the **forces driving the shell structure in nuclei** and how do they change in nuclei far from stability?
- What remains of the  **$Z = 28$  and  $N = 50$  “magic numbers”** in  $^{78}\text{Ni}$ ?
- Do we understand **shape coexistence in nuclei**, and what are the mechanisms controlling its appearance?

## Changes in nuclear structure far from stability

- Shell evolution towards  $^{78}\text{Ni}$
- Shape coexistence “west” of  $^{208}\text{Pb}$



# Shell evolution towards $^{78}\text{Ni}$

- Migration of  $\pi f_{7/2}$ ,  $\pi f_{5/2}$  as  $\nu g_{9/2}$  is filled (tensor interaction)

68Zn	69Zn	70Zn	71Zn	72Zn	73Zn	74Zn	75Zn	76Zn	77Zn	78Zn	79Zn	80Zn	81Zn
67Cu	68Cu	69Cu	70Cu	71Cu	72Cu	73Cu	74Cu	75Cu	76Cu	77Cu	78Cu	79Cu	80Cu
66Ni	67Ni	68Ni	69Ni	70Ni	71Ni	72Ni	73Ni	74Ni	75Ni	76Ni	77Ni	78Ni	
65Co	66Co	67Co	68Co	69Co	70Co	71Co	72Co	73Co	74Co	75Co			





FIG. 3. Energy of the lowest levels from experiment [2,5,6] compared to large-scale shell-model calculation [25].

K. Flanagan et al.,  
 PRL 103 (2009) 142501

# Shell evolution towards $^{78}\text{Ni}$

- Migration of  $\pi f_{7/2}$ ,  $\pi f_{5/2}$  as  $\nu g_{9/2}$  is filled (tensor interaction)

68Zn	69Zn	70Zn	71Zn	72Zn	73Zn	74Zn	75Zn	76Zn	77Zn	78Zn	79Zn	80Zn	81Zn
67Cu	68Cu	69Cu	70Cu	71Cu	72Cu	73Cu	74Cu	75Cu	76Cu	77Cu	78Cu	79Cu	80Cu
66Ni	67Ni	68Ni	69Ni	70Ni	71Ni	72Ni	73Ni	74Ni	75Ni	76Ni	77Ni	78Ni	
65Co	66Co	67Co	68Co	69Co	70Co	71Co	72Co	73Co	74Co	75Co			



K. Flanagan et al.,  
 PRL 103 (2009) 142501

Proposal P-495:  $^{72,74,76}\text{Zn}(d, ^3\text{He})$   
 to directly measure the  $f_{7/2}$ - $f_{5/2}$  gap



LOI I-191

Letter of Intent to the ISOLDE and Neutron Time-of-Flight Committee  
 (Following HIE-ISOLDE proposal INTC-P-495)

Single-particle proton states in  $^{69}\text{Cu}$

# Shell evolution and shape coexistence

Y Tsunoda, T Otsuka et al., PRC 89 (2014) 031301





$^{68}\text{Ni}$




# Shell evolution and shape coexistence

Y Tsunoda, T Otsuka et al., PRC 89 (2014) 031301






$^{68}\text{Ni}$

T Otsuka and Y Tsunoda, JPG 43 (2016) 024009



# The n-deficient Pb region



B A Marsh et al., Nature Physics (2018)



# The n-deficient Pb region

## Transfer reactions in Hg

- $^{185g,m}\text{Hg}$  (d,p), (p,d), ( $\alpha$ , $^3\text{He}$ )
- Beam intensity  $\approx 10^5$  pps  $\rightarrow$  feasible!





Data: NNDC, figure courtesy of Liam Gaffney  
 Original figure in R. Julin et al.,  
 J. Phys. G 27 (2001) R109

# The n-deficient Pb region

## Transfer reactions in Hg

- $^{185g,m}\text{Hg}$  (d,p) and (p,d)
- Beam intensity  $\approx 10^5$  pps  $\rightarrow$  feasible!



### LOI I-195

Letter of Intent to the ISOLDE and Neutron Time-of-Flight Committee

Investigating single-particle configurations  
in deformed Hg and Cd isotopes

Data: NNDC, figure courtesy of Liam Gaffney  
 Original figure in R. Julin et al.,  
 J. Phys. G 27 (2001) R109

# Study of the Pigmy Dipole Resonance

## Low-energy dipole strength

- First observation in 1961  
 $\gamma$  rays from neutron capture  
 G.A. Bartholomew, Annu. Rev. Nucl. Sci. 11 (1961) 259
- First use of “pygmy resonance” (PDR)  
 J.S. Brzosko et al., Can. J. Phys 47 (1969) 2849
- Description as a collective excitation  
 Mohan et al., Phys. Rev. C 3 (1971) 1740  
 “Three-Fluid Hydrodynamical Model of Nuclei”:  
 Neutron excess oscillates against the N=Z core
- Different experimental probes  
 to investigate the isospin nature of these states






Figure A. Bracco et al.,  
 Eur. Phys. J. A 51 (2015) 99  
 Data from K. Govaert et al.,  
 Phys. Rev. C 57 (1998) 2229  
 and J. Endres et al.,  
 Phys. Rev. C 85 (2012) 064331

# Study of the Pigmy Dipole Resonance


D. Vretenar et al., J. Phys. G 35 (2008) 014039



Core in phase  
+neutron skin

# Study of the Pigmy Dipole Resonance

- Isoscalar probe:  
( $\alpha, \alpha'$ ) inelastic scattering
- SpecMAT:  
 $\gamma$  rays from decay of bound states



LOI I-194


Letter of Intent to the ISOLDE and Neutron Time-of-Flight Committee  
 Study of the Pygmy Dipole Resonance using an Active Target

- Cases:  $^{90}\text{Sr}$ ,  $^{194}\text{Hg}$ ,  $^{146,148,150}\text{Gd}$   
 $\approx 10$  MeV/nucleon

# Status

Design: O. Poleshchuk, KU Leuven

- GET Electronics (2048 channels):  
purchased and commissioned
- Pad plane and field cage  
designed in CERN (EP-DT-DD MPT)  
to be ordered shortly
- Chamber finalized  
and ordered
- Scintillators + SiPMs purchased,  
commissioning ongoing  
48x48x48 mm<sup>3</sup>  
Photopeak efficiency 8% at 1 MeV  
Nominal resolution 3.9% at 661 keV  
No degradation in 3-T magnetic field






# Status

Design: O. Poleshchuk, KU Leuven


- GET Electronics (2048 channels):  
purchased and commissioned
- Pad plane and field cage  
designed in CERN (EP-DT-DD MPT)  
to be ordered shortly
- Chamber finalized  
and ordered
- Scintillators + SiPMs purchased,  
commissioning ongoing  
48x48x48 mm<sup>3</sup>  
Photopeak efficiency 8% at 1 MeV  
Nominal resolution 3.9% at 661 keV  
No degradation in 3-T magnetic field



# Status

- GET Electronics (2048 channels): purchased and commissioned
- Pad plane and field cage designed in CERN (EP-DT-DD MPT) to be ordered shortly
- Chamber finalized and ordered
- Scintillators + SiPMs purchased, commissioning ongoing  
48x48x48 mm<sup>3</sup>  
Photopeak efficiency 8% at 1 MeV  
Nominal resolution 3.9% at 661 keV  
No degradation in 3-T magnetic field



Design: O. Poleshchuk, KU Leuven



# Status

- GET Electronics (2048 channels): purchased and commissioned
- Pad plane and field cage designed in CERN (EP-DT-DD MPT) to be ordered shortly
- Chamber finalized and ordered
- Scintillators + SiPMs purchased, commissioning ongoing  
 $48 \times 48 \times 48 \text{ mm}^3$   
 Photopeak efficiency 8% at 1 MeV  
 Nominal resolution 3.9% at 661 keV  
 No degradation in 3-T magnetic field

Design: O. Poleshchuk, KU Leuven




Simulations for:  
 electric field, mechanical stress, gas flow

# Status


- GET Electronics (2048 channels): purchased and commissioned
- Pad plane and field cage designed in CERN (EP-DT-DD MPT) to be ordered shortly
- Chamber finalized and ordered
- Scintillators + SiPMs purchased, commissioning ongoing  
 48x48x48 mm<sup>3</sup>  
 Photopeak efficiency 8% at 1 MeV  
 Nominal resolution 3.9% at 661 keV  
 No degradation in 3-T magnetic field

Design: O. Poleshchuk, KU Leuven



# Installation

- Installation through a platform and rails from the back of the ISS
- Cables (8m) running to the electronics racks outside the cage
- To be coordinated with the installation of the new ISS array



# Planning

## January – March 2019: characterisation of the full system

- Characterisation of the  $\gamma$ -ray array
- Leak tests of the detector chamber
- Commissioning of the gas control system
- Tests of the electric field cage
- Characterisation of particle tracks

## 2019 Move to CERN

- Safety clearance
- Installation



# Summary

## Thanks to the SpecMAT team!

*A. Arokja Raj, H. De Witte, O. Poleshchuk, J. Refsgaard, M. Renaud, J. Yang*  
*With us in the past: S. Ceruti, M. Babo, T. Marchi, C. Swartz*

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement n. 617156