


Variation of Reactor Antineutrino Yield at RENO

Hyunkwan Seo for the RENO Collaboration
Seoul National University

XVIII International Workshop on Neutrino Telescopes
Palazzo Franchetti, Istituto Veneto, Venezia, Italy, March. 18-22, 2019



RENO Collaboration



Reactor Experiment for Neutrino Oscillation

(9 institutions and 40 physicists)



- Chonnam National University
- Dongshin University
- GIST
- Gyeongsang National University
- KAIST
- Kyungpook National University
- Seoul National University
- Seoyeong University
- Sungkyunkwan University

- Total cost : **\$10M**
- Start of project : **2006**
- The first experiment running with both near & far detectors from **Aug. 2011**

YongGwang (靈光) :



RENO Experimental Set-up



New Results from RENO

- Precise measurement of $|\Delta m_{ee}^2|$ and θ_{13} using ~2200 days of data (Aug. 2011 – Feb 2018)



“Measurement of Reactor Antineutrino Oscillation Amplitude and Frequency at RENO” (Phys. Rev. Lett. 121, 201801 (2018. 11. 15))

- Fuel-composition dependent reactor antineutrino yield

“Fuel-composition dependent reactor antineutrino yield at RENO”
(submitted to PRL (arXiv: 1806.00574))

Measured Spectra of IBD Prompt Signal

Clear excess at 5 MeV




Near Live time = 1807.88 days
of IBD candidate = 850,666
Background : $2.03 \pm 0.06\%$


Far Live time = 2193.04 days
of IBD candidate = 103,212
Background : $4.76 \pm 0.20\%$

Far/Near Shape analysis

Energy-dependent disappearance of reactor antineutrinos




Results of θ_{13} and $|\Delta m_{ee}^2|$



$$\sin^2 2\theta_{13} = 0.0896 \pm 0.0048(\text{stat.}) \pm 0.0047(\text{syst.}) \quad (\pm 7.6\%)$$



$$|\Delta m_{ee}^2| = 2.68 \pm 0.12(\text{stat.}) \pm 0.07(\text{syst.}) (\times 10^{-3} \text{ eV}^2) \quad (\pm 5.2\%)$$

Observed L/E Dependent Oscillation



$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \approx 1 - \sin^2 2\theta_{13} \sin^2 \left(\Delta m_{ee}^2 \frac{L}{4E_\nu} \right)$$

Comparison of θ_{13} and $|\Delta m_{ee}^2|$



Motivation for the study of fuel composition dependent reactor antineutrino yield

Reactor Antineutrino Anomaly

- ~6% deficit of measured reactor neutrino flux compared to the prediction with new predicted flux evaluation in 2011 by Huber and Mueller.
- Deficit of observed reactor neutrino fluxes relative to the prediction (Huber + Mueller model) indicates an overestimated flux or possible oscillation to sterile neutrinos.



The possibility that reactor anomaly is due to miscalculation of one or more of the ^{235}U , ^{239}Pu , ^{238}U and ^{241}Pu antineutrino fluxes is investigated by observing **fuel-composition dependent variation of reactor antineutrino yield and spectrum.**

C. Giunti, Phys. Lett. B 764, 145 (2017)


F. P. An et al. (Daya Bay Collaboration), PRL 118, 251801 (2017)

RENO Collaboration, arXiv:1806.00574 (submitted to PRL)


Reactor for Antineutrino Source

Reactor: A copious and isotropic source of electron antineutrinos

$\sim 3 \text{ GW}_{\text{th}}$ or $\sim 1 \text{ GW}_{\text{elec}}$ per reactor




3 GW_{th} reactor
 $\rightarrow \sim 6 \times 10^{20} \bar{\nu}_e/\text{sec}$



- 3-4% accurate neutrino source
- 0.13% uncertainty of IBD cross section


[* P. Huber, Phys. Rev. C84, 024617 (2011)
T. Mueller *et al.*, Phys. Rev. C83, 054615 (2011)]

Evolution of Fuel Isotope Fraction in Single Reactor




The **Fission fraction** of an isotope varies with **fuel-burning** ¹²

Observable Reactor Neutrino Spectrum




Evolution of Fuel Composition at RENO

Average fission fraction $f_{235}: f_{239}: f_{238}: f_{241} = 0.573 : 0.299 : 0.073 : 0.055$



Effective fission fraction of ^{235}U
(weighted by each reactor's thermal power and baseline)



8 groups of near IBD samples with equal statistics according to ^{235}U isotope fraction

Effective Fission fraction for each isotope

$$F_i(t) = \frac{\sum_{r=1}^6 \frac{W_{th,r}(t)\bar{p}_r(t)f_{i,r}(t)}{L_r^2\bar{E}_r(t)}}{\sum_{r=1}^6 \frac{W_{th,r}(t)\bar{p}_r(t)}{L_r^2\bar{E}_r(t)}}$$

Predicted IBD Yield per Fission

IBD yield per fission for each isotope


(Total # of produced IBD events)

$$y_i = \int \sigma(E_\nu) \phi_i(E_\nu) dE_\nu$$

IBD cross section Antineutrino spectrum
(H-M model)

(i : each isotope)

(H-M model)



Average IBD yield per fission

(for each 8 group, j)

$$\bar{y}_{f,j} = \sum_{i=1}^4 \bar{F}_{i,j} y_i$$

$\bar{F}_{i,j}$: Effective Fission fraction for each isotope

	H-M model ($10^{-43} \text{ cm}^2/\text{fission}$)
y_{235}	6.70 +/- 0.14
y_{239}	4.38 +/- 0.11
y_{238}	10.07 +/- 0.82
y_{241}	6.07 +/- 0.13

Measurement of IBD Yield per Fission using RENO data

Measurement of IBD yield per fission ($\bar{y}_{f,j}$) for each group

- Number of IBD events after subtracting background is obtained for each group, j . Then $\bar{y}_{f,j}$ is determined by solving the following equation.

$$N_j = \bar{y}_{f,j} \sum_{r=1}^6 \frac{N_p}{4\pi L_r^2} \int dt \left[\frac{W_{th,r}(t) \bar{P}_{sur}^r(t)}{\sum_i f_{i,r}(t) E_i} \right] \epsilon_d(t)$$

of Observed IBD # of Target proton # of fission Detection Efficiency

‘Measured IBD yield per fission’ corresponding to average IBD yield per fission

Average IBD yield per fission

(for each 8 group, j)

$$\bar{y}_{f,j} = \sum_{i=1}^4 \bar{F}_{i,j} \gamma_i \quad \bar{F}_{i,j} : \text{Effective Fission fraction for each isotope}$$


Fuel-Composition Dependent Reactor Neutrino Yield

Measured total averaged IBD yield per fission (\bar{y}_f)

$$= (5.84 \pm 0.13) \times 10^{-43} \text{ cm}^2/\text{fission}$$

Ratio (Data / H-M model) for the total average IBD yield

$$= 0.940 \pm 0.021 \rightarrow (6.0 \pm 2.1)\% \text{ deficit}$$



Averaged IBD yield per fission (\bar{y}_f) vs $\bar{F}_{i,j}$


→ slope means **different neutrino yield** for each isotope

→ rules out the no fuel-dependent variation at **6.6 σ**

The scaled model indicates the **reactor antineutrino anomaly**

Measurement of y_{235} and y_{239}

The best-fit measured yields per fission of ^{235}U and ^{239}Pu



The best-fit values

$$y_{235} = 6.15 \pm 0.19 \text{ (} 2.8\sigma \text{ deficit)}$$

$$y_{239} = 4.18 \pm 0.26 \text{ (} 0.8\sigma \text{ deficit)}$$

H-M model

$$y_{235} = 6.70 \pm 0.14$$


$$y_{239} = 4.38 \pm 0.11$$

Reevaluation of the y_{235} may **mostly solve** the reactor antineutrino **anomaly**.

But ^{239}Pu is **not entirely** ruled out as a possible source of the anomaly.

Correlation of 5 MeV excess with fuel ^{235}U

2.9 σ indication of 5 MeV excess coming from ^{235}U fuel isotope fission !!



← (End of reactor cycle)

(Beginning of reactor cycle) →

Summary

- **More precise measurement of θ_{13} and $|\Delta m_{ee}^2|$ (2200 days)**

$$\sin^2 2\theta_{13} = 0.0896 \pm 0.0068 \text{ (7.6 \%)}$$

$$|\Delta m_{ee}^2| = 2.68 \pm 0.14 (\times 10^{-3} \text{ eV}^2) \text{ (5.2 \%)}$$

- **Observation of fuel composition dependent IBD yield at 6.6σ CL**

- **Measured IBD yield per fission (10^{-43} cm^2)**

^{235}U : 6.15 ± 0.19 (smaller than the H-M prediction at 2.8σ)

^{239}Pu : 4.18 ± 0.26 (smaller than the H-M prediction at 0.8σ)

- **First hint for 2.9σ correlation between 5 MeV excess and ^{235}U fission fraction**

Thanks for your attention!