



Form Factors and High-Mass Moments in $B \rightarrow K\pi ll$

Javier Virto

Universitat de Barcelona

[arXiv:1908.02267 \[hep-ph\]](https://arxiv.org/abs/1908.02267) in collaboration with S. Descotes-Genon, A. Khodjamirian



Workshop on the Implications of LHCb Measurements – CERN – October 16th, 2019

1. **Definition:** $\mathcal{F}_i(q^2) \sim \langle V(k) | \bar{q} \Gamma_i b | B(q+k) \rangle$

2. Necessary for:

- Semileptonic decays: $B \rightarrow \rho l \nu$, $B_s \rightarrow K^* l \nu$, ...
- Non-Leptonic decays: $B \rightarrow K^* \pi$, ...
- “Rare” FCNC decays: $B \rightarrow K^* \bar{\nu} \nu$, $B \rightarrow K^* \ell^+ \ell^-$

Local $B \rightarrow K^*$ Form Factors



- ▶ Two main approaches: (1) **Lattice QCD** (large q^2) (2) **LCSRs** (low q^2)
- ▶ Two approaches to **LCSRs**, in terms of (Left) K^* LCDAs (Right) B LCDAs
- ▶ q^2 dependence can be parametrized model-independently

Subject of this talk

However:

- ρ, K^*, \dots are not stable in QCD (e.g. $K^* \rightarrow K\pi$ strong decay)
- Form factor calculations done in the narrow-width limit

This talk:

$$B \rightarrow K^* X \quad \text{---} \longrightarrow \quad B \rightarrow K\pi X$$

Naively, corrections from finite width are

$$\mathcal{W} \sim 1 + \text{coeff.} \times \frac{\Gamma}{M} + \dots$$

Target precision: $\sim 10\%$ $\Gamma/M \sim 20\%(\rho), 6\%(K^*), 0.5\%(\phi)$

But there are also “non-resonant” effects (higher resonances, S, D-waves, ...)

$B \rightarrow K\pi$ Form factors

Definition of Lorentz-Invariant Form Factors:

$$\begin{aligned}i\langle K^-(k_1)\pi^+(k_2)|\bar{s}\gamma^\mu b|\bar{B}^0(q+k)\rangle &= F_\perp k_\perp^\mu \\-i\langle K^-(k_1)\pi^+(k_2)|\bar{s}\gamma^\mu\gamma_5 b|\bar{B}^0(q+k)\rangle &= F_t k_t^\mu + F_0 k_0^\mu + F_\parallel k_\parallel^\mu \\\langle K^-(k_1)\pi^+(k_2)|\bar{s}\sigma^{\mu\nu}q_\nu b|\bar{B}^0(q+k)\rangle &= F_\perp^T k_\perp^\mu \\\langle K^-(k_1)\pi^+(k_2)|\bar{s}\sigma^{\mu\nu}q_\nu\gamma_5 b|\bar{B}^0(q+k)\rangle &= F_0^T k_0^\mu + F_\parallel^T k_\parallel^\mu\end{aligned}$$

Functions $F_i^{(T)}(k^2, q^2, q \cdot \bar{k})$. Partial-wave expansion:


$$\begin{aligned}F_{0,t}(k^2, q^2, q \cdot \bar{k}) &= \sum_{\ell=0}^{\infty} \sqrt{2\ell+1} F_{0,t}^{(\ell)}(k^2, q^2) P_\ell^{(0)}(\cos\theta_K) \\F_{\perp,\parallel}(k^2, q^2, q \cdot \bar{k}) &= \sum_{\ell=1}^{\infty} \sqrt{2\ell+1} F_{\perp,\parallel}^{(\ell)}(k^2, q^2) \frac{P_\ell^{(1)}(\cos\theta_K)}{\sin\theta_K}\end{aligned}$$

Light-Cone Sum Rules with B -meson LCDAs

Khodjamirian, Mannel, Offen 2006

[Analyticity+Unitarity+Duality]

Consider a correlation function: $\mathcal{P}_{ab}(k, q) = i \int d^4x e^{ik \cdot x} \langle 0 | T \{ j_a(x), j_b(0) \} | \bar{B}^0(q+k) \rangle$




► Traditionally, $h(k) = K^* + \text{continuum} \Rightarrow 2 \text{Im} \mathcal{P}_{ab}(k, q) \sim f_K^* F^{BK^*} \delta(k^2 - m_{K^*}) + \dots$

Light-Cone Sum Rules with B -meson LCDAs

Khodjamirian, Mannel, Offen 2006

[Analyticity+Unitarity+Duality]

Consider a correlation function: $\mathcal{P}_{ab}(k, q) = i \int d^4x e^{ik \cdot x} \langle 0 | T \{ j_a(x), j_b(0) \} | \bar{B}^0(q+k) \rangle$



► Traditionally, $h(k) = K^* + \text{continuum} \Rightarrow 2 \text{Im} \mathcal{P}_{ab}(k, q) \sim f_K^* F^{BK*} \delta(k^2 - m_{K^*}^2) + \dots$

► Generalization for unstable mesons Cheng, Khodjamirian, Virto 2017: $h(k) = K\pi + \dots$

LCSRs with B -meson DAs, natural for this generalization.

Light-Cone Sum Rules for P -wave $B \rightarrow K\pi$ Form Factors

$$\int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \omega_i(s, q^2) f_+^*(s) F_i^{(T)(\ell=1)}(s, q^2) = \mathcal{P}_i^{(T),\text{OPE}}(q^2, \sigma_0, M^2)$$

- s_0 – Effective threshold
- $\omega_i(s, q^2)$ – (known) kinematic factors
- $\langle K^-(k_1)\pi^+(k_2)|\bar{s}\gamma_\mu d|0\rangle = f_+(k^2) \bar{k}_\mu + \frac{m_K^2 - m_\pi^2}{k^2} f_0(k^2) k_\mu$
- $\mathcal{P}_i^{(T),\text{OPE}}$ – OPE result for the correlation function

$$\int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \omega_i(s, q^2) f_+^*(s) F_i^{(T)(\ell=1)}(s, q^2) = \mathcal{P}_i^{(T), \text{OPE}}(q^2, \sigma_0, M^2)$$

- Generalize LCSRs in [Khodjamirian, Mannel, Offen 2006](#) beyond the K^* case, including LCSRs for $A_0, T_{2,3}$
- Recalculate $\mathcal{P}_i^{(T), \text{OPE}}$ including 3-particle contributions, and extended consistently to twist-4 accuracy. Full (numerical) agreement with [Gubernari, Kokulu, van Dyk 2018](#) (not input parameters)
- Revisit $s_0 \Rightarrow$ significantly lower value!! – f_{K^*} is derived quantity
- Study of Narrow-width limit, Finite-Width effects, and effects beyond the K^*
- Applications to $B \rightarrow K\pi\ell\ell$

$K\pi$ form factor $f_+(s)$ from $\tau \rightarrow K\pi\nu_\tau$

Differential decay rate of $\tau \rightarrow K\pi\nu_\tau$:

$$\frac{d\Gamma}{ds} = \frac{N_\tau}{s^3} \left(1 - \frac{s}{m_\tau^2}\right)^2 \left(1 + 2\frac{s}{m_\tau^2}\right) \lambda_{K\pi}^{3/2} |\tilde{f}_+(s)|^2 \left\{ 1 + \frac{3(\Delta m^2)^2}{(1 + 2s/m_\tau^2) \lambda_{K\pi}} |\tilde{f}_0(s)|^2 \right\}$$

with the normalization [Total BR will give $|f_+(0)|^2 = 0.99$, consistent with $f_+^{LQCD}(0) = 0.97$]

$$N_\tau = \frac{G_F^2 |V_{us}|^2 |f_+(0)|^2 m_\tau^3}{1536\pi^3} S_{EW}^{\text{had}}$$

Belle fits to models: [This gives $f_{K^*} \simeq 205$ MeV, compared to $f_{K^*} = 217(5)$ MeV (NWL)]



$$\tilde{f}_+(s) = \sum_R \frac{\xi_R m_R^2}{m_R^2 - s - i\sqrt{s}\Gamma_R(s)}, \quad f_0(s) = f_+(0) \cdot \sum_{R_0} \frac{\xi_{R_0} s}{m_{R_0}^2 - s - i\sqrt{s}\Gamma_{R_0}(s)},$$

Model 1 : $\xi_{K^*(892)} = 1, \xi_{K_0^*(800)} = 1.27, \xi_{K_0^*(1430)} = 0.954 e^{i0.62}$

Model 2 : $\xi_{K^*(892)} = 0.988 e^{-i0.07}, \xi_{K^*(1410)} = 0.074 e^{i1.37}, \xi_{K_0^*(800)} = 1.57$

$K\pi$ form factor $f_+(s)$ from $\tau \rightarrow K\pi\nu_\tau$

Data from Belle, arXiv:0706.2231 [hep-ex]



Effective threshold: 2-point SVZ sum rule

Knowing $|f_+(s)|$ we can extract s_0 from a QCD sum rule:

$$\begin{aligned}\Pi_{\mu\nu}(k) &= i \int d^4x e^{ikx} \langle 0 | T \{ \bar{d}(x) \gamma_\mu s(x), \bar{s}(0) \gamma_\nu d(0) \} | 0 \rangle \\ &= (k_\mu k_\nu - k^2 g_{\mu\nu}) \Pi(k^2) + k_\mu k_\nu \tilde{\Pi}(k^2)\end{aligned}$$

$$\Pi(M^2, s_0) \equiv \frac{1}{\pi} \int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \text{Im} \Pi(s) = \int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \frac{\lambda_{K\pi}^{3/2}(s)}{32\pi^2 s^3} |f_+(s)|^2$$

$$\begin{aligned}\Pi^{\text{OPE}}(M^2, s_0) &= \frac{1}{8\pi^2} \int_{m_s^2}^{s_0} ds e^{-s/M^2} \frac{(s - m_s^2)^2 (2s + m_s^2)}{s^3} \\ &\quad + \frac{\alpha_s(M)}{\pi} \frac{M^2}{4\pi^2} \left(1 - e^{-s_0/M^2} \right) + \frac{V_4}{M^2} + \frac{V_6}{2M^4}\end{aligned}$$

Effective threshold: 2-point SVZ sum rule

Borel parameter M^2	Effective threshold s_0	
1.00 GeV ²	1.28 ± 0.18 GeV ² (Model 1)	1.26 ± 0.18 GeV ² (Average)
	1.25 ± 0.18 GeV ² (Model 2)	
1.25 GeV ²	1.33 ± 0.12 GeV ² (Model 1)	1.31 ± 0.12 GeV ² (Average)
	1.31 ± 0.12 GeV ² (Model 2)	
1.50 GeV ²	1.36 ± 0.09 GeV ² (Model 1)	1.35 ± 0.09 GeV ² (Average)
	1.34 ± 0.09 GeV ² (Model 2)	

Table 3: Values for the effective threshold s_0 extracted from the SVZ sum rules.

Significantly low value compared to the usual $s_0^{K^*} \simeq 1.7 \text{ GeV}^2 \sim (\sqrt{s_0^P} + m_s)^2$

Models for $B \rightarrow K\pi$ form factors

Assume that the P -wave $K\pi$ state couples to its interpolating current $\bar{s}\Gamma d$ resonantly, through a set of Breit-Wigner-type vector resonances:

$$\langle K(k_1)\pi(k_2)|\bar{s}\gamma^\mu d|X\rangle = \sum_{R,\eta} BW_R(k^2)\langle K(k_1)\pi(k_2)|R(k,\eta)\rangle\langle R(k,\eta)|\bar{s}\gamma^\mu d|X\rangle$$

$$f_+(s) = -\sum_R \frac{m_R f_R g_{RK\pi} e^{i\phi_R(s)}}{m_R^2 - s - i\sqrt{s}\Gamma_R(s)}$$

$$F_i^{(T),(\ell=1)}(s, q^2) = \sum_R \frac{Y_{R,i}^{(T)}(s, q^2) g_{RK\pi} \mathcal{F}_{R,i}^{(T)}(q^2) e^{i\phi_R(s)}}{m_R^2 - s - i\sqrt{s}\Gamma_R(s)}$$

This model is totally equivalent to the model fitted by Belle for $f_+(s)$.

Light-Cone Sum Rule + BW model

$$\sum_R \mathcal{F}_{R,i}^{(T)}(q^2) d_{R,i}^{(T)} I_R(s_0, M^2) = \mathcal{P}_i^{(T), \text{OPE}}(q^2, \sigma_0, M^2)$$

with

$$I_R(s_0, M^2) = \frac{m_R}{16 \pi^2} \int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \frac{g_{RK\pi} \lambda_{K\pi}^{3/2}(s) |f_+(s)|}{s^{5/2} \sqrt{(m_B^2 - s)^2 + s \Gamma_R^2(s)}}$$

and

$$d_{R,\perp} = -d_{R,-} = (m_B + m_R)^{-1}, \quad d_{R,\parallel} = \frac{(m_B + m_R)}{2}, \quad d_{R,t} = -m_R,$$
$$d_{R,\perp}^T = -d_{R,-}^T = 1, \quad d_{R,\parallel}^T = \frac{(m_B^2 - m_R^2)}{2}.$$

Narrow-width limit

Consider the sum rule with a single resonance R :

$$\mathcal{F}_{R,i}^{(T)}(q^2) d_{R,i}^{(T)} I_R(s_0, M^2) = \mathcal{P}_i^{(T), \text{OPE}}(q^2, \sigma_0, M^2)$$

$$I_R(s_0, M^2) = 3 m_R f_R \mathcal{B}(R \rightarrow K^+ \pi^-) \int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \frac{m_R}{\sqrt{s}} \left[\frac{1}{\pi} \frac{\sqrt{s} \Gamma_R(s)}{(m_R^2 - s)^2 + s \Gamma_R^2(s)} \right]$$

$$\xrightarrow{\Gamma_R^{\text{tot}} \rightarrow 0} 3 m_R f_R \mathcal{B}(R \rightarrow K^+ \pi^-) e^{-m_R^2/M^2}$$

$$\Rightarrow 3 m_R f_R d_{R,i}^{(T)} \mathcal{F}_{R,i}^{(T)}(q^2) e^{-m_R^2/M^2} \mathcal{B}(R \rightarrow K^+ \pi^-) = \mathcal{P}_i^{(T), \text{OPE}}(q^2, \sigma_0, M^2)$$

This agrees with Khodjamirian, Mannel, Offen 2006

Finite-width effects

Consider the sum rule with a single K^* :

$$\mathcal{F}_{K^*,i}^{(T)}(q^2) d_{K^*,i}^{(T)} l_{K^*}(s_0, M^2) = \mathcal{P}_i^{(T),\text{OPE}}(q^2, \sigma_0, M^2)$$


Define the “Width ratio” \mathcal{W}_{K^*} :

$$\mathcal{W}_{K^*} \equiv \frac{\mathcal{F}_{K^*,i}^{(T)}(q^2)}{\mathcal{F}_{K^*,i}^{(T)}(q^2)_{\text{NWL}}} = \frac{l_{K^*}(s_0, M^2)|_{\Gamma_{K^*} \rightarrow 0}}{l_{K^*}(s_0, M^2)} = \frac{2m_{K^*} f_{K^*} e^{-m_{K^*}^2/M^2}}{l_{K^*}(s_0, M^2)}$$

- \mathcal{W}_{K^*} is independent of the form factor type
- \mathcal{W}_{K^*} is independent of q^2

⇒ BRs are corrected by $|\mathcal{W}_{K^*}|^2$, ratios are uncorrected! + true in q^2 bins.

Finite-width effects



$$\mathcal{W}_{K^*} \simeq 1 + 1.9 \frac{\Gamma_{K^*}}{m_{K^*}}$$
$$\mathcal{W}_{K^*} = 1.09 \pm 0.01$$

\Rightarrow BRs are corrected by a factor $|\mathcal{W}_{K^*}|^2 \simeq 1.2$

Beyond the $K^*(892)$

Consider the sum rule with $R = \{K^*(892), K^*(1410)\}$:


$$\sum_R \mathcal{F}_{R,i}^{(\tau)}(q^2) d_{R,i}^{(\tau)} I_R(S_0, M^2) = \mathcal{P}_i^{(\tau), \text{OPE}}(q^2, \sigma_0, M^2)$$

		$M^2 = 1.00 \text{ GeV}^2$	$M^2 = 1.25 \text{ GeV}^2$	$M^2 = 1.50 \text{ GeV}^2$
Model 1	$I_{K^*(892)}$	0.1506(23)	0.1781(16)	0.1992(13)
	$I_{K^*(1410)}$	0.0050(07)	0.0062(07)	0.0072(06)
Model 2	$I_{K^*(892)}$	0.1491(22)	0.1766(20)	0.1975(16)
	$I_{K^*(1410)}$	0.0048(07)	0.0061(06)	0.0070(06)

Table 8: Values for the quantities I_R for $R = \{K^*(892), K^*(1410)\}$ for the different values of the Borel parameter M^2 and for the two models for the $K\pi$ form factor. The $K^*(1410)$ contribution is very suppressed in the sum rules, with $I_{K^*(1410)}/I_{K^*(892)} \simeq 0.03$ in all cases.

Beyond the $K^*(892)$

Set $\mathcal{F}_{K^*(1410)} = \alpha \mathcal{F}_{K^*(892)}$ with α a floating parameter



$\alpha = 1$: $\mathcal{F}_{K^*,\perp}(0) = 0.28$; $\alpha = 10$: $\mathcal{F}_{K^*,\perp}(0) = 0.22$; $\alpha = 50$: $\mathcal{F}_{K^*,\perp}(0) = 0.11$.


High $K\pi$ -Mass Moments in $B \rightarrow K\pi\ell\ell$

Differential decay rate including S,P,D waves -- [$d\Omega = d\cos\theta_\ell d\cos\theta_K d\phi$]

$$\frac{d\Gamma}{dq^2 dk^2 d\Omega} = \frac{1}{4\pi} \sum_{i=1}^{41} f_i(\Omega) \tilde{f}_i(q^2, k^2)$$

The 41 moments $\tilde{f}_i(q^2, k^2)$ have been measured by LHCb ([arXiv:1609.04736](https://arxiv.org/abs/1609.04736)) in the bins

$$\sqrt{k^2} \in [1.33, 1.53]\text{GeV}, \quad q^2 \in [1.1, 6]\text{GeV}^2$$




High $K\pi$ -Mass Moments in $B \rightarrow K\pi\ell\ell$

Differential decay rate including S,P,D waves – – [$d\Omega = d\cos\theta_\ell d\cos\theta_K d\phi$]

$$\frac{d\Gamma}{dq^2 dk^2 d\Omega} = \frac{1}{4\pi} \sum_{i=1}^{41} f_i(\Omega) \tilde{\Gamma}_i(q^2, k^2)$$

The 41 moments $\tilde{\Gamma}_i(q^2, k^2)$ have been measured by LHCb ([arXiv:1609.04736](https://arxiv.org/abs/1609.04736)) in the bins

$$\sqrt{k^2} \in [1.33, 1.53] \text{ GeV}, \quad q^2 \in [1.1, 6] \text{ GeV}^2$$



High $K\pi$ -Mass Moments in $B \rightarrow K\pi\ell\ell$

Differential decay rate including S, P, D waves – – $[d\Omega = d\cos\theta_\ell d\cos\theta_K d\phi]$

$$\frac{d\Gamma}{dq^2 dk^2 d\Omega} = \frac{1}{4\pi} \sum_{i=1}^{41} f_i(\Omega) \tilde{r}_i(q^2, k^2)$$

The 41 moments $\tilde{r}_i(q^2, k^2)$ depend on S, P, D -wave amplitudes:

i	$f_i(\Omega)$	$\Gamma_i^{L, \text{tr}}(q^2)/\mathbf{k}q^2$	$\eta_i^{L \rightarrow R}$
1	$P_0^0 Y_0^0$	$ H_0^L ^2 + H_{\parallel}^L ^2 + H_{\perp}^L ^2 + S^L ^2 + D_0^L ^2 + D_{\parallel}^L ^2 + D_{\perp}^L ^2$	+1
2	$P_1^0 Y_0^0$	$2 \left[\frac{2}{\sqrt{5}} \text{Re}(H_0^L D_0^{L*}) + \text{Re}(S^L H_0^{L*}) + \sqrt{\frac{3}{5}} \text{Re}(H_{\parallel}^L D_{\parallel}^{L*} + H_{\perp}^L D_{\perp}^{L*}) \right]$	+1
3	$P_2^0 Y_0^0$	$\frac{\sqrt{5}}{7} (D_{\parallel}^L ^2 + D_{\perp}^L ^2) - \frac{3}{\sqrt{5}} (H_{\parallel}^L ^2 + H_{\perp}^L ^2) + \frac{2}{\sqrt{5}} H_0^L ^2 + \frac{10}{7\sqrt{5}} D_0^L ^2 + 2 \text{Re}(S^L D_0^{L*})$	+1
4	$P_3^0 Y_0^0$	$\frac{6}{\sqrt{35}} \left[-\text{Re}(H_{\parallel}^L D_{\parallel}^{L*} + H_{\perp}^L D_{\perp}^{L*}) + \sqrt{3} \text{Re}(H_0^L D_0^{L*}) \right]$	+1
5	$P_4^0 Y_0^0$	$\frac{2}{7} \left[-2(D_{\parallel}^L ^2 + D_{\perp}^L ^2) + 3 D_0^L ^2 \right]$	+1
6	$P_0^0 Y_2^0$	$\frac{1}{2\sqrt{5}} \left[(D_{\parallel}^L ^2 + D_{\perp}^L ^2) + (H_{\parallel}^L ^2 + H_{\perp}^L ^2) - 2 S^L ^2 - 2 D_0^L ^2 - 2 H_0^L ^2 \right]$	+1
7	$P_1^0 Y_2^0$	$\left[\frac{\sqrt{3}}{5} \text{Re}(H_{\parallel}^L D_{\parallel}^{L*} + H_{\perp}^L D_{\perp}^{L*}) - \frac{2}{\sqrt{5}} \text{Re}(S^L H_0^{L*}) - \frac{4}{5} \text{Re}(H_0^L D_0^{L*}) \right]$	+1
8	$P_2^0 Y_2^0$	$\left[\frac{1}{14} (D_{\parallel}^L ^2 + D_{\perp}^L ^2) - \frac{2}{7} D_0^L ^2 - \frac{1}{10} (H_{\parallel}^L ^2 + H_{\perp}^L ^2) - \frac{2}{5} H_0^L ^2 - \frac{2}{\sqrt{5}} \text{Re}(S^L D_0^{L*}) \right]$	+1
9	$P_3^0 Y_2^0$	$-\frac{3}{5\sqrt{7}} \left[\text{Re}(H_{\parallel}^L D_{\parallel}^{L*} + H_{\perp}^L D_{\perp}^{L*}) + 2\sqrt{3} \text{Re}(H_0^L D_0^{L*}) \right]$	+1
10	$P_4^0 Y_2^0$	$-\frac{2}{7\sqrt{5}} \left[D_{\parallel}^L ^2 + D_{\perp}^L ^2 + 3 D_0^L ^2 \right]$	+1
11	$P_1^1 \sqrt{2} \text{Re}(Y_2^1)$	$-\frac{3}{\sqrt{10}} \left[\sqrt{\frac{2}{3}} \text{Re}(H_{\parallel}^L S^{L*}) - \sqrt{\frac{2}{15}} \text{Re}(H_{\parallel}^L D_0^{L*}) + \sqrt{\frac{2}{5}} \text{Re}(D_{\parallel}^L H_0^{L*}) \right]$	+1
12	$P_2^1 \sqrt{2} \text{Re}(Y_2^1)$	$-\frac{3}{2} \left[\text{Re}(H_{\parallel}^L H_{\parallel}^{L*}) + \sqrt{\frac{1}{5}} \text{Re}(D_{\parallel}^L S^{L*}) + \frac{5}{2} \text{Re}(D_{\parallel}^L D_{\parallel}^{L*}) \right]$	+1

High $K\pi$ -Mass Moments in $B \rightarrow K\pi\ell\ell$

Combinations of moments depending **only on P -wave**:



$$\begin{aligned} |\widehat{A}_{\parallel}^L|^2 + |\widehat{A}_{\parallel}^R|^2 &= \frac{1}{36} (5\tilde{r}_1 - 7\sqrt{5}\tilde{r}_3 + 5\sqrt{5}\tilde{r}_6 - 35\tilde{r}_8 - 5\sqrt{15}\tilde{r}_{19} + 35\sqrt{3}\tilde{r}_{21}) \\ |\widehat{A}_{\perp}^L|^2 + |\widehat{A}_{\perp}^R|^2 &= \frac{1}{36} (5\tilde{r}_1 - 7\sqrt{5}\tilde{r}_3 + 5\sqrt{5}\tilde{r}_6 - 35\tilde{r}_8 + 5\sqrt{15}\tilde{r}_{19} - 35\sqrt{3}\tilde{r}_{21}) \\ \text{Im}(\widehat{A}_{\perp}^L \widehat{A}_{\parallel}^{L*} + \widehat{A}_{\perp}^R \widehat{A}_{\parallel}^{R*}) &= \frac{5}{36} (\sqrt{15}\tilde{r}_{24} - 7\sqrt{3}\tilde{r}_{26}) \\ \text{Re}(\widehat{A}_{\perp}^L \widehat{A}_{\parallel}^{L*} - \widehat{A}_{\perp}^R \widehat{A}_{\parallel}^{R*}) &= \frac{1}{36} (-5\sqrt{3}\tilde{r}_{29} + 7\sqrt{15}\tilde{r}_{31}) \end{aligned}$$

Binned LHCb results ([arXiv:1609.04736](https://arxiv.org/abs/1609.04736)) imply:

$$\begin{aligned} \tau_B \langle |\widehat{A}_{\parallel}^L|^2 + |\widehat{A}_{\parallel}^R|^2 \rangle &\equiv \langle M_{\parallel} \rangle = (1.07 \pm 1.13) \times 10^{-8} \\ \tau_B \langle |\widehat{A}_{\perp}^L|^2 + |\widehat{A}_{\perp}^R|^2 \rangle &\equiv \langle M_{\perp} \rangle = (0.94 \pm 1.06) \times 10^{-8} \\ \tau_B \langle \text{Im}(\widehat{A}_{\perp}^L \widehat{A}_{\parallel}^{L*} + \widehat{A}_{\perp}^R \widehat{A}_{\parallel}^{R*}) \rangle &\equiv \langle M_{\text{im}} \rangle = (-0.75 \pm 0.79) \times 10^{-8} \\ \tau_B \langle \text{Re}(\widehat{A}_{\perp}^L \widehat{A}_{\parallel}^{L*} - \widehat{A}_{\perp}^R \widehat{A}_{\parallel}^{R*}) \rangle &\equiv \langle M_{\text{re}} \rangle = (0.27 \pm 0.50) \times 10^{-8} \end{aligned}$$

High $K\pi$ -Mass Moments in $B \rightarrow K\pi\ell\ell$

Example: $\langle M_{\parallel} \rangle$:




Bounds: From $\langle M_{\parallel} \rangle$: $\alpha \lesssim 11$; From $\langle M_{\perp} \rangle$: $\alpha \lesssim 17$; From $\langle M_{\text{re}} \rangle$: $\alpha \lesssim 18$.

High $K\pi$ -Mass Moments in $B \rightarrow K\pi\ell\ell$

Upper bounds on P -wave from differential BR:

$$\frac{d\Gamma}{dq^2 dk^2} = \tilde{\Gamma}_1 = |\hat{A}_{\parallel}^L|^2 + |\hat{A}_{\parallel}^R|^2 + |\hat{A}_{\perp}^L|^2 + |\hat{A}_{\perp}^R|^2 + |\hat{A}_0^L|^2 + |\hat{A}_0^R|^2 + \dots$$



$$\begin{aligned} 10^8 \cdot \langle \mathcal{B} \rangle_{[0.10, 0.98]} &= 1.41 \pm 0.27 \rightarrow \alpha \lesssim 5 \\ 10^8 \cdot \langle \mathcal{B} \rangle_{[1.10, 2.50]} &= 1.60 \pm 0.29 \rightarrow \alpha \lesssim 6 \\ 10^8 \cdot \langle \mathcal{B} \rangle_{[2.50, 4.00]} &= 1.37 \pm 0.26 \rightarrow \alpha \lesssim 5 \\ 10^8 \cdot \langle \mathcal{B} \rangle_{[4.00, 6.00]} &= 1.12 \pm 0.26 \rightarrow \alpha \lesssim 4 \\ 10^8 \cdot \langle \mathcal{B} \rangle_{[6.00, 8.00]} &= 0.98 \pm 0.23 \rightarrow \alpha \lesssim 3 \end{aligned}$$

Bounds are easily improved with some info on S-wave form factors.

Summary

- Absolutely no excuse to do the transition $K^* \rightarrow K\pi$ in your life
- Recalculation of all $B \rightarrow K^*$ form factors from LCSRs with B -DAs with twist-4 accuracy
- Finite-Width effects are **20%** at the level of BRs, universal and q^2 -independent \Rightarrow Global factor 1.2 in BRs; but ratios (e.g. P'_5 unaffected.)
- Higher resonance effects can have dramatic effect on $B \rightarrow K^*$.
High mass BRs and Moments very efficient to bound this possibility
- Measurements of angular moments in bins across the q^2 and k^2 spectra \Rightarrow **very useful**

Thank You

Extra

Form Factor	This work	Ref. [12]	Ref. [24]	Ref. [15]	Ref. [17]
$\mathcal{F}_{K^*,\perp}(0) = V^{BK^*}(0)$	0.26(15)	0.39(11)	0.36(18)	0.32(11)	0.34(4)
$\mathcal{F}_{K^*,\parallel}(0) = A_1^{BK^*}(0)$	0.20(12)	0.30(8)	0.25(13)	0.26(8)	0.27(3)
$\mathcal{F}_{K^*,-}(0) = A_2^{BK^*}(0)$	0.14(13)	0.26(8)	0.23(15)	0.24(9)	0.23(5)
$\mathcal{F}_{K^*,t}(0) = A_0^{BK^*}(0)$	0.30(7)	–	0.29(8)	0.31(7)	0.36(5)
$\mathcal{F}_{K^*,\perp}^T(0) = T_1^{BK^*}(0)$	0.22(13)	0.33(10)	0.31(14)	0.29(10)	0.28(3)
$\mathcal{F}_{K^*,\parallel}^T(0) = T_2^{BK^*}(0)$	0.22(13)	0.33(10)	0.31(14)	0.29(10)	0.28(3)
$\mathcal{F}_{K^*,-}^T(0) = T_3^{BK^*}(0)$	0.13(12)	–	0.22(14)	0.20(8)	0.18(3)

Table 6: *Results for the form factors at $q^2 = 0$ in the narrow-width limit, compared to corresponding results in the literature. The approach in Ref. [17] is a completely different LCSR approach, in terms of K^* DAs.*

$\mathcal{F}^{BK^*}(q^2 = 0)$	V^{BK^*}	$A_1^{BK^*}$	$A_2^{BK^*}$	$A_0^{BK^*}$	$T_{1,2}^{BK^*}$	$T_3^{BK^*}$
Ref. [12]	0.39	0.30	0.26	–	0.33	–
Inputs [12], no g_+	0.38	0.29	0.26	0.31	0.33	0.25
Inputs [12], with g_+	0.27	0.21	0.14	0.31	0.24	0.14
Our inputs, but $s_0 = 1.7 \text{ GeV}^2$	0.33	0.26	0.17	0.38	0.29	0.17
Our inputs, our s_0 , no g_+	0.36	0.28	0.25	0.30	0.31	0.23
Our inputs, our s_0 , with g_+	0.26	0.20	0.14	0.30	0.22	0.13

Table 7: *Deconstruction of the different effects explaining the difference between our results for the form factors at $q^2 = 0$ and those in Ref. [12]. The difference stems mainly from the inclusion of the twist-four two-particle contributions. See the text for more details.*

Light-Cone Sum Rules with B -meson LCDAs

Khodjamirian, Mannel, Offen 2006

Consider a correlation function of the type:

$$\mathcal{P}_{ab}(k, q) = i \int d^4x e^{ik \cdot x} \langle 0 | T \{ j_a(x), j_b(0) \} | \bar{B}^0(q+k) \rangle$$

which obeys a dispersion relation:

$$\mathcal{P}_{ab}^{\text{OPE}}(k^2, q^2) = \frac{1}{\pi} \int_{s_{\text{th}}}^{\infty} ds \frac{\text{Im} \mathcal{P}_{ab}(s, q^2)}{s - k^2}$$

Duality + Borel transformation:

$$\frac{1}{\pi} \int_{s_{\text{th}}}^{s_0} ds e^{-s/M^2} \text{Im} \mathcal{P}_{ab}(s, q^2) = \mathcal{P}_{ab}^{\text{OPE}}(q^2, \sigma_0, M^2),$$

Light-Cone Sum Rules with B -meson LCDAs

Khodjamirian, Mannel, Offen 2006

From Unitarity:

$$2 \operatorname{Im} \mathcal{P}_{ab}(k, q) = \sum_h \int d\tau_h \langle 0 | j_a | h(k) \rangle \underbrace{\langle h(k) | j_b | \bar{B}^0(q+k) \rangle}_{\text{form factor}}$$

▷ Traditionally,

$$h(k) = K^* + \text{continuum} \quad \Rightarrow \quad 2 \operatorname{Im} \mathcal{P}_{ab}(k, q) \sim f_K^* F^{BK^*} \delta(k^2 - m_{K^*}) + \dots$$

▷ Generalization for unstable mesons [Cheng, Khodjamirian, Virto 2017](#)

$$h(k) = K\pi + \dots$$

LCSRs with B -meson DAs, natural for this generalization.