

# Formation and Evolution of quarkonia states in rapidly varying strong magnetic field.

Partha Bagchi

Variable Energy Cyclotron Centre, Kolkata, India

December 14, 2018

XXIII DAE-BRNS HIGH ENERGY PHYSICS SYMPOSIUM 2018

December 10-14, 2018

IIT Madras

# Outline of Talk

- Review of Conventional Mechanism of quarkonia suppression
- Non Adiabaticity: An Important Aspect of Evolution of Quarkonia
- Quarkonia in Magnetic Field
- Results
- Summary



# Conventional Mechanism for Quarkonia Suppression

- Matsui and Satz<sup>1</sup> proposed  $J/\psi$  suppression as a signal for QGP due to Debye screening of the potential between  $q\bar{q}$ .
- If at a temperature  $T_D$ , the Debye screening length of the medium becomes less than the radius of quarkonia, then  $q\bar{q}$  may not form bound states.
- In the above picture, suppression of quarkonia occurs when the temperature of QGP achieves a value higher than  $T_D$ .

---

<sup>1</sup>T. Matsui and H. Satz, Phys.Lett. B178,416 (1986)

# Adiabatic Approximation

- If the QGP temperature remains below  $T_D$ , **no quarkonia suppression is expected** due to color screening(?) in the conventional mechanism.

## Description of quarkonia through effective potential

- $q\bar{q}$  potential changes **slowly** from initial temperature ( $V(T = T_i)$ ) to the final temperature ( $V(T_f)$ ).
- Initial quarkonium state evolves to the state corresponding to  $V(T_f)$  which is also a bound state for  $T_f < T_D$  with same quantum number as initial state, hence no quarkonium suppression for  $T < T_D$ .  $\implies$  **Adiabatic**

# Evolution of Fireball Created in Heavy Ion Collisions




Figure: Nuclear collision evolution epoch.

## Some Important Observations

- The fireball created in Heavy Ion Collision is **rapidly** evolving with time.
- If quarkonia is described in potential model then  $q\bar{q}$  potential is no-doubt time dependent.

### Note:

Matsui and Satz picture considers the static QGP only.

- **One need to solve Schrödinger equation for **time-dependent** Hamiltonian.**

# Adiabaticity Violation: An Important Aspect of Quarkonia Evolution

## Several possible Example of Adiabaticity Violation

- During Thermalisation <sup>1</sup>
- Cooling Phase <sup>2</sup>
- In presence of initial fluctuation <sup>3</sup>
- During Freeze-out
- **In presence of transient magnetic field**
  - Formation <sup>4</sup>
  - Spin Mixing <sup>5</sup>
  - **Spacial Excitation** <sup>6</sup>

---

<sup>1</sup> Bagchi and Srivastava, Mod.Phys.Lett. A30 (2015) no.32, 1550162

<sup>2</sup> Dutta and Borghini, Mod.Phys.Lett. A30 (2015) no.37, 1550205


<sup>3</sup> Bagchi et al., Springer Proc.Phys. 203 (2018) 493-495

<sup>4</sup> Guo et al. Physics Letters B 751 (2015) 215219

<sup>5</sup> Dutta et al., Eur.Phys.J. C78 (2018) no.6, 525

<sup>6</sup> Bagchi et al., arXiv:1805.04082

## During Thermalisation



**Figure:** Survival Probability  $p$  of  $J/\psi$  and  $\Upsilon$  vs. temperature of medium. Plots are given upto the temperature  $T_D$  for  $J/\psi$  and  $\Upsilon$  .



# Initial Fluctuation in Heavy Ion Collisions





Figure: Initial energy density fluctuation (Phys.Rev. C92 (2015) no.5, 054902).

# Formation of Quarkonia in Presence of Magnetic Field



**Figure:** Formation of quarkonia in presence of magnetic field (Physics Letters B 751 (2015) 215219).

## Quark Anti-quark Potential in QGP Medium

- In medium Debye screened potential between quark ( $q$ ) and anti-quark ( $\bar{q}$ )<sup>1</sup>

$$V(r) = -\frac{\alpha}{r} \exp(-m_D r) + \frac{\sigma}{m_D} (1 - \exp(-m_D r)) \quad (1)$$

- $m_D$  is the Debye mass<sup>2</sup>  $\Rightarrow$  Static limit ( $p_0 \rightarrow 0, |\vec{p}| = 0$ ) of the longitudinal part of the gluon self energy  $\pi_{\mu\nu}$
- $m_D$  for three flavor case<sup>3</sup>

$$m_D = gT \sqrt{1 + N_f/6} \quad (2)$$

<sup>1</sup>H. Satz, J. Phys. Conf. Ser. **455**, 012045 (2013).

<sup>2</sup>E. Braaten and A. Nieto, Phys. Rev. Lett. **73**, 2402 (1994)

<sup>3</sup>F. Karsch, M.T. Mehr, and H. Satz, Z. Phys. C **37**, 617 (1988).

## Quark Anti-quark Potential in Presence of Magnetic Field

- The effect of magnetic field in the fermion self energy is incorporated through the fermion propagator.
- In the strong field limit:

$$S_0(k) = i \frac{m + \gamma \cdot k_{\parallel}}{k_{\parallel}^2 - m^2} (1 - i\gamma_1\gamma_2) e^{\frac{-k_{\perp}^2}{|q_f B|}} \quad (3)$$

- B is along Z-axis.
- The self energy is calculated by using thermal propagator in imaginary time formalism.
- Debye mass is then obtained as<sup>1</sup>:

$$m_D^2 = g'^2 T^2 + \frac{g^2}{4\pi^2 T} \sum_f |q_f B| \int_0^{\infty} dp_z \frac{e^{\beta \sqrt{p_z^2 + m_f^2}}}{\left(1 + e^{\beta \sqrt{p_z^2 + m_f^2}}\right)^2} \quad (4)$$

<sup>1</sup>Hasan et al., arXiv:1802.06874

# Quark Anti-quark Potential in Presence of Magnetic Field

Continued....

- First term is the contribution from the gluon loops and this is solely dependent on temperature
- $g'^2 = 4\pi\alpha'_s(T)$  where  $\alpha'_s(T)$  is the usual temperature dependent running coupling where the renormalization scale is taken as  $2\pi T$

- 

$$\alpha'_s(T) = \frac{2\pi}{\left(11 - \frac{2}{3}N_f\right) \ln\left(\frac{\Lambda}{\Lambda_{QCD}}\right)} \quad (5)$$

Where  $\Lambda = 2\pi T$  and  $\Lambda_{QCD} \sim 200$  MeV

# Quark Anti-quark Potential in Presence of Magnetic Field

Continued...

- Second term is the contribution from the fermion loop and this term strongly depends on magnetic field.
- $g^2 = 4\pi\alpha_s^{\parallel}(k_z, q_f B)$ , where  $\alpha_s^{\parallel}(k_z, q_f B)$  is the magnetic field dependent coupling and doesn't depend on temperature. <sup>1 2</sup>
- 

$$\alpha_s^{\parallel}(k_z, q_f B) = \frac{1}{\alpha_s^0(\mu_0)^{-1} + \frac{11N_c}{12\pi} \ln\left(\frac{k_z^2 + M_B^2}{\mu_0^2}\right) + \frac{1}{3\pi} \sum_f \frac{q_f B}{\sigma}} \quad (6)$$

$$\text{where, } \alpha_s^0(\mu_0) = \frac{12\pi}{11N_c \ln\left(\frac{\mu_0^2 + M_B^2}{\Lambda_V^2}\right)}$$


and,  $M_B = 1 \text{ GeV}$ ,  $\sigma = 0.18 \text{ GeV}^2$ ,  $\mu_0 = 1.1 \text{ GeV}$ ,  
 $\Lambda_V = 0.385 \text{ GeV}$ .

<sup>1</sup> Andreichikov et al., Phys. Rev. Lett. 110, 162002 (2013)

<sup>2</sup> Ferrer et al., Phys. Rev. D91, 054006 (2015)

## Evolution of Magnetic Field in Non Central Collisions

- In Heavy Ion Collisions there are certain possibilities of production of huge magnetic field for non-central collision.
- The magnetic field will last for only few fm/c time<sup>1</sup>.



**Figure:** Magnetic field for  $\sigma_e = 5.8 \text{ MeV}$ ,  $z = 0.2 \text{ fm}$ ,  $t_0 = 0.2 \text{ fm}$ . Solid, dashed, and dotted lines stand for  $B$ ,  $B_{init}$  and  $B_{val}$ ,  $\gamma = 2000$ .

- $\sigma_e$  (Electrical conductivity) = 0, for  $t < t_0$  (QGP formation time)

<sup>1</sup>Kirill Tuchin, Phys. Rev. C 93, 014905 (2016)

# Time-Dependent Potential for Studying Quarkonia Wave-Function Evolution

- Magnetic Field is Transient in Nature
- Decays to order of magnitude within few  $fm/c$  time.
- The evolution of the wave function, thus, cannot be taken to be adiabatic and it should be treated in terms of a time dependent perturbation theory (**is one of the tool**).
- Survival probability of quarkonia should be calculated under this perturbation.



# Results

- We have calculated dissociation energy of  $J/\psi$  ( $T = 1.7T_c$ ) and  $\Upsilon(1S)$  ( $T = 3.0T_c$ ) for different values of magnetic field <sup>1</sup>
- In the strong field limit, the effect of the temperature is suppressed.  
⇒ **strongly bound quarkonia**

---

<sup>1</sup>F. Karsch, M.T. Mehr, and H. Satz, Z. Phys. C **37**, 617 (1988).

# Results

## Dissociation Energy




Figure: Magnetic field vs dissociation energy for charmonia.

## Results

- Considering magnetic field starting from  $15m_\pi^2$ , decays with time like

$$B(t) = \frac{B_0}{(1.0+0.706896(t+6.23841t^2-1.39341t^3+0.108236t^4))}$$

- The temperature starting from  $1.7T_c$  for  $J/\psi$  ( $3.0T_c$  for  $\Upsilon(1S)$ ) decays like

$$T(t) = T_0 \left( \frac{\tau_0}{\tau_0+t} \right)^{\frac{1}{3}}$$

- Then we have calculated the transition probability of quarkonia from its ground states to continuum states using 1st order perturbation theory.

# Results

## Dissociation Probability




Figure: Magnetic field vs dissociation probability.

## Summary

- The presence of strong magnetic field makes quarkonia strongly bound .  
⇒ more or less true for all available potential present in the community still now
- Even non-adiabatic evolution can not dissociate  $\Upsilon(1S)$  at an initial temperature  $T = 3T_c$ .  
⇒ **contradictory with experimental results**
- **Possibilities:**
  - 1: There will be no(/very weak) magnetic field present when medium formed
  - 2: Behavior of quarkonia may be drastically opposite in presence of weak(/intermediate) magnetic field in comparison with the presence of strong field.

# Thank You !