



Contribution ID: 41

Type: Talk

Enhancement of Gravitational Waves Induced by Scalar Perturbations due to a Sudden Transition from an Early Matter Era to the Radiation Era

Thursday 5 September 2019 17:20 (20 minutes)

We study gravitational waves induced from the primordial scalar perturbations at second order around the reheating of the Universe. We consider reheating scenarios in which a transition from an early matter dominated era to the radiation dominated era completes within a timescale much shorter than the Hubble time at that time. We find that an enhanced production of induced gravitational waves occurs just after the reheating transition because of fast oscillations of scalar modes well inside the Hubble horizon. This enhancement mechanism just after an early matter-dominated era is much more efficient than a previously known enhancement mechanism during an early matter era, and we show that the induced gravitational waves could be detectable by future observations if the reheating temperature T_R is in the range $T_R \lesssim 7 \times 10^{-2} \text{ GeV}$ or $20 \text{ GeV} \lesssim T_R \lesssim 2 \times 10^7 \text{ GeV}$. This is the case even if the scalar perturbations on small scales are not enhanced relative to those on large scales, probed by the observations of the cosmic microwave background. This talk will be based on our paper, arXiv:1904.12879.

Author: Dr TERADA, Takahiro (KEK)

Co-authors: Mr INOMATA, Keisuke (Kavli IPMU, the University of Tokyo); Prof. KOHRI, Kazunori (KEK); Dr NAKAMA, Tomohiro (The Hong Kong University of Science and Technology)

Presenter: Dr TERADA, Takahiro (KEK)

Session Classification: Parallel Sessions: Gravitational Waves and Black Holes (C.A.R.L., H08)

Track Classification: Gravitational Waves and Black Holes