



**Measurement of top-quark  $\vec{p}$  olarization in t-channel  
single top-quark production using  $pp$  collisions at  $\sqrt{s}$   
=13TeV with the ATLAS detector**

Runyu Bi

---



DPF 2019 – August 1st, 2019




# Introduction

## \*Top polarization

- **Spin polarization**: represented by a polarization vector in 3D phase space,  $\vec{P}$ . The polarization depends upon the specific top quark(antiquark) sample.
- **Single Top**: t-channel in electroweak interaction produces highly polarized top quarks due to V-A nature.
- **Detectable**: Because of top's large mass, it decays before hadronization to an on-shell W boson. The decay products preserve the spin information of the top.



- **Interesting phenomenology**: pp collision at LHC results a large degree of polarization.




[arXiv:hep-ph/0011349](https://arxiv.org/abs/hep-ph/0011349)

# Introduction

## \*Motivations

- **Large lumi @LHC**: from  $9.7 \text{ fb}^{-1}$  to  **$140.5 \text{ fb}^{-1}$** .
- **Never measured**: the proposed fiducial measurement at reconstruction level of the single top polarization is a first. Proposed experimental measurement strategy: [Eur. Phys. J. C77 \(2017\) 200](#)
- **High polarization expected**: Mahlon, Parke ([Phys.Lett.B 476\(2000\)](#)) predicated a high polarized top ensemble in LHC, with a set of cuts comparable to our selection criteria.
- **Sensitive to new physics**:
  - Four-fermion operator, top couplings
  - CP violation if existing a nonzero  $P_y$  component.



# Introduction

## \* Motivations — Previous Analyses

- In 2016, CMS published the first measurement at 8TeV of the top polarization in t-channel, represented by a differential distribution of the top cross section as a function of the leptonic angle,  $\cos \theta_l$ . A smaller than prediction measurement was shown.




$$A_\mu = 0.26 \pm 0.03 \text{ (stat)} \pm 0.10 \text{ (syst)}$$

$$\Rightarrow P_Z = 0.52 \pm 0.20$$

**Prediction:**  $A_\mu = 0.44$

$$\Rightarrow P_Z = 0.88$$

- In 2017, ATLAS published two polarization measurements at 8TeV. Although different approaches, both analyses were consistent with a high value of polarization with the SM predictions.




# Introduction

## \* Motivations — Previous Analyses

- In 2016, CMS published the first measurement of the top polarization in t-channel, represented by a differential distribution of the top cross section as a function of the leptonic angle,  $\cos \theta_l$ . A smaller than prediction measurement was shown.



$$A_\mu = 0.26 \pm 0.03 \text{ (stat)} \pm 0.10 \text{ (syst)} \quad \Rightarrow \quad P_Z = 0.52 \pm 0.20$$



★ Only the z-components were measured.

★ All three measurements were unfolded onto the parton level, measuring the top quark polarization both detected and undetected.

★ The top quark and anti-quark data were not separated.



J. High Energ. Phys. (2017) 2017: 17.


# Introduction

## \*Analysis Goal

- The polarization of the top quark can be determined through the angular distribution of its decay products.


$$\longrightarrow \frac{1}{\Gamma} \frac{d\Gamma}{d(\cos \theta_l)} = \frac{1}{2}(1 + \alpha_l P \cos \theta_l)$$

- Performing a cut-based fiducial measurement of the single top polarization in t-channel with full ATLAS Run II dataset at 13TeV (expected 140.2 fb<sup>-1</sup>).
- We have devised a template fit method to simultaneously measure the polarization vector in terms of three orthogonal axes: (P<sub>x</sub>, P<sub>y</sub>, P<sub>z</sub>).
- The polarization of the top quark and anti-quark will be measured **separately**.
- The results will then be compared to the theoretical predictions, which are (SM LO).



## \*Definitions

- In this analysis, we follow the listed choices to define our angular observables. They are:
  - Spectator quark (the untagged light jet) momentum as the top spin axis.
  - Lepton as the primary analyzer ( $\alpha_l = 1$ ).
  - Top quark and W boson are fully reconstructed.
  - Lepton momentum in the top rest frame is determined.



# Introduction

## \*Template Fit Strategy

- **Construct templates** in the variable  $\cos \theta_l$  for pure polarized ensembles:  $P_Z = \pm 1, P_X = \pm 1, P_Y = \pm 1$ .
  - Six templates needed to fully cover the differential cross section coming from all spin configurations + interference terms.
- **Obtain dataset for arbitrary polarizations** as a linear combination of these.
- Fit the coefficients to obtain the polarization in real/simulated data.
- The full expression of the differential decay distribution of  $t \rightarrow Wb \rightarrow l\nu b$  is:


$$\frac{1}{\Gamma} \frac{d\Gamma}{d\Omega d\Omega^*} = \frac{3}{64\pi^2} \frac{1}{N} \left\{ \left[ |a_{1\frac{1}{2}}|^2 (1 + \lambda \cos \theta^*)^2 + 2|a_{0-\frac{1}{2}}|^2 \sin^2 \theta^* \right] (1 + \vec{P} \cdot \vec{u}_L) \right. \\ \left. + \left[ 2|a_{0\frac{1}{2}}|^2 \sin^2 \theta^* + |a_{-1-\frac{1}{2}}|^2 (1 - \lambda \cos \theta^*)^2 \right] (1 - \vec{P} \cdot \vec{u}_L) \right. \\ \left. + \lambda 2\sqrt{2} \left[ \text{Re}(a_{0\frac{1}{2}} a_{1\frac{1}{2}}^* e^{-i\phi^*}) (1 + \lambda \cos \theta^*) \right. \right. \\ \left. \left. + \text{Re}(a_{-1-\frac{1}{2}} a_{0-\frac{1}{2}}^* e^{-i\phi^*}) (1 - \lambda \cos \theta^*) \right] \sin \theta^* \vec{P} \cdot \vec{u}_T \right. \\ \left. + \lambda 2\sqrt{2} \left[ \text{Im}(a_{0\frac{1}{2}} a_{1\frac{1}{2}}^* e^{-i\phi^*}) (1 + \lambda \cos \theta^*) \right. \right. \\ \left. \left. + \text{Im}(a_{-1-\frac{1}{2}} a_{0-\frac{1}{2}}^* e^{-i\phi^*}) (1 - \lambda \cos \theta^*) \right] \sin \theta^* \vec{P} \cdot \vec{u}_N \right\} .$$

Eur. Phys. J. C (2017) 77: 200.

where the  $a_{\lambda_W, \lambda_b}$  the coefficients are the transition amplitudes for  $t \rightarrow Wb$ , and  $\lambda$ 's are helicities. The unit vectors are:

- $\vec{u}_L = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$  in the direction of the W boson momentum in the top quark rest frame,
- $\vec{u}_T = (\cos \theta \cos \phi, \cos \theta \sin \phi, -\sin \theta)$ ,  $\vec{u}_N = (\sin \phi, -\cos \phi, 0)$  are two orthonormal vectors.
- $\lambda = 1$  for top quarks,  $\lambda = -1$  for top antiquarks.


PolManip package!



An illustration of angles that were used to define the top decay process.  $\theta$  and  $\phi$  are the polar and azimuthal angles of the W boson momentum in top rest frame, and  $\theta^*$  and  $\phi^*$  are the polar and azimuthal angles of the charged lepton momentum in W rest frame.

# Introduction



## \*Template Fit Strategy



# Introduction

## \*PolManip

- **A decay model for top quarks that modifies the polarization:** It takes the output of a single top event generator (i.e. Protos: LHE file), and re-decays the top quark into a lepton, neutrino and b quark according to a user-specified polarization state.
- **Convenient output**
- **Validated in ATLAS.**



**Validation plots:** comparisons are made between the original Photos sample (Protos), and the PolManip SM sample (SM gen).

## \*Event Selection Criteria — 4 regions

### • Selection Region (SR):


- PR
- $m_{l,b} < 153 \text{ GeV}$
- $34 \text{ GeV} < m_{top} < 206 \text{ GeV}$
- Trapezoidal Cut on  $\eta_{top}$  vs.  $\eta_j$
- $m_{j,top} > 280 \text{ GeV}$  (select real top)
- $H_T > 170 \text{ GeV}$  (discriminate against  $W$ +jets with softer jets)

### • W+jets CR == Anti-selection Region :

- Enriched by selecting events passing the PR criteria, but vetoing all SR requirements.
- PR-SR.

### • $t\bar{t}$ CR

- Passing all SR requirements, but requiring 2 b-tagged jets.



### • Preselection Region (PR):

#### • Exactly one tight charged lepton

- $p_T > 30 \text{ GeV}$ ,  $|\eta| < 2.5$
- Vetoing if existing a secondary high- $p_T$  ( $p_T > 30 \text{ GeV}$ ) charged loose leptons.

#### • Exactly 2 jets. Exactly 1 b-tagged.

- $p_T > 30 \text{ GeV}$  ( $p_T > 35 \text{ GeV}$  in transition region  $2.75 \leq |\eta| < 3.5$  to avoid mis-modeling between the central and forward calorimeters.)
- Spectator jet ( $|\eta| < 4.5$ ), b-jet (60 %WP (bin selection) within  $|\eta| < 2.5$ )

#### • MET > 35 GeV.

- $m_T(\text{lepton-MET})$  [or  $M_{tW}$ ] > 60 GeV.
- Additional multijet rejection (“triangular cut”)


## \*Event Selection

### • Multijet estimation



- Approach with data-driven methods.
- Data-driven jet-electron model and generic simulated di-jet events in the electron channel.
- Data-driven anti-muon model in the muon channel.

Scale factors


Process	$e$ -channel	$\mu$ -channel	$e+\mu$ -channels
$t$ -channel	$1.06 \pm 0.04$	$1.05 \pm 0.06$	$1.05 \pm 0.03$
$W$ +jets	$0.93 \pm 0.03$	$0.98 \pm 0.03$	$0.954 \pm 0.020$
$t\bar{t}, Wt, s$ -channel	$1.008 \pm 0.016$	$0.996 \pm 0.017$	$1.003 \pm 0.012$



**ATLAS Work in Progress**  
 $\sqrt{s} = 13 \text{ TeV}, 140.5 \text{ fb}^{-1}$   
 $t\bar{t}$  control region (post-fit)





**ATLAS Work in Progress**  
 $\sqrt{s} = 13 \text{ TeV}, 140.5 \text{ fb}^{-1}$   
 $W$ +jets control region (post-fi






# Control Plots: W+Jets CR

## Kinematics







# Control Plots: W+Jets CR

## Angular Observables






## Kinematics



# Control Plots: $t\bar{t}$ CR




## Angular Observables



# Top Polarization Measurement

## \*Template Fit

- **Octant Fit:** We slice the three-dimensional polarization phase space into 8 octants, and perform a template fit on the populations of the 8  $\cos \theta_i$  bins in these regions.
- **Asimov dataset** with an SM polarization is set up to develop the fitting procedure, and to predict the statistical and systematic uncertainties of the fit.
  - Asimov dataset: PolManip SM gen + total background (**no real data!**)
  - Input polarization:  $\vec{P}^t = (0.0, 0.0, +0.9)$ ,  $\vec{P}^{\bar{t}} = (-0.14, 0.0, -0.86)$



$P$ components	Expected value: stat.+syst. (stat.)
$P_x^t$	0.00 +0.09 / -0.09 ( $\pm 0.02$ )
$P_y^t$	0.00 +0.03 / -0.03 ( $\pm 0.01$ )
$P_z^t$	0.90 +0.13 / -0.13 ( $\pm 0.02$ )
$P_x^{\bar{t}}$	-0.14 +0.16 / -0.16 ( $\pm 0.03$ )
$P_y^{\bar{t}}$	0.00 +0.04 / -0.04 ( $\pm 0.02$ )
$P_z^{\bar{t}}$	-0.86 +0.18 / -0.17 ( $\pm 0.04$ )


# Top Polarization Measurement

## \*Template Fit

- **Systematics dominated measurement:** JES, JER and MET are the main sources of the systematic uncertainties. This is expected because:
  - Spectator jet defines our spin axis.
  - Top quark rest frame is reconstructed from both jets and the neutrino, which means the systematics JET JER and MET have a major impact on this determination.

$P$ components	Expected value: stat.+syst. (stat.)
$P_x^t$	0.00 +0.09 / -0.09 ( $\pm 0.02$ )
$P_y^t$	0.00 +0.03 / -0.03 ( $\pm 0.01$ )
$P_z^t$	0.90 +0.13 / -0.13 ( $\pm 0.02$ )
$P_x^{\bar{t}}$	-0.14 +0.16 / -0.16 ( $\pm 0.03$ )
$P_y^{\bar{t}}$	0.00 +0.04 / -0.04 ( $\pm 0.02$ )
$P_z^{\bar{t}}$	-0.86 +0.18 / -0.17 ( $\pm 0.04$ )

Polarization vector should be located inside the physical region represented by this unit circle.



Retrieved the polarization of the Asimov dataset

- A polarization measurement of the top quark and antiquark (separately) based on template fit method with full Run II data is presented.
- Signal and background are well modelled.
- The fitting procedure has shown to be robust through fitting Asimov dataset. Looking forward to unblinding.
- Results with Run II 13 TeV data expected soon!

**Thank you!**





# Backup

Backup