

A strange program for the LHC


Diego Martínez Santos

GAIN

(on behalf of the LHCb Collaboration)

Introduction


- LHCb experiment at LHC
 - Designed mostly for **b** and **c** decays
 → ~zero trigger efficiency otherwise
 - But there is also an ~infinite **strangeness** production at LHC (kaon xs ~ 1.2 **barn**)
 - Infinite production times zero efficiency requires L'Hopital
 - In 2011 we managed to get world best result in $K_S \rightarrow \mu\mu$
 - Major improvements in the trigger for **s** decays done for Run-II (2016-2018), and ongoing for Upgrade (≥ 2021)



Trigger system: status and prospects

L0
 (Hardware)

Main bottleneck for K. Can't be changed



Typical PT

~30-40 GeV



HLT1
 (Software)

Not designed for K, but flexible.

B-physics

~ 1-2 GeV

HLT2
 (Software)

K triggers being implemented


s-physics

~0.08 GeV

Trigger system: status and prospects

L0
 (Hardware)

Main bottleneck for K. Can't be changed



Typical PT

~30-40 GeV



HLT1
 (Software)

Not designed for K, but flexible.

B-physics

~ 1-2 GeV

HLT2
 (Software)

K triggers being implemented

s-physics


~0.08 GeV

$\epsilon(2011-2012) \sim 1-2\%$

$\epsilon(\text{Run-II})$ improved HLT $\sim 18\%$ (dimuons)

Maximum allowed by L0 $\sim 30\%$

Trigger system: status and prospects




$\epsilon(2011-2012) \sim 1-2\%$

$\epsilon(\text{Run-II})$ improved HLT $\sim 18\%$ (dimuons)

Maximum allowed by L0 $\sim 30\%$

Trigger system: status and prospects



Main bottleneck for K. Can't be changed



Not designed for K, but flexible.



K triggers being implemented

LHCb Upgrade



(Note: This logo may not be official)




$\epsilon(2011-2012) \sim 1-2\%$
 $\epsilon(\text{Run-II})$ improved HLT $\sim 18\%$ (dimuons)
 Maximum allowed by L0 $\sim 30\%$


$\epsilon(\text{Upgrade}) \sim 80-100\%$?
 Simulation studies show that rate would be under control

$K_S \rightarrow \mu\mu$: motivation

- SM prediction: $BR(K_S \rightarrow \mu\mu) = (5.18 \pm 1.50_{LD} \pm 0.02_{SD}) \times 10^{-12}$
JHEP05(2018) 024, JHEP 0401 (2004) 009, NPB 366 (1991) 189
- $K_S \rightarrow \mu\mu$ sensitive to different physics than $K_L \rightarrow \mu\mu$, NP can be bigger than SM by ~ 1 order of magnitude or even saturate current EXP limit







Example of a SUSY scenario from V.Chobanova et al., JHEP05(2018) 024



Leptoquark scenarios from Bobeth & Buras, JHEP02(2018)101

$K_S \rightarrow \mu\mu$ latest result

LHCb-CONF-2019-002



[LHCb-CONF-2019-002]

Full LHCb dataset analysed (9 fb⁻¹)


Benefits from huge (~1 order of magnitude) improvements in trigger for Run II

P_T muon thresholds at HLT: 80 MeV

No evidence for signal (1.4σ)

$K_S \rightarrow \mu\mu$ latest result

LHCb-CONF-2019-002



Full LHCb dataset analysed (9 fb⁻¹)

No evidence for signal (1.4σ)

New world best upper limit
(improved by a factor ~4)

$BR(K_S \rightarrow \mu\mu) < 2.1 \times 10^{-10}$
@ 90% CL

Sept. 2019

At 1σ: $B(K_S^0 \rightarrow \mu^+\mu^-) = 0.94_{-0.64}^{+0.72} \times 10^{-10}$



The HyperCP evidence

- The HyperCP collaboration found evidence for $\Sigma \rightarrow p \mu \mu$ decays, and provided a BR:

$$\mathcal{B}(\Sigma^+ \rightarrow p \mu^+ \mu^-) = (8.6_{-5.4}^{+6.6} \pm 5.5) \cdot 10^{-8}$$

[Phys.Rev.Lett. 94 (2005) 021801]



- Consistent w/ SM: $1.6 < \text{BR}[x10^{-8}] < 9$
X G He et al, PRD 72 (2005) 074003
- This evidence had wide relevance since all 3 observed events had the same dimuon invariant mass (214 MeV)
- Suggested the existence of a new neutral particle at that mass



$\Sigma \rightarrow p\mu\mu$



LHCb-PAPER-2017-049
 arXiv:1712.08606
 PRL 120, 221803 (2018)

- **Current result $\Sigma \rightarrow p\mu\mu$** : Found 4σ evidence $\text{BR}(\Sigma \rightarrow p\mu\mu) : 2.1^{+1.6}_{-1.2} \times 10^{-8}$, no evidence of resonant dilepton state
- **Run-II**: We expect ~ 150 signal events \rightarrow measure AFB
- **Upgrade(s)**: Full differential decay rate



10y ago we thought this channel was \sim impossible and instead now we are even thinking on an amplitude analysis....

$K_S \rightarrow \pi^0 \mu\mu$ sensitivity study



LHCb-upgrade

Phase-II-upgrade? →

$|a_S| = 1.2 \pm 0.2$ from NA48 fixing b_S from VMD
PLB599 (2004) 197-211,



Table 4: Projected statistical uncertainties on a_S under various conditions.

Configuration	Phase I	Phase II
BR & q^2 fit	0.25	0.10
BR & q^2 fit with NA48 constraint	0.19	0.10
BR & q^2 fit fixing b_S	0.06	0.024
a_S measurement from BR alone (fixing b_S)	0.06	0.024

Much more bkg than $K_S \rightarrow \mu\mu$,
but also 1000x more signal

$K_S \rightarrow \gamma\mu\mu$?

$K_S \rightarrow \pi^0\mu\mu$ analysis can also be extended to other neutrals, eg: $K_S \rightarrow \gamma\mu\mu$
 But harder to separate from $K_S \rightarrow \pi\pi$ as the mass of the neutral gets lighter
 (unless a cut on the energy is used)



Semileptonic decays


- Semileptonic Hyperon Decays (SHD)

Very interesting in view of LUV hints in semileptonic B decays

Many muonic modes have still very poor precision (20%, 100%)

- ☺ High BR (10^{-4}): Massive yields in LHCb acceptance

$$R_{B_1 B_2}^{\text{NP}} \simeq \frac{\left(\epsilon_S^{s\mu} \frac{f_S(0)}{f_1(0)} + 12 \epsilon_T^{s\mu} \frac{g_1(0)}{f_1(0)} \frac{f_T(0)}{f_1(0)} \right)}{\left(1 - \frac{3}{2} \delta \right) \left(1 + 3 \frac{g_1(0)^2}{f_1(0)^2} \right)} \Pi(\Delta, m_\mu)$$



(extrapolations from 1412.8484)

Gonzalez-Alonso & JMC, NA62 Physics Handbook

<https://indico.cern.ch/event/590880/contributions/2485320/>

Semileptonic decays

- Semileptonic Hyperon Decays (SHD)


Very interesting in view of LUV hints in semileptonic B decays

Many muonic modes have still very poor precision (20%, 100%)

- 😊 High BR (10^{-4}): Massive yields in LHCb acceptance
- 😞 Challenging peaking backgrounds:


For each
 $B1 \rightarrow B2 \mu\nu$ there is always a
 $B1 \rightarrow B2\pi$ (inc. $\rightarrow B2\mu\nu$)

😊 Can be separated in search planes



Semileptonic decays

- Semileptonic Hyperon Decays (SHD)



Expected $O(7k)$ signal events per $\text{fb}^{-1} \rightarrow$ very good stat precision

Semileptonic decays

- Semileptonic Hyperon Decays (SHD)

Very interesting in view of LUV hints in semileptonic B decays


Many muonic modes have still very poor precision (20%, 100%)

- 😊 High BR (10^{-4}): Massive yields in LHCb acceptance
- 😞 Challenging peaking backgrounds:

For each
 $B1 \rightarrow B2 \mu \nu$ there is always a
 $B1 \rightarrow B2 \pi$ (inc. $\rightarrow B2 \mu \nu$)

😊 Can be separated in search planes

Fast Simulation



Lepton Flavour Violation

- Lepton Flavour Violation is a hot topic nowadays

LHCb can do:

$$K_S \rightarrow e\mu$$

No limit exists so far

$$K_L \rightarrow e\mu < 4.7 \times 10^{-12} \text{ BNL, } PRL \text{ 81 (1998) 5734-5737}$$

$K_S \rightarrow e\mu$ is a LFV model discriminator


Lepton Flavour Violation

- Lepton Flavour Violation is a hot topic nowadays

LHCb can do:

$$K_S \rightarrow e\mu$$

$$K^+ \rightarrow \pi^+\mu^-e^+$$



Lepton Flavour Violation


- Lepton Flavour Violation is a hot topic nowadays

LHCb can do:


$K_S \rightarrow e\mu$

$K^+ \rightarrow \pi^+\mu^-e^+$

Maybe $K^+ \rightarrow \pi^+\mu^+e^-$





Competition w/
NA62 to be clarified




Charged kaons

- K^+ mass in $K \rightarrow 3\pi$
- Under study sensitivity to $K^+ \rightarrow \pi^+ \mu \mu$ vs NA62
- Benefits from the new dimuon triggers (the same way as $K_S \rightarrow \mu \mu$)



B and L violation (very low priority)

CLAS collaboration (Jefferson Lab): Limits on B and L violation



Reaction	\mathcal{B}_{UL}
$\Lambda \rightarrow K^+ e^-$	2×10^{-6}
$\Lambda \rightarrow K^+ \mu^-$	3×10^{-6}
$\Lambda \rightarrow K^- e^+$	2×10^{-6}
$\Lambda \rightarrow K^- \mu^+$	3×10^{-6}
$\Lambda \rightarrow \pi^+ e^-$	6×10^{-7}
$\Lambda \rightarrow \pi^+ \mu^-$	6×10^{-7}
$\Lambda \rightarrow \pi^- e^+$	4×10^{-7}
$\Lambda \rightarrow \pi^- \mu^+$	6×10^{-7}
$\Lambda \rightarrow \bar{p}\pi^+$	9×10^{-7}
$\Lambda \rightarrow K_S^0 \nu$	2×10^{-5}

[arXiv:1507.03859](https://arxiv.org/abs/1507.03859) [hep-ex]

We can easily do many of CLAS' decays

...as well as others:

- $\Sigma \rightarrow 3\mu$
- $\Lambda \rightarrow \pi 3\mu$

...and many other crazy (J conserving) combinations.

Currently very low priority, since we assume that BSM contributions can only be as much as $BR \sim 10^{-56}$

Conclusions




Conclusions

- **s** decays are awesome
 - High interest for BSM
 - Ultimate experimental precision $\sim 10^{-11}$ - 10^{-12}
- There is an LHC**s** community in the LHC**b** village
 - Trigger is constantly improving
 - We aim for LHCb upgrade to reach efficiencies **s** as high as for **b**'s
- Full LHCb dataset result : $K_S \rightarrow \mu\mu$
- Run-II (2016-2018) data analysis ongoing $\Sigma \rightarrow p\mu\mu$, $K_S \rightarrow (\gamma/\pi^0)\mu\mu\dots$

Backup


$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) = 2.65_{-0.39}^{+0.43} \times 10^{-9}, \quad \text{BR}(B_d \rightarrow \mu^+ \mu^-) = 1.09_{-0.68}^{+0.74} \times 10^{-10}.$$



<https://arxiv.org/pdf/1904.08399.pdf>

Strangeness decays

- So far a kaons showed great success on indirect searches: $c, b, t, CKM \dots$
- High theoretical interest, most notably to test departures from MFV paradigm (eg, flavor generic)



- Useful to understand “Hints” for BSM in b sector
 - Eg: deviations in $b \rightarrow s \mu \mu$: are they replicated in $s \rightarrow d \mu \mu$?
 - Y \rightarrow interesting
 - N \rightarrow interesting
- Potentially immense samples : high(est) ultimate experimental precision

Efficiencies

* More details in: arXiv:1808.03477 [hep-ex]

Channel	$\chi_s/\chi_s(K_S)$	eff/eff(K_S)	eff/eff(K_S) w/ Downstream tracks	Mass resolution	
				σ_L (MeV/ c^2)	σ_D (MeV/ c^2)
$K_S^0 \rightarrow \mu^+ \mu^-$	1	1.0 (1.0)	1.8 (1.8)	~ 3.0	~ 8.0
$K_S^0 \rightarrow \pi^+ \pi^-$	1	1.1 (0.30)	1.9 (0.91)	~ 2.5	~ 7.0
$K_S^0 \rightarrow \pi^0 \mu^+ \mu^-$	1	0.93 (0.93)	1.5 (1.5)	~ 35	~ 45
$K_S^0 \rightarrow \gamma \mu^+ \mu^-$	1	0.85 (0.85)	1.4 (1.4)	~ 60	~ 60
$K_S^0 \rightarrow \mu^+ \mu^- \mu^+ \mu^-$	1	0.37 (0.37)	1.1 (1.1)	~ 1.0	~ 6.0
$K_L^0 \rightarrow \mu^+ \mu^-$	~ 1	$2.7 (2.7) \times 10^{-3}$	0.014 (0.014)	~ 3.0	~ 7.0
$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	~ 2	$9.0 (0.75) \times 10^{-3}$	$41 (8.6) \times 10^{-3}$	~ 1.0	~ 4.0
$K^+ \rightarrow \pi^+ \mu^+ \mu^-$	~ 2	$6.3 (2.3) \times 10^{-3}$	0.030 (0.014)	~ 1.5	~ 4.5
$\Sigma^+ \rightarrow p \mu^+ \mu^-$	~ 0.13	0.28 (0.28)	0.64 (0.64)	~ 1.0	~ 3.0
$\Lambda \rightarrow p \pi^-$	~ 0.45	0.41 (0.075)	1.3 (0.39)	~ 1.5	~ 5.0
$\Lambda \rightarrow p \mu^- \bar{\nu}_\mu$	~ 0.45	0.32 (0.31)	0.88 (0.86)	—	—
$\Xi^- \rightarrow \Lambda \mu^- \bar{\nu}_\mu$	~ 0.04	$39 (5.7) \times 10^{-3}$	0.27 (0.09)	—	—
$\Xi^- \rightarrow \Sigma^0 \mu^- \bar{\nu}_\mu$	~ 0.03	$24 (4.9) \times 10^{-3}$	0.21 (0.068)	—	—
$\Xi^- \rightarrow p \pi^- \pi^-$	~ 0.03	0.41(0.05)	0.94 (0.20)	~ 3.0	~ 9.0
$\Xi^0 \rightarrow p \pi^-$	~ 0.03	1.0 (0.48)	2.0 (1.3)	~ 5.0	~ 10
$\Omega^- \rightarrow \Lambda \pi^-$	~ 0.001	$95 (6.7) \times 10^{-3}$	0.32 (0.10)	~ 7.0	~ 20

Sensitivity of (semi)leptonic kaon decays in a nutshell

- $K_{\ell 3}$

$$\Gamma(K_{\ell 3}(\gamma)) = \underbrace{\frac{G_F^2 m_K^5}{192\pi^3}}_{\text{Measured in } \mu \text{ decay}} \underbrace{|\tilde{V}_{us}^\ell|^2 f_+(0)^2}_{\left(1 + \epsilon_L^{s\ell} + \epsilon_R^s - \tilde{V}_L\right) V_{us}^{\text{SM}}} \underbrace{I_K^\ell(\lambda_{+,0}, \epsilon_S^{s\ell}, \epsilon_T^{s\ell})}_{\text{Phase-space Int.}} \underbrace{\left(1 + \delta^c + \delta_{\text{em}}^{c\ell}\right)^2}_{\text{Rad. and isosp. corr.}}$$


- $K_{\ell 2}$

$$\Gamma_{K_{\ell 2}(\gamma)} = \frac{G_F^2 m_K m_\ell^2}{8\pi} \left(1 - \frac{m_\ell^2}{m_P^2}\right)^2 |\tilde{V}_{us}^\ell|^2 f_{K^\pm}^2 \left(1 - 4\epsilon_R^s - \underbrace{\frac{2B_0}{m_\ell} \epsilon_P^{s\mu}}_{\chi \text{ enh.}}\right)$$

- ▶ $|\tilde{V}_{us}^\ell|$ only accessible through CKM unitarity and LUV tests
- ▶ ϵ_R^s cannot be completely disentangled from $\epsilon_P^{s\ell}$
- ▶ $\epsilon_{S,T}^{s\ell}$ accessible through the spectra/angular distribution

Kaon decays alone cannot disentangle all NP possibilities

$K_S \rightarrow \pi^+ \pi^- e e$ sensitivity study



Based on simulation:

Expected a signal yield of

$$N = 120^{+280}_{-100}$$


For the full Run-I dataset

Expected background yield
is not well known yet

K⁰ tagging?

$pp \rightarrow K^0 K^- X$, $pp \rightarrow K^{*+} X \rightarrow K^0 \pi^+ X$ and $pp \rightarrow K^0 \Lambda^0 X$.

Toy MC for 50fb^{-1}



Lifetime acceptance and $K_L \rightarrow \mu\mu$ background

K_L and K_S are distinguishable only by the decaytime...

... and that is in theory. In practice, LHCb decaytime acceptance is not great for kaons

$$\epsilon(t) \sim e^{-\beta t} \quad \text{With } \beta \gtrsim 5 \times \Gamma_S (\gg \Gamma_L).$$

This makes the two lifetime distributions to look similar

But the overall efficiency ratio is of course different


$$\frac{\epsilon_{K_L^0}}{\epsilon_{K_S^0}} = \frac{\Gamma_L \int_{0.1\tau_S}^{1.45\tau_S} e^{-t(\Gamma_S+\beta)} dt}{\Gamma_S \int_{0.1\tau_S}^{1.45\tau_S} e^{-t(\Gamma_L+\beta)} dt} \approx 2.2 \times 10^{-3}$$

And makes $K_L \rightarrow \mu\mu$ to become a negligible background for the current level of precision


But can be relevant when we approach the 10^{-11} level

$$\beta \sim 86 \text{ ns}^{-1}$$

LHCb 2015 Trigger Diagram



LHCb Upgrade Trigger Diagram



Normalization of event yield

Converting a signal yield into a branching ratio

$$N(K_S^0 \rightarrow \pi\mu\mu) = \overset{K_S^0 \text{ production crosssection}}{\sigma(K_S^0)} BR(K_S^0 \rightarrow \pi\mu\mu) \overset{\text{Absolute efficiency}}{\epsilon} \overset{\text{Integrated luminosity}}{L}$$

How? (normalization of event yield)

Converting a signal yield into a branching ratio

$$N(K_S^0 \rightarrow \pi\mu\mu) = \overset{K_S^0 \text{ production crosssection}}{\sigma(K_S^0)} BR(K_S^0 \rightarrow \pi\mu\mu) \overset{\text{Absolute efficiency}}{\epsilon} \overset{\text{Integrated luminosity}}{L}$$

$$\frac{N(K_S^0 \rightarrow \pi\mu\mu)}{N(K_S^0 \rightarrow \pi\pi)} = \frac{\cancel{\sigma(K_S^0)} BR(K_S^0 \rightarrow \pi\mu\mu) \cancel{\epsilon} L}{\cancel{\sigma(K_S^0)} BR(K_S^0 \rightarrow \pi\pi) \cancel{\epsilon'} L}$$

Introduce in the ntuples a $K_S^0 \rightarrow \pi\pi$ decays counter

Very well known
(69.20±0.05)%

Dilepton mass distribution

Take formulae from hep-ph/9808289

$$\frac{d\Gamma}{dz} = \frac{\alpha^2 M_K}{12\pi(4\pi)^4} \lambda^{3/2}(1, z, r_\pi^2) \sqrt{1 - 4\frac{r_\ell^2}{z}} \left(1 + 2\frac{r_\ell^2}{z}\right) |W(z)|^2, \quad (3)$$

$$z = m^2 \rightarrow d\Gamma/dm = 2m d\Gamma/dz$$


$$W_i(z) = G_F M_K^2 (a_i + b_i z) + W_i^{\pi\pi}(z), \quad (11)$$

$$W_i^{\pi\pi}(z) = \frac{1}{r_\pi^2} \left[\alpha_i + \beta_i \frac{z - z_0}{r_\pi^2} \right] F(z) \chi(z),$$

Remind of Bmm sensitivity


B mesons

We check that we get right the expected increase of B meson yields (i.e, a factor ~ 2)



D mesons

For D mesons the increase is slightly smaller ($\sim 1.6- 1.7$)




Strange particles

Increase for most of them is ~40%

A bit less for baryons (note: baryons, not anti-baryons)

However, the momentum is also different w.r.t 7 TeV.




In particular, for the K0s decaying in the VELO the increase is "only" ~30% → This is the number we really care for Ks → μμ studies


Leptons

Increase in tau yield consistent with ~ 2 , expected by the fact that most of them come from b 's and c 's


Check with more stats if the asymmetry +/- is still there




→ the long-distance (LD) contributions:



→ the short-distance (SD) contributions:






$K_S \rightarrow \pi^0 \mu \mu$ sensitivity study


The background discrimination

V. Chobanova et al,
CERN-LHCb-PUB-2016-017

BDT response FULL



BDT response PARTIAL



- As usual: BDT trained against combinatorial background
- Specific backgrounds: $K_S \rightarrow \pi \pi$, $K_L \rightarrow \pi \pi \pi$, $K_{S/L} \rightarrow \mu \mu \gamma \gamma$ (negligible)





Don't affect the sensitivity estimate

$K_S \rightarrow \pi^0 \mu \mu$ sensitivity study

Fit, FULL



V. Chobanova et al,
CERN-LHCb-PUB-2016-017







$K_S \rightarrow \pi^0 \mu \mu$ sensitivity study

Fit, PARTIAL





V. Chobanova et al,
CERN-LHCb-PUB-2016-017




Strangeness production/detection at LHCb

- The pp collisions @ LHC produce a 'kaon flux' of 10^{13} K_S per fb^{-1} of luminosity in the LHCb acceptance
- Charged decay products can be reconstructed using Long Tracks or Downstream Tracks
- We use Long Tracks for R_{nS}
- Downstream will be investigated (extra yield, but worse reconstruction quality)




K_L, K_S produced in equal amounts. Acceptance ratio is $\sim 2 \times 10^{-3}$ (for Long Tracks)

Ongoing stuff



K⁺ studies



Large samples of charged kaon decays are available



K⁺ mass is not very well known

K⁺ → πμμ ?

$$K_S \rightarrow X^0 \mu \mu$$

- The $K_S \rightarrow \pi^0 \mu \mu$ PARTIAL analysis can be recasted for general/inclusive $K_S \rightarrow X^0 \mu \mu$. With X being whatever neutral system:
 - $K_S \rightarrow \gamma \mu \mu$. Can also be completed with photon reconstruction
 - $K_S \rightarrow (1+1-) \mu \mu$. Some of them are also being searched for explicitly
 - Some exotic, eg, 17 MeV neutral boson of Phys. Rev. Lett. 116, 042501 (2016)

Limits can be provided as a function of X^0 mass




$K_S \rightarrow \mu\mu$ full Run-I analysis

[arXiv:1706.00758](https://arxiv.org/abs/1706.00758) [hep-ex]

- Analysed full Run-I (2011-2012) data
- Events classified using a BDT trained against combinatorial background
- Dedicated muon identification algorithm trained against $K_S \rightarrow \pi\pi$
- Mass resolution 4 MeV

Background



$K_L \rightarrow \mu\mu$ negligible: (down to 10^{-11} precision)


$K \rightarrow \pi\mu\nu$: negligible

$\Lambda \rightarrow p\pi$ removed by a cut in the Armenteros-Podolanski plot.

- **Combinatorial background**
- **$K_S \rightarrow \pi\pi$ double misid**

$K_S \rightarrow \pi^+ \pi^- e e$ sensitivity study

C.Marin et al,
CERN-LHCb-PUB-2016-016



Based on simulation:

Expected a signal yield of

$$N = 120_{-100}^{+280}$$

For the full Run-I dataset

Expected background yield
is not well known yet

Why? ($K_S \rightarrow \pi^0 \mu \mu$ and SM errors on $K_L \rightarrow \pi^0 \mu \mu$)

$$\mathcal{B}(K_L \rightarrow \pi^0 \mu^+ \mu^-)_{\text{SM}} = \{1.4 \pm 0.3, 0.9 \pm 0.2\} \cdot 10^{-11}$$

$$\mathcal{B}(K_L \rightarrow \pi^0 l^+ l^-) = (C_{\text{dir}}^l \pm C_{\text{int}}^l |a_S| + C_{\text{mix}}^l |a_S|^2 + C_{\gamma\gamma}^l + C_S^l) \cdot 10^{-12}$$

$$|a_S| = 1.20 \pm 0.20$$

Dominant uncertainty, that makes difficult potential BSM interpretation of $K_L \rightarrow \pi^0 \mu \mu$

$$C_{\text{dir}}^e = (4.62 \pm 0.24) [(\text{Im} Y_A)^2 + (\text{Im} Y_V)^2],$$

$$C_{\text{int}}^e = (11.3 \pm 0.3) \text{Im} Y_V,$$

$$C_{\text{mix}}^e = 14.5 \pm 0.5,$$

$$C_{\gamma\gamma}^e \approx C_S^e \approx 0,$$

$$C_{\text{dir}}^\mu = (1.09 \pm 0.05) [2.32 (\text{Im} Y_A)^2 + (\text{Im} Y_V)^2]$$

$$C_{\text{int}}^\mu = (2.63 \pm 0.06) \text{Im} Y_V,$$

$$C_{\text{mix}}^\mu = 3.36 \pm 0.20,$$

$$C_{\gamma\gamma}^\mu = 5.2 \pm 1.6,$$

$$C_S^\mu = (0.04 \pm 0.01) \text{Re} Y_S + 0.0041 (\text{Re} Y_S)^2.$$

It comes from the **experimental uncertainty** on $\text{BR}(K_S \rightarrow \pi^0 \mu \mu)$ measured by NA48


$K_S^0 \rightarrow \pi^0 \mu^+ \mu^-$	NA48	$(2.9^{+1.5}_{-1.2}) \times 10^{-9}$
---------------------------------------	------	--------------------------------------

~50% relative error

Improved measurements of $\text{BR}(K_S \rightarrow \pi^0 \mu \mu)$ will translate into improved BSM constraints from $K_L \rightarrow \pi^0 \mu \mu$

$K_S \rightarrow \mu\mu$ prospects

Run- I: $BR < 8 (10) \times 10^{-10}$ @90(95)%CL



- Extrapolating from Run-I result
- Full Run-II analysis ongoing: expected to improve by a **factor 4 to 10** Run-I's sensitivity
 - Better trigger
 - Better reco/selection
- Future: start to investigate tagged decays, which would allow to access NP in the K_S - K_L interference
 [D'Ambrosio&Kitahara PRL 119, 201802 (2017)]

Could well become the strongest limit on a BR by an LHC experiment