

# Azimuthal anisotropy in 5.02 TeV Pb+Pb and 5.44 TeV Xe+Xe collisions with the ATLAS experiment

XIV Polish Workshop  
on Relativistic Heavy-Ion Collisions

Kraków, Poland 6-7 April 2019





Klaudia Burka  
AGH UST



# Motivation


- **Azimuthal anisotropy** results from different pressure gradients in different spatial directions



## Particle azimuthal distribution

**Singles:** 
$$\frac{dN}{d\phi} \propto 1 + \sum_n 2v_n \cos[n(\phi - \Phi_n)]$$


**Pairs:** 
$$\frac{dN}{d\Delta\phi} \propto 1 + \sum_n 2v_n^a v_n^b \cos[n(\Delta\phi)]$$



- Pb+Pb 5.02 TeV  $\rightarrow 0.49\text{nb}^{-1}$
- Xe+Xe 5.44 TeV  $\rightarrow 3\mu\text{b}^{-1}$

# ATLAS detector $\rightarrow v_n$ measurement

## ATLAS detector




- Pb+Pb 5.02 TeV  $\rightarrow 0.49\text{nb}^{-1}$
- Xe+Xe 5.44 TeV  $\rightarrow 3\mu\text{b}^{-1}$

### Inner Detector (Pixel+SCT)

Flow measurements is based on charged tracks reconstructed in ID

- $|\eta| < 2.5$
- $2\pi$   $\phi$  acceptance
- $p_T > 0.5$  GeV




Pixel detector

SCT detector

# ATLAS detector $\rightarrow$ $v_n$ measurement




## ATLAS detector



- Pb+Pb 5.02 TeV  $\rightarrow$   $0.49\text{nb}^{-1}$
- Xe+Xe 5.44 TeV  $\rightarrow$   $3\mu\text{b}^{-1}$

### Forward Calorimeter ( $3.2 < |\eta| < 4.9$ )

- Flow vectors
- Centrality definition




# Methodology

## ✧ Two-particle correlations (2PC) and Scalar-product (SP) methods


- Correlating tracks with Q-vectors at forward rapidities
- Non-flow suppressed by large  $\Delta\eta$ -gap ( $\Delta\eta > 2$ )

arXiv:0809.2949 [nucl-ex]



$$u_n, Q_n = \sum_j w_j e^{in\phi_j}$$

$$v_2^2\{2\} = \langle v_2 \rangle^2 + \sigma_v^2$$



## ✧ Multi-particle cumulants:

- Correlating tracks at mid-rapidity with each other
- Analytically suppress non-flow
- Sensitive to flow fluctuations

$$v_2^2\{4\} \approx \langle v_2 \rangle^2 - \sigma_v^2$$


$$v_n\{2\} = \sqrt[2]{\langle v_n^2 \rangle},$$

$$v_n\{4\} = \sqrt[4]{2\langle v_n^2 \rangle^2 - \langle v_n^4 \rangle},$$


$$v_n\{6\} = \sqrt[6]{\langle v_n^6 \rangle - 9\langle v_n^2 \rangle \langle v_n^4 \rangle + 12\langle v_n^2 \rangle^3}$$

# $v_n$ harmonics in Pb+Pb collisions $\rightarrow p_T$


## ULTRA-CENTRAL



## MID-CENTRAL




## PERIPHERAL



- ✧  $v_n$  measured up to  $p_T = 60$  GeV  $\rightarrow v_2(p_T)$  positive at highest  $p_T$   
 $\rightarrow$  provide information about parton energy loss
- ✧ The ordering:  $v_n > v_{n+1}$  in mid-central and peripheral collisions
  - $v_3 > v_4 > v_5 \approx v_2$  for the most central collisions at  $p_T = 3-5$  GeV
- ✧ The  $v_7$  harmonic is found to be non-zero for centralities 0-50%

# $v_n$ harmonics in Pb+Pb collisions $\rightarrow \eta$





- ✧  $v_n$  flow harmonics show very weak  $\eta$ -dependence in the range of  $|\eta| < 2.5$
- ✧ The exceptions are peripheral collisions and  $v_3$ ,  $v_4$  harmonics for which the differences between  $v_n(\eta=0)$  and  $v_n(\eta=2.5)$  become significant and reach about 10-25%

Eur. Phys. J. C (2018) 78: 997

# $v_n$ harmonics in Pb+Pb collisions $\rightarrow$ centrality

✧ Elliptic flow is strongly dependent on event centrality and is largest in mid-central events  $N_{\text{part}} = 70-110$

✧ Higher order  $v_n$  show weak centrality dependence






# Methods comparison in Pb+Pb collisions





## SCALAR-PRODUCT vs. EVENT-PLANE

- ✧ The SP method  $\rightarrow$  always  $\sqrt{\langle v_n^2 \rangle}$
- ✧ The EP method  $\rightarrow$  values between  $\langle v_n \rangle$  and  $\sqrt{\langle v_n^2 \rangle}$
- ✧ A small difference seen for  $v_2$ 
  - Largest for 20-50% centralities reaching  $\sim 3\%$
- ✧ For  $n > 2$  the EP and SP results are consistent



# Methods comparison in Pb+Pb collisions

## SCALAR-PRODUCT vs. TWO-PARTICLE CORRELATION




✧  $v_2 \rightarrow$  the largest differences seen in 0-5%

- It decreases with centrality
- $v_2\{2PC\}$  and  $v_2\{SP\}$  matches within  $\sim 2\text{-}5\%$  for mid-central collisions

✧  $v_3 \rightarrow$  match within  $\sim 5\%$  for  $p_T < 4$  GeV

✧ Differences due to the factorization breakdown


# Universal scaling of $v_n$ harmonics






- ✧ Simultaneous scaling along the  $p_T$  and  $v_n$  axes was performed
  - Universal shapes for the  $v_n$  ( $n=2,3$ ) across the different centrality classes
  - Similarity in properties of the QGP evolving from different initial conditions

# Azimuthal anisotropy in Xe+Xe collisions

## CENTRAL



## MID-CENTRAL



✧  $v_n$  measured up to  $n=5$  in a wide  $p_T$  range (20 GeV for  $v_2$ )

✧  $v_2$  is dominant except the most central collisions

✧  $v_n$  measured with higher order correlations smaller



● suppressed non-flow

● impact of fluctuations

ATLAS-CONF-2018-011

# Initial state fluctuations in Xe+Xe & Pb+Pb collisions

ATLAS-CONF-2018-011





✧ Integrated  $v_n$  higher in the most central events for Xe+Xe

- Smaller collision system  $\rightarrow$  larger initial fluctuations

✧ Reduced  $v_n\{\text{Xe+Xe}\}$  value in mid-central and peripheral

- Viscous effects



✧ Ratio  $v_n\{\text{Xe+Xe}\}/v_n\{\text{Pb+Pb}\}$  consistent with theoretical predictions: Giacalone *et al.* Phys.Rev.C97,034904(2018)



# Flow fluctuations in Xe+Xe collisions

✧ Comparisons of 2PC/SP to cumulant results:

- $v_2\{2PC/SP\} \gg v_2\{4\}$  (central collisions)
- $v_3\{2PC/SP\} \gg v_3\{4\}$

✧ Indicate strong flow fluctuations

✧ The ordering  $v_2\{2PC\} > v_2\{4\} \approx v_2\{6\}$  is observed, which indicates that  $v_2$  fluctuations are close to Gaussian



$$v_n\{2\} = \sqrt{\bar{v}_n^2 + \delta_n^2}, \quad v_n\{4\} = v_n\{6\} = \bar{v}_n$$

- ✧ If  $v_2 \sim$  Gaussian:  $v_2\{6\}/v_2\{4\} = 1$
- ✧  $v_2$  in Xe+Xe deviates further from Gauss  $\rightarrow$  deformed nucleus



# Summary

- ✧ New results from Pb+Pb and Xe+Xe collisions are obtained (Eur. Phys. J. C (2018) 78: 997, ATLAS-CONF-2018-011)
- ✧ Thanks to the excellent ATLAS detector and rich datasets:
  - Measured flow harmonics up  $v_7$  and to a very high  $p_T$  in Pb+Pb
  - Performed a comprehensive study of flow in Xe+Xe collisions at 5.44 TeV and compared to Pb+Pb at 5.02 TeV

**Thank you for your attention!**

This work was supported in part by the National Science Centre, Poland, grant no. 2016/23/N/ST2/01339 and by PL-Grid Infrastructure


# Universal scaling of $v_n$ harmonics – $v_3$



- ✧ Simultaneous scaling along the  $p_T$  and  $v_n$  axes was performed
  - Universal shapes for the  $v_n$  ( $n=2,3$ ) across the different centrality classes
  - Similarity in properties of the QGP evolving from different initial conditions




# Universal scaling of $v_n$ harmonics – scale factors



- ✧ the  $p_T$ -scale factors are quite comparable between  $v_2$  and  $v_3$ 
  - in the 0–10% most central events, larger difference  $\rightarrow$  could be due to larger jet-bias and factorization-breaking effects in the  $v_2$  as compared to  $v_3$
- ✧ the  $y$ -scale factors are different  $\rightarrow$  to be expected as the  $y$ -scale factors corresponds to the changing collision geometry, which becomes more and more elliptic from central to mid-central events resulting in a large increase in  $v_2$ , while  $v_3$ , which is driven by fluctuations, changes only gradually.


**Eur. Phys. J. C (2018) 78: 997**

# Xe+Xe/Pb+Pb flow harmonics ratios



ATLAS-CONF-2018-011

# Xe+Xe/Pb+Pb flow harmonics ratios



ATLAS-CONF-2018-011