



Contribution ID: 12

Type: **not specified**

## Measurements of single diffraction using forward proton tagging at ATLAS

*Monday, 24 June 2019 10:50 (30 minutes)*

Inclusive single diffractive dissociation ( $pp \rightarrow pX$ ) is studied using data collected by the ATLAS experiment at the LHC. The intact proton is reconstructed and measured in the ALFA forward spectrometer, while charged particles from the dissociative system (X) are reconstructed and measured using the ATLAS inner tracking detector and calorimeters. Differential cross sections are presented as a function of the proton fractional momentum loss, the four-momentum transfer squared, and the size of a rapidity gap measured from the edge of the ATLAS calorimeters. The results are interpreted in the framework of Regge phenomenology. If available, measurements of the charged particle multiplicity in single-diffractive events will also be presented, using protons reconstructed in the AFP detectors.

### Additional comments

**Primary authors:** ADAMCZYK, Leszek (AGH University of Science and Technology (PL)); ADAMCZYK, Leszek (AGH University of Science and Technology (PL))

**Presenters:** ADAMCZYK, Leszek (AGH University of Science and Technology (PL)); ADAMCZYK, Leszek (AGH University of Science and Technology (PL))

**Session Classification:** Session 1

**Track Classification:** Diffraction and Central Exclusive Production