


Improved constraints on a t-channel simplified model of Majorana dark matter


Kirtimaan A. Mohan
Michigan State University

In collaboration with
Dipan Sengupta, Tim Tait, Bin Yan, C-P. Yuan
arXiv: 1903.05650, JHEP 1905 (2019) 115
University of Washington, 13, Aug, 2019

Theories of Dark Matter




Spectrum of Theory Space



From Tim Tait

Outline

- Objective: Take a Simplified model and calculate everything with better precision.
- Direct Detection constraints @ 1-loop.
- Include Renormalization Group Evolution effects.
- LHC constraints @NLO.
- Understand importance of improving precision. Is it worth the effort?



A Simplified Model

- Construction—Inspired by more complete models, consider models that contain dark matter as well as the most important mediator(s).
- Example—Consider a class of models in which dark matter interacts with quarks through colored scalar mediators—looks like the MSSM, but simpler with three parameters; dark matter mass, mediator mass, coupling strength.

$$\{M_\chi, M_{\tilde{q}_L}, g_{DM}\}$$

$$\mathcal{L}_{int} = \sum_{q=u,d,s,c,b,t} g_{DM} (\tilde{q}_L^* \bar{\chi} P_L q + h.c.)$$

- Dark matter can be Dirac or Majorana fermion.

The Lagrangians

- Colored scalar mediators interact with quarks and singlet dark matter.
- Three possible charges—corresponding to three possible models (u_R , d_R , q_L).
- Motivated by MFV we set all masses and couplings equal.

$$\mathcal{L}_{FV} = (\delta g_{DM} \tilde{u}^* Y^{u\dagger} Y^u \bar{\chi} P_R u + h.c.) + \delta m^2 \tilde{u}^* Y^{u\dagger} Y^u \tilde{u} + \mathcal{O}(Y^4)$$

$$(3, 1)_{2/3}$$

$$\mathcal{L}_{u_R} \supset \sum_{u=u,c,t} [(D_\mu \tilde{u})^* (D^\mu \tilde{u}) - M_{\tilde{u}}^2 \tilde{u}^* \tilde{u} + g_{DM} (\tilde{u}^* \bar{\chi} P_R u + \bar{u} P_L \chi)]$$

$$(3, 1)_{-1/3}$$

$$\mathcal{L}_{d_R} \supset \sum_{d=d,s,b} [(D_\mu \tilde{d})^* (D^\mu \tilde{d}) - M_{\tilde{d}}^2 \tilde{d}^* \tilde{d} + g_{DM} (\tilde{d}^* \bar{\chi} P_R d + \bar{d} P_L \chi)]$$

$$(3, 2)_{-1/6}$$


$$\mathcal{L}_{q_L} \supset \sum_{q=u,c,t,d,s,b} [(D_\mu \tilde{q})^* (D^\mu \tilde{q}) - M_{\tilde{q}}^2 \tilde{q}^* \tilde{q} + g_{DM} (\tilde{q}^* \bar{\chi} P_L q + \bar{q} P_R \chi)]$$



Rest of this talk— We will look at q_L model with majorana fermions.


Before we dive into the details...

Leading Order Constraints



Constraints Dominated by LHC

Improved Constraints



Constraints Dominated by SI Direct Detection

Constraints improve by an order of magnitude in some places.

Direct Detection


Spin Independent cross-section

Look for elastic scattering of WIMPS with nuclei.

$$\frac{d\sigma}{dE} = \frac{m_A}{2\mu_A^2 v^2} \cdot (\sigma_0^{\text{SI}} \cdot F_{\text{SI}}^2(E) + \sigma_0^{\text{SD}} \cdot F_{\text{SD}}^2(E))$$

$$\sigma_0^{\text{SI}} = \sigma_p \cdot \frac{\mu_A^2}{\mu_p^2} \cdot [Z \cdot f^p + (A - Z) \cdot f^n]^2$$

$$f_N/m_N = \sum_{q=u,d,s} f_{Tq}(f_q) + \sum_{q=u,d,s,c,b} \frac{3}{4} [q(2) + \bar{q}(2)] (g_q^{(1)} + g_q^{(2)}) - \frac{8\pi}{9\alpha_s} f_{TG}(f_G) + \frac{3}{4} G(2) (g_G^{(1)} + g_G^{(2)}) .$$




Source: KIPAC

$$\begin{aligned} \langle N | m_q \bar{q} q | N \rangle / m_N &\equiv f_{Tq} , \\ 1 - \sum_{u,d,s} f_{Tq} &\equiv f_{TG} , \\ \langle N(p) | \mathcal{O}_{\mu\nu}^q | N(p) \rangle &= \frac{1}{m_N} (p_\mu p_\nu - \frac{1}{4} m_N^2 g_{\mu\nu}) (q(2) + \bar{q}(2)) , \\ \langle N(p) | \mathcal{O}_{\mu\nu}^g | N(p) \rangle &= \frac{1}{m_N} (p_\mu p_\nu - \frac{1}{4} m_N^2 g_{\mu\nu}) G(2) . \end{aligned}$$

Nuclear matrix elements



$$0.94 f_G + 0.09 f_q + 0.29 (g_G^{(1)} + g_G^{(2)}) + 0.46 (g_q^{(1)} + g_q^{(2)})$$




Spin-dependent cross section

$$\sigma_N^{\text{SD}}(E_R) = 32\mu^2 G_F^2 [(J_N + 1)/J_N] [\langle S_p \rangle a_p + \langle S_n \rangle a_n]^2 F_{\text{SD}}^2(E_R)$$

- Due to axial vector coupling
- Depend on the nuclear spin J_N
- Dark matter couplings to proton and neutron a_p, a_n
- Nuclear form factor F_{SD}
- Expectation values $\langle S_{\{p,n\}} \rangle$ of the proton and neutron spin in the target nucleus.



Direct Detection Leading Order



$$\mathcal{L}_{int} = \sum_{q=u,d,s,c,b,t} g_{DM} (\tilde{q}_L^* \bar{\chi} P_L q + h.c.)$$


$$\mathcal{M}_{DD} \approx \frac{ig_{DM}^2}{M_{\tilde{q}_L}^2 - M_\chi^2} \frac{1}{8} [(\bar{\chi} \gamma^\mu \chi)(\bar{u} \gamma_\mu u) - (\bar{\chi} \gamma^\mu \gamma^5 \chi)(\bar{u} \gamma_\mu \gamma^5 u)]$$

SI **SD**




0 for Majorana

$$\sigma_p = \frac{4}{\pi} \left(\frac{M_\chi m_p}{M_\chi + m_p} \right)^2 |\langle \mathcal{M}_{DD} \rangle_{NR}|^2 .$$

- LO calculation tells us that model has only a spin dependent cross-section.
- Limits from direct detection are weak—large values of g_{DM} allowed.



DD @ 1-Loop



$$\mathcal{O}_{\mu\nu}^q \equiv \frac{1}{2} \bar{q} i \left(D_\mu \gamma_\nu + D_\nu \gamma_\mu - \frac{1}{2} g_{\mu\nu} \not{D} \right) q$$

$$\mathcal{O}_{\mu\nu}^g \equiv \left(G_\mu^{a\rho} G_{\rho\nu}^a + \frac{1}{4} g_{\mu\nu} G_{\alpha\beta}^a G^{a\alpha\beta} \right) .$$

Determine Wilson Coefficients for effective operators

Spin-2
Operators

$$\mathcal{L}_q^{\text{eff}} = f_q m_q \bar{\chi} \tilde{\chi} \bar{q} q + \frac{g_q^{(1)}}{m_\chi} \bar{\chi} i \partial^\mu \gamma^\nu \tilde{\chi} \mathcal{O}_{\mu\nu}^q + \frac{g_q^{(2)}}{m_\chi^2} \bar{\chi} (i \partial^\mu) (i \partial^\nu) \tilde{\chi} \mathcal{O}_{\mu\nu}^q ,$$

$$\mathcal{L}_g^{\text{eff}} = f_G \bar{\chi} \tilde{\chi} G_{\mu\nu}^a G^{a\mu\nu} + \frac{g_G^{(1)}}{m_\chi} \bar{\chi} i \partial^\mu \gamma^\nu \tilde{\chi} \mathcal{O}_{\mu\nu}^g + \frac{g_G^{(2)}}{m_\chi^2} \bar{\chi} (i \partial^\mu) (i \partial^\nu) \tilde{\chi} \mathcal{O}_{\mu\nu}^g .$$

Spin 0

Evaluate matrix element for the elastic scattering process in the non-relativistic limit.

$$f_N/m_N = \sum_{q=u,d,s} f_q f_{Tq} + \sum_{q=u,d,s,c,b} \frac{3}{4} (q(2) + \bar{q}(2)) (g_q^{(1)} + g_q^{(2)})$$


$$- \frac{8\pi}{9\alpha_s} f_{TG} f_G + \frac{3}{4} G(2) (g_G^{(1)} + g_G^{(2)}) .$$

Tools: FeynArts, FORM, PackageX

$$0.94 f_G + 0.09 f_q + 0.29 (g_G^{(1)} + g_G^{(2)}) + 0.46 (g_q^{(1)} + g_q^{(2)})$$

RGE

- Nucleon DM cross-sections at Non-Relativistic velocities.
- At what scale do we define coupling and masses? If at scale $\mu \sim 0$, then to compare with LHC we should run up. If at $\mu \sim \text{LHC energy}$, then to compare we should run down.
- RGE not necessary if no comparisons being made at different energy scales.



Details of RGE

Operators for Spin Independent Interactions

$O_q^{(0)} = m_q \bar{q}q$	$O_g^{(0)} = G_{\mu\nu}^A G^{A\mu\nu}$	Spin 0	Sum Rules Relate operators
$O_q^{(2)\mu\nu} = \frac{1}{2} \bar{q} \left(\gamma^{\{\mu} i D_-^{\nu\}} - \frac{g^{\mu\nu}}{4} i \not{D}_- \right) q$	$O_g^{(2)\mu\nu} = -G^{A\mu\lambda} G^{A\nu}_{\lambda} + \frac{g^{\mu\nu}}{4} (G_{\alpha\beta}^A)^2$	Spin 2	
Quark	Gluon		

Spin Dependent Operators

$$A_q^\mu = \bar{q} \gamma^\mu \gamma_5 q$$

Determine Anomalous dimensions

$$\frac{d}{d \log \mu} O_i = -\gamma_{ij} O_j, \quad \frac{d}{d \log \mu} c_i = \gamma_{ji} c_j$$

Evolve and Match at each threshold

$$c_i(\mu_l) = R_{ij}(\mu_l, \mu_h) c_j(\mu_h) .$$

$$c_i(\mu_Q) = M_{ij}(\mu_Q) c'_j(\mu_Q)$$

$$R = \left(\begin{array}{ccc|cc} & & & R_{qq} & \\ & & & \vdots & \\ & & & R_{qq} & \\ \hline R_{gq} & \dots & R_{gq} & R_{gg} & \end{array} \right), \quad \begin{aligned} R_{qq}^{(0)} &= 1, & R_{qg}^{(0)} &= 2[\gamma_m(\mu_h) - \gamma_m(\mu_l)]/\tilde{\beta}(\mu_h), \\ R_{gq}^{(0)} &= 0, & R_{gg}^{(0)} &= \tilde{\beta}(\mu_l)/\tilde{\beta}(\mu_h) \end{aligned}$$

$$R_{qq}^{(2)} - R_{qq'}^{(2)} = r(0) + \mathcal{O}(\alpha_s), \quad R_{qq'}^{(2)} = \frac{1}{n_f} \left[\frac{16r(n_f) + 3n_f}{16 + 3n_f} - r(0) \right] + \mathcal{O}(\alpha_s)$$

$$R_{qg}^{(2)} = \frac{16[1 - r(n_f)]}{16 + 3n_f} + \mathcal{O}(\alpha_s),$$

$$R_{gq}^{(2)} = \frac{3[1 - r(n_f)]}{16 + 3n_f} + \mathcal{O}(\alpha_s), \quad R_{gg}^{(2)} = \frac{16 + 3n_f r(n_f)}{16 + 3n_f} + \mathcal{O}(\alpha_s)$$

$$\langle O_q'^{(0)} \rangle = \langle O_q^{(0)} \rangle + \mathcal{O}(1/m_Q)$$

$$M = \left(\begin{array}{ccc|cc} 1 & & & 0 & 0 \\ & \ddots & & \vdots & \vdots \\ & & 1 & 0 & 0 \\ \hline 0 & \dots & 0 & M_{gQ} & M_{gg} \end{array} \right)$$


$$M_{gQ}^{(0)} = -\frac{\alpha'_s(\mu_Q)}{12\pi} \left\{ 1 + \frac{\alpha'_s(\mu_Q)}{4\pi} \left[11 - \frac{4}{3} \log \frac{\mu_Q}{m_Q} \right] + \mathcal{O}(\alpha_s^2) \right\}$$

$$M_{gg}^{(0)} = 1 - \frac{\alpha'_s(\mu_Q)}{3\pi} \log \frac{\mu_Q}{m_Q} + \mathcal{O}(\alpha_s^2)$$


Operators Mix

$$c_j(\mu_0) = R_{jk}(\mu_0, \mu_c) M_{kl}(\mu_c) R_{lm}(\mu_c, \mu_b) M_{mn}(\mu_b) R_{ni}(\mu_b, \mu_t) c_i(\mu_t)$$

How important is RGE?



Spin-2 Wilson coefficients do not run as strongly.



Factor ~2 increase in matrix element when performing RGE.

Factor ~4 enhancement in cross-section

Those large logs!



Port Huron logging mills

A closer look at the Wilson Coefficients



$$\frac{g_G^{(1)}}{m_\chi} = \alpha_s \alpha_{DM} \left[f_1(m_q, M_{\tilde{q}_L}, m_\chi) \log \left(\frac{m_q}{M_{\tilde{q}_L}} \right) + f_2(m_q, M_{\tilde{q}_L}, m_\chi) \right]$$

- For light quarks, large logs dominate the loop integral.
- Including RGE ensures large logs cancel



$$\Delta g_G^{(1)} = \frac{\alpha_s g_{DM}^2 m_\chi}{24\pi(M^2 - m_\chi^2)^2} \log \left(\frac{M}{m} \right)$$

$$\Delta g_G^{(1)} \Big|_{\mu_l} \simeq \frac{m_\chi g_{DM}^2}{72\pi^2(M_{\tilde{q}}^2 - m_\chi^2)^2} \left[3\pi\alpha_s(\mu_h) \log \left(\frac{\mu_l}{\mu_h} \right) + \alpha_s(M_{\tilde{q}}) \log \left(\frac{M_{\tilde{q}}}{m_b} \right) \left(3\pi - 5\alpha_s(\mu_h) \log \left(\frac{\mu_l}{\mu_h} \right) \right) \right]$$

SI Limits (Loop)





SD Limits (LO)




Constraints improve by an order of magnitude.

LHC Constraints







- Colored scalar mediator pair production— production cross-section (mostly QCD) depends on mass of mediator alone.
- Acceptance depends on mass of dark matter candidate also.





- Associated production of colored mediator and dark matter candidate— depends on all three model parameters.






K factors






Pure QCD
 Independent of
 DM coupling (g_{DM})




Tools : FeynRules, NLOCT, MadGraph5_aMC@NLO, MadAnalysis5

Complementarity of DD & LHC experiments



Dark matter annihilation cross-section



$$\langle\sigma v\rangle \simeq N_c^f g_{DM}^4 \left[\frac{m_f^2 \sqrt{1 - \frac{m_f^2}{m_\chi^2}}}{64\pi(m_{\tilde{q}}^2 + m_\chi^2 - m_f^2)^2} + \beta^2 \left\{ \frac{m_\chi^2 \sqrt{m_\chi^4 + m_{\tilde{q}}^4}}{32\pi(m_\chi^2 + m_{\tilde{q}}^2)^4} + \mathcal{O}(m_f^2) \right\} \right]$$

Conclusions

- Simplified models— useful to capture essential features of classes of models; one can make generic statements.
- Importantly - one is able to compare constraints from experiments probing a wide range of energy scales.
- Can easily evaluate LHC constraints @ NLO precision — Madgraph UFO file available.
- Demonstrated importance of going beyond LO for determining direct-detection constraints (order of magnitude increase for this particular model).
- Demonstrated importance of RGE effects (factor 4 increase).
- Able to make proper comparison between experiments that operate at different energy scales.
- Tool to calculate direct detection constraints with RGE will be hosted on the msu website soon. CalcHep model file to run with micromegas also available.

Thank you

$$\begin{aligned}
m_u &= 2.2 \text{ MeV}, & m_d &= 4.7 \text{ MeV}, & m_s &= 95 \text{ MeV}, \\
m_c &= 1.3 \text{ GeV}, & m_b &= 4.2 \text{ GeV}, & m_t &= 172 \text{ GeV}, \\
m_Z &= 91.188 \text{ GeV}, & \alpha_s(m_Z) &= 0.1184, \\
m_n &= 0.9396 \text{ GeV} & m_p &= 0.9383 \text{ GeV} .
\end{aligned}$$

$$\begin{aligned}
[f_{T_u}]_p &= 0.018, & [f_{T_d}]_p &= 0.030, & [f_{T_s}]_p &= 0.043, \\
[f_{T_u}]_n &= 0.015, & [f_{T_d}]_n &= 0.034, & [f_{T_s}]_n &= 0.043, \\
f_{T_G}|_{\text{NNNLO}} &= 0.80 .
\end{aligned}$$

$$\begin{aligned}
[u(2) + \bar{u}(2)]_p &= 0.3481, & [d(2) + \bar{d}(2)]_p &= 0.1902, \\
[s(2) + \bar{s}(2)]_p &= 0.0352, & [c(2) + \bar{c}(2)]_p &= 0.0107 , \\
[G(2)]_p &= [G(2)]_n = 0.4159 .
\end{aligned}$$

$$\begin{aligned}
\Delta u^{(p)} &= 0.84, & \Delta d^{(p)} &= -0.43, & \Delta s^{(p)} &= -0.09, \\
\Delta u^{(n)} &= \Delta d^{(p)}, & \Delta d^{(n)} &= \Delta u^{(p)}, & \Delta s^{(n)} &= \Delta s^{(p)} .
\end{aligned}$$

$$f_{T_G} = -\frac{9\alpha_s(\mu)}{4\pi\beta(\mu)} \left[1 - (1 + \gamma_m(\mu)) \sum_{u,d,s} f_{T_q} \right] .$$