



Contribution ID: 126

Type: not specified

## Probing low scale leptogenesis with inflationary blue-tilted GWs

*Monday 23 January 2023 12:20 (20 minutes)*

In Low scale leptogenesis (LSL) models there is a possibility of dynamically generating a small RH neutrino mass when a long-lived scalar field is coupled weakly to the RH neutrinos despite having a large VEV,  $v_\Phi$ . Considering such a scenario, the correlation shared by the non-standard scalar era driven by  $\Phi$  and  $M_{iS}$  offers a unique opportunity to study the fingerprints of LSL on primordial gravitational waves. We study the gravitational waves originating due to the inflationary blue-tilted tensor power spectrum and propagating through the scalar epoch, which depending upon  $M_{iS}$  provides two important insights. Firstly, if LSLs are taken seriously even for very high scale reheating GWs with a significant blue tilt don't violate the BBN bound. Secondly, It provides an opportunity to test LSLs with a low frequency and a complementary high frequency doubly peaked GW background. If recent results on GWs from PTAs are taken at face value and used at low frequencies as a measure, allows one to get possible signatures of LSL mechanisms at higher frequencies.

**Primary authors:** Mr DATTA, Satyabrata (Saha Institute of Nuclear Physics); Dr SAMANTA, Rome (Czech Academy of Sciences, CEICO)

**Presenter:** Mr DATTA, Satyabrata (Saha Institute of Nuclear Physics)