

Hadronic molecules of heavy hadrons with tensor force

Yasuhiro Yamaguchi (RIKEN, Japan)


in collaboration with

Hugo García-Tecocoatzi (UNAM), Alessandro Giachino (INFN Genoa, Genoa Univ.), Atsushi Hosaka (RCNP, Osaka Univ.), Elena Santopinto (INFN Genoa), Sachiko Takeuchi (Japan Coll. Social Work), Makoto Takizawa (Showa Pharmaceutical Univ.).

The 24th European conference on few-body problems in physics (EFB24)

University of Surrey, UK 1-6 September 2019


- 1 Introduction
 - Exotic hadrons - Hadronic molecules
 - Hidden-charm pentaquark P_c
- 2 Model setup
 - One Pion Exchange Potential
 - Compact 5-quark potential
- 3 Numerical results
 - Hidden-charm molecules
- 4 Summary



Hadron structure: Constituent quark model


Introduction

- Ordinary Hadrons: Baryon (qqq) and Meson ($q\bar{q}$)



* q : "Constituent quark"


- Exotic Hadrons ($\neq qqq, q\bar{q}$): **Multiquark? Multihadron?**



Constituent quark picture and beyond

Introduction

▷ e.g. $c\bar{c}$ mesons (Charmonium)



N. Brambilla, et al. Eur.Phys.J.C **71**(2011)1534,

S. Godfrey and N. Isgur, PRD**32**(1985)189

one-g exchange

+

Confinement


⇓

$$V(r) = -\frac{\alpha_s}{r} + cr + \dots$$

Constituent quark picture and beyond

Introduction

- ▷ e.g. $c\bar{c}$ mesons (Charmonium) and **Unexpected X, Y, Z**



N. Brambilla, et al. Eur.Phys.J.C **71**(2011)1534,

S. Godfrey and N. Isgur, PRD**32**(1985)189

one-g exchange

+

Confinement

⇓


$$V(r) = -\frac{\alpha_s}{r} + cr + \dots$$

- Exotics $\neq c\bar{c}$ have been observed in the Experiments (BaBar, Belle, BESIII, LHCb,...) \Rightarrow **Q. Structure? Physics?**

Hadronic molecules?

Introduction

- Exotics as Hadronic molecule = Hadron composite system




Hadronic molecule

$X(3872) = D\bar{D}^*$ molecule?

$$D^+ D^{*-} \quad \underline{\underline{3879.84 \text{ MeV}}}$$

$$D^0 \bar{D}^{*0} \quad \underline{\underline{3871.69 \text{ MeV}}}$$
$$\underline{\underline{3871.68 \text{ MeV}}}$$


- Expected **near the thresholds**
→ Hadron-hadron (quasi) bound state
- ⇒ Analogous to **Atomic Nuclei**
Deuteron $\sim pn$ bound state ($B = 2.2 \text{ MeV}$)



Hadronic molecules and π exchange potential

Introduction



- Driving force of Nuclei \Rightarrow long range force: π exchange
— \rightarrow generating **the loosely bound state**



Hadronic molecules and π exchange potential

Introduction

- Driving force of Nuclei \Rightarrow long range force: π exchange
 \rightarrow generating **the loosely bound state**





K.Ikeda *et al.*, Lect.Notes.Phys.**818**(2010)165

- Strong attraction from **Tensor term ($S - D$ mixing)**

Hadronic molecules and π exchange potential


Introduction

- Driving force of Nuclei \Rightarrow long range force: π exchange
 \rightarrow generating **the loosely bound state**




K.Ikeda *et al.*, Lect.Notes.Phys.**818**(2010)165

- Strong attraction from **Tensor term ($S - D$ mixing)**
 \Rightarrow Important role in the heavy hadronic molecules?



N.A.Tornqvist, Z.Phys.**C61**(1994)525

Hidden-charm pentaquarks




R.Aaij, *et al.* (LHCb collaboration) PRL**115**(2015)072001, PRL**122**(2019)222001

Two hidden-charm pentaquarks !! (2015)

Introduction: pentaquark

- Observation of the Hidden-charm Pentaquark ($c\bar{c}uud$) in $\Lambda_b^0 \rightarrow J/\psi K^- p$ Decay? R.Aaij, *et al.* (LHCb collaboration) PRL**115**(2015)072001




$$P_c(4380): \quad M = 4380 \text{ MeV} \quad \Gamma = 205 \text{ MeV} \quad P_c(4450): \quad M = 4449.8 \text{ MeV} \quad \Gamma = 39 \text{ MeV}$$

- J^P ? ($3/2^-, 5/2^+$), ($3/2^+, 5/2^-$), or ($5/2^+, 3/2^-$)
- There have been a lot of articles investigating the P_c states...
Hadronic molecule? Compact state? Kinematical effect?

New LHCb analysis in 2019!

Introduction: pentaquark

- R. Aaij, *et al.* Phys.Rev.Lett. 122 (2019) no.22, 222001



- $P_c(4450)$ in 2015 $\rightarrow P_c(4440)$ and $P_c(4457)$
 $P_c(4440)$ $M = 4440.3$ MeV $P_c(4457)$ $M = 4457.3$ MeV
 $\Gamma = 20.6$ $\Gamma = 6.4$ MeV
- Observation of **New state!** $P_c(4312)$ $M = 4311.9$ MeV
 $\Gamma = 9.8$ MeV
- $P_c(4380)$ in 2015? "these fits can neither confirm nor contradict the existence of the $P_c(4380)^{+u}$ "

Hidden-charm meson-baryon molecule...?

Introduction: pentaquark

- ▶ P_c states reported close to **the $\bar{D}^{(*)}\Sigma_c^{(*)}$ thresholds**
- ▶ **π exchange** in the heavy hadron system

Hidden-charm meson-baryon molecule...?

Introduction: pentaquark

- ▶ P_c states reported close to **the $\bar{D}^{(*)}\Sigma_c^{(*)}$ thresholds**
- ▶ **π exchange** in the heavy hadron system
enhanced by **the heavy quark spin symmetry!**

N.Isgur, M.B.Wise, PLB232(1989)113

$\Rightarrow \bar{D}(0^-) - \bar{D}^*(1^-), \Sigma_c(1/2^+) - \Sigma_c^*(3/2^-)$ **mixing**

Hidden-charm meson-baryon molecule...?


Introduction: pentaquark

- ▶ P_c states reported close to **the $\bar{D}^{(*)}\Sigma_c^{(*)}$ thresholds**
- ▶ **π exchange** in the heavy hadron system
enhanced by **the heavy quark spin symmetry!**

N.Isgur, M.B.Wise, PLB232(1989)113

$\Rightarrow \bar{D}(0^-) - \bar{D}^*(1^-), \Sigma_c(1/2^+) - \Sigma_c^*(3/2^-)$ **mixing**

$\Rightarrow P_c$ states: $\bar{D}^{(*)}\Sigma_c^{(*)}$ molecules?





Hadronic molecule

Compact state: 5-quark configuration

Introduction: pentaquark

- S. Takeuchi and M. Takizawa, PLB**764** (2017) 254-259.
 P_c states by the quark cluster model
- 5-quark configurations





$$S_{q^3} = 1/2, 3/2, S_{c\bar{c}} = 0, 1 \quad S_{q^3} = 1/2, S_{c\bar{c}} = 0, 1$$

Compact state: 5-quark configuration

Introduction: pentaquark

- S. Takeuchi and M. Takizawa, PLB**764** (2017) 254-259.
 P_c states by the quark cluster model
- 5-quark configurations





$$S_{q^3} = 1/2, 3/2, S_{c\bar{c}} = 0, 1 \quad S_{q^3} = 1/2, S_{c\bar{c}} = 0, 1$$

- $[q^3 8_c 3/2]$: Color magnetic int. is attractive!

Compact state: 5-quark configuration

Introduction: pentaquark

- S. Takeuchi and M. Takizawa, PLB**764** (2017) 254-259.
 P_c states by the quark cluster model
- 5-quark configurations



$$S_{q^3} = 1/2, 3/2, S_{c\bar{c}} = 0, 1 \quad S_{q^3} = 1/2, S_{c\bar{c}} = 0, 1$$


- $[q^3 8_c 3/2]$: Color magnetic int. is attractive!
 \Rightarrow Couplings to (qqc) baryon- $(q\bar{c})$ meson, e.g. $\bar{D}\Sigma_c$, are allowed!

Mixing of Compact state and Hadronic Molecule!

Model setup in this study

- Hadronic molecule (MB) + Compact state ($5q$)


$MB + 5q$



Model setup in this study

- **Hadronic molecule (MB) + Compact state ($5q$)**
⇒ MB coupled to $5q$ (Feshbach Projection)


$MB + 5q$



Model setup in this study

- **Hadronic molecule (MB) + Compact state ($5q$)**
⇒ MB coupled to $5q$ (Feshbach Projection)

Interaction of hadrons (M and B)



- ▶ **Long range** interaction: One pion exchange potential (OPEP)
- ▶ **Short range** interaction: $5q$ potential


Y.Y, A. Giachino, A. Hosaka, E. Santopinto, S. Takeuchi, M. Takizawa, PRD**96**(2017)114031

Y.Y, H.G-Tecocoatzi, A.Giachino, A.Hosaka, E.Santopinto, S.Takeuchi, M.Takizawa, 1907.04684[hep-ph]

Model setup in this study

- **Hadronic molecule (MB) + Compact state ($5q$)**
⇒ MB coupled to $5q$ (Feshbach Projection)

Interaction of hadrons (M and B)



- ▶ **Long range** interaction: One pion exchange potential (OPEP)
- ▶ **Short range** interaction: $5q$ potential (→ **Local Gaussian**)

Spin dependence → Spin structure of $5q$


Y.Y, A. Giachino, A. Hosaka, E. Santopinto, S. Takeuchi, M. Takizawa, PRD**96**(2017)114031

Y.Y, H.G-Tecocoatzi, A.Giachino, A.Hosaka, E.Santopinto, S.Takeuchi, M.Takizawa, 1907.04684[hep-ph]

$\bar{D}^{(*)} Y_c$ Interaction: Long range force

HQS and OPEP

- One pion exchange potential



$\bar{D}^{(*)}$: \bar{D} or \bar{D}^*

Y_c : Λ_c , Σ_c or Σ_c^*

$$V_{\bar{D}^{(*)} Y_c - \bar{D}^{(*)} Y_c}^{\pi} = \frac{g_{D^* D \pi} g_{Y_c Y_c \pi}}{f_{\pi}^2} \left[\vec{S}_1 \cdot \vec{S}_2 C(r) + S_{S_1 S_2} T(r) \right]$$


(Contact term is removed)

- Form factor with Cutoff Λ (determined by the hadron size)

$$F(q^2) = \frac{\Lambda^2 - m_{\pi}^2}{\Lambda^2 - q^2}, \quad \Lambda_{\bar{D}} \sim 1130 \text{ MeV}, \Lambda_{Y_c} \sim 840 \text{ MeV}$$


Y.Y, A. Giachino, A. Hosaka, E. Santopinto, S. Takeuchi, M. Takizawa, Phys.Rev. **D96** (2017), 114031

2. Short range force: 5-quark potential



Model: 5-quark potential

- 5-quark potential \Rightarrow **Local Gaussian potential** is employed.
- ▷ Massive M_{5q} (few hundred MeV above $\bar{D}^*\Sigma_c^*$) \rightarrow **Attractive**


$$\Rightarrow -f S_i S_j e^{-\alpha r^2}$$

Channel $i, j = \bar{D}^{(*)}\Lambda_c, \bar{D}^{(*)}\Sigma_c^{(*)}$ with S -wave

Model: 5-quark potential

- 5-quark potential \Rightarrow **Local Gaussian potential** is employed.
- ▷ Massive M_{5q} (few hundred MeV above $\bar{D}^*\Sigma_c^*$) \rightarrow **Attractive**

$$\Rightarrow -f S_i S_j e^{-\alpha r^2}$$

Channel $i, j = \bar{D}^{(*)}\Lambda_c, \bar{D}^{(*)}\Sigma_c^{(*)}$ with S -wave

Free Parameters

Strength f and Gaussian para. α (\rightarrow may be fixed in the future)
(f vs E will be shown latter. $\alpha = 1 \text{ fm}^{-2}$ is fixed.)

Model: 5-quark potential

- 5-quark potential \Rightarrow **Local Gaussian potential** is employed.
- ▷ Massive M_{5q} (few hundred MeV above $\bar{D}^*\Sigma_c^*$) \rightarrow **Attractive**

$$\Rightarrow -f S_i S_j e^{-\alpha r^2}$$

Channel $i, j = \bar{D}^{(*)}\Lambda_c, \bar{D}^{(*)}\Sigma_c^{(*)}$ with S -wave

Free Parameters

Strength f and Gaussian para. α (\rightarrow may be fixed in the future)
(f vs E will be shown latter. $\alpha = 1 \text{ fm}^{-2}$ is fixed.)

Relative strength S_i

Spectroscopic factors \Rightarrow determined by **the spin structure** of $5q$

Model: 5-quark potential

- 5-quark potential \Rightarrow **Local Gaussian potential** is employed.
- ▷ Massive M_{5q} (few hundred MeV above $\bar{D}^*\Sigma_c^*$) \rightarrow **Attractive**

$$\Rightarrow -f S_i S_j e^{-\alpha r^2}$$

Channel $i, j = \bar{D}^{(*)}\Lambda_c, \bar{D}^{(*)}\Sigma_c^{(*)}$ with S -wave

Free Parameters

Strength f and Gaussian para. α (\rightarrow may be fixed in the future)
(f vs E will be shown latter. $\alpha = 1 \text{ fm}^{-2}$ is fixed.)

Relative strength S_i

Spectroscopic factors \Rightarrow determined by **the spin structure** of $5q$

Spectroscopic factor S_i (Spin structure)

$5q$ potential

- S-factor: $S_i = \langle (\bar{D}Y_c)_i | 5q \rangle$

Table: Spectroscopic factors S_i for each meson-baryon channel.

J		$S_{c\bar{c}}$	S_{3q}	$\bar{D}\Lambda_c$	$\bar{D}^*\Lambda_c$	$\bar{D}\Sigma_c$	$\bar{D}\Sigma_c^*$	$\bar{D}^*\Sigma_c$	$\bar{D}^*\Sigma_c^*$
1/2	(i)	0	1/2	0.4	0.6	-0.4	—	0.2	-0.6
	(ii)	1	1/2	0.6	-0.4	0.2	—	-0.6	-0.3
	(iii)	1	3/2	0.0	0.0	-0.8	—	-0.5	0.3
3/2	(i)	0	3/2	—	0.0	—	-0.5	0.6	-0.7
	(ii)	1	1/2	—	0.7	—	0.4	-0.2	-0.5
	(iii)	1	3/2	—	0.0	—	-0.7	-0.8	-0.2
5/2	(i)	1	3/2	—	—	—	—	—	-1.0

Spectroscopic factor S_i (Spin structure)

$5q$ potential


- S-factor: $S_i = \langle (\bar{D}Y_c)_i | 5q \rangle$

Table: Spectroscopic factors S_i for each meson-baryon channel.

J		$S_{c\bar{c}}$	S_{3q}	$\bar{D}\Lambda_c$	$\bar{D}^*\Lambda_c$	$\bar{D}\Sigma_c$	$\bar{D}\Sigma_c^*$	$\bar{D}^*\Sigma_c$	$\bar{D}^*\Sigma_c^*$
1/2	(i)	0	1/2	0.4	0.6	-0.4	—	0.2	-0.6
	(ii)	1	1/2	0.6	-0.4	0.2	—	-0.6	-0.3
	(iii)	1	3/2	0.0	0.0	-0.8	—	-0.5	0.3
3/2	(i)	0	3/2	—	0.0	—	-0.5	0.6	-0.7
	(ii)	1	1/2	—	0.7	—	0.4	-0.2	-0.5
	(iii)	1	3/2	—	0.0	—	-0.7	-0.8	-0.2
5/2	(i)	1	3/2	—	—	—	—	—	-1.0

- $\bar{D}Y_c$ with **Large S_i** will play an important role.

Numerical Results for Hidden-charm sector




Bound state and Resonance

- Coupled-channel Schrödinger equation for $\bar{D}\Lambda_c$, $\bar{D}^*\Lambda_c$, $\bar{D}\Sigma_c$, $\bar{D}\Sigma_c^*$, $\bar{D}^*\Sigma_c$, $\bar{D}^*\Sigma_c^*$ (6 MB components).
- For $J^P = 1/2^-, 3/2^-, 5/2^-$ (Negative parity)

Results (f^{5q} vs E) of charm $\bar{D}Y_c$ for $J^P = 1/2^-$

- Energy with $V_\pi + V^{5q}(f^{5q})$. (Y.Yamaguchi *et al*, PRD**96** (2017), 114031)




Dashed line: Thresholds, **Red line: Energy obtained**

- For small f^{5q} , **No bound state**
 \Rightarrow The OPEP attraction is not enough to generate a state
- $5q$ potential helps to generate the states **near the thresholds**
 \Leftrightarrow **Large S-factor** (Spin structure)


Results (f^{5q} vs E) for $J^P = 3/2^-, 5/2^-$

- Energy with $V_\pi + V^{5q}(f^{5q})$. (Y.Yamaguchi *et al*, PRD**96** (2017), 114031)

$$J^P = 3/2^-$$



$$J^P = 5/2^-$$



- For small f^{5q} , **No bound state**
 \Rightarrow The OPEP attraction is not enough to generate a state
- $5q$ potential helps to generate the states **near the thresholds**
 \Leftrightarrow **Large S-factor** (Spin structure)

In 2019, New P_c states by LHCb


- $f^{5q} = 45$ is fixed to reproduce new P_c 's

Y.Y., H.Garcia-Tecocoatzi, A.Giachino, A.Hosaka, E.Santopinto, S.Takeuchi, M.Takizawa, 1907.04684[hep-ph]

In 2019, New P_c states by LHCb

- $f^{5q} = 45$ is fixed to reproduce new P_c 's

Y.Y., H.Garcia-Tecocoatz, A.Giachino, A.Hosaka, E.Santopinto, S.Takeuchi, M.Takizawa, 1907.04684[hep-ph]




- (E, Γ) in our pred.
 \leftrightarrow consistent with Exp for
 $P_c(4312)$, $P_c(4440)$, $P_c(4457)$

- Missing three ($1/2^-$, $3/2^-$, $5/2^-$) states below $\bar{D}^* \Sigma_c^*$
- Broad $P_c(4380) \leftrightarrow$ Our prediction with 4376 MeV
 \rightarrow Further theoretical and experimental studies are necessary...

In 2019, New P_c states by LHCb

- $f^{5q} = 45$ is fixed to reproduce new P_c 's


Y.Y., H.Garcia-Tecocoatz, A.Giachino, A.Hosaka, E.Santopinto, S.Takeuchi, M.Takizawa, 1907.04684[hep-ph]



- (E, Γ) in our pred.
 \leftrightarrow consistent with Exp for
 $P_c(4312)$, $P_c(4440)$, $P_c(4457)$

- Missing three ($1/2^-$, $3/2^-$, $5/2^-$) states below $\bar{D}^* \Sigma_c^*$
- Broad $P_c(4380) \leftrightarrow$ Our prediction with 4376 MeV
 \rightarrow Further theoretical and experimental studies are necessary...

J^P assignment for $P_c(4440)$ and $P_c(4457)$



▷ J^P assignment

$$P_c(4440) : 3/2^-$$

$$P_c(4457) : 1/2^-$$

⇒ $E(1/2^-) > E(3/2^-)$

- OPEP of the $\bar{D}^* \Sigma_c$ channel

$$1/2^- : {}^2S, \quad {}^4D$$


$$\begin{pmatrix} 4C & 2\sqrt{2}T \\ 2\sqrt{2}T & -2C + 4T \end{pmatrix}$$

$$3/2^- : {}^4S, \quad {}^2D, \quad {}^4D$$

$$\begin{pmatrix} -2C & -2T & -4T \\ -2T & 4C & 2T \\ -4T & 2T & -2C \end{pmatrix}$$

* C : Central force, T : Tensor force

J^P assignment for $P_c(4440)$ and $P_c(4457)$



▷ J^P assignment

$$P_c(4440) : 3/2^-$$

$$P_c(4457) : 1/2^-$$

$$\Rightarrow E(1/2^-) > E(3/2^-)$$

- OPEP of the $\bar{D}^* \Sigma_c$ channel

$$1/2^- : {}^2S, {}^4D$$

$$\begin{pmatrix} 4C & 2\sqrt{2}T \\ 2\sqrt{2}T & -2C + 4T \end{pmatrix}$$

$$3/2^- : {}^4S, {}^2D, {}^4D$$

$$\begin{pmatrix} -2C & -2T & -4T \\ -2T & 4C & 2T \\ -4T & 2T & -2C \end{pmatrix}$$

* C : Central force, T : Tensor force

- $S - D, D - D$ couplings producing the attraction from the tensor force

$$\Rightarrow 1/2^- : {}^2S - {}^4D$$

$$3/2^- : {}^4S - {}^2D, {}^4S - {}^4D, {}^2D - {}^4D \quad \updownarrow \text{14 MeV}$$

More attractive!

Summary

- Hidden-charm pentaquarks as Hadronic molecule + Compact multiquark
- $\bar{D}^{(*)} Y_c$ Interaction
 - Long range force: OPEP with the tensor force, enhanced by the heavy quark symmetry
 - Short range force: Coupling to Compact $5q$ states
- The OPEP is not enough to generate the bound state. \rightarrow OPEP + $5q$ potential generates the states
- Applying this model to New hidden-charm pentaquarks by LHCb in 2019
 - \Rightarrow our prediction is consistent with EXP
- The J^P assignment $P_c(4440)$: $3/2^-$ and $P_c(4457)$: $1/2^-$ understood by the tensor force of the OPEP

Y.Y., A. Giachino, A. Hosaka, E. Santopinto, S. Takeuchi, M. Takizawa, Phys.Rev. **D96** (2017), 114031

Y.Y., H.Garcia-Tecocoatz, A.Giachino, A.Hosaka, E.Santopinto, S.Takeuchi, M.Takizawa, 1907.04684[hep-ph]

Back up

Experimental mass and Our prediction

State	Mass	Width	Our pred. (M, Γ, J^P)
$P_c(4312)^+$	$4311.9 \pm 0.7^{+6.8}_{-0.6}$	$9.8 \pm 2.7^{+3.7}_{-4.5}$	$(4312, 5, \frac{1}{2}^-)$
$P_c(4380)^+$	$4380 \pm 8 \pm 29$	$205 \pm 18 \pm 86$	$(4376, 8, \frac{3}{2}^-)$
$P_c(4440)^+$	$4440.3 \pm 1.3^{+4.1}_{-4.7}$	$20.6 \pm 4.9^{+8.7}_{-10.1}$	$(4442, 26, \frac{3}{2}^-)$
$P_c(4457)^+$	$4457.3 \pm 0.6^{+4.1}_{-1.7}$	$6.4 \pm 2.0^{+5.7}_{-1.9}$	$(4462, 6.6, \frac{1}{2}^-)$
			$(4524, 1.5, \frac{1}{2}^-)$
			$(4521, 23, \frac{3}{2}^-)$
			$(4511, 55, \frac{5}{2}^-)$

(in units of MeV)

Coupled-channels

Channels	$\bar{D}Y_c(2S+1L)$	
$1/2^-$	$\bar{D}\Lambda_c(2S), \bar{D}^*\Lambda_c(2S),$ $\bar{D}\Sigma_c(2S), \bar{D}\Sigma_c^*(4D),$ $\bar{D}^*\Sigma_c(2S, 4D), \bar{D}^*\Sigma_c^*(2S, 4D, 6D)$	(10 ch)
$3/2^-$	$\bar{D}\Lambda_c(2D), \bar{D}^*\Lambda_c(4S, 2D, 4D),$ $\bar{D}\Sigma_c(2D), \bar{D}\Sigma_c^*(4S, 4D),$ $\bar{D}^*\Sigma_c(4S, 2D, 4D), \bar{D}^*\Sigma_c^*(4S, 2D, 4D, 6D, 6G)$	(15 ch)
$5/2^-$	$\bar{D}\Lambda_c(2D), \bar{D}^*\Lambda_c(2D, 4D, 4G),$ $\bar{D}\Sigma_c(2D), \bar{D}\Sigma_c^*(4D, 4G),$ $\bar{D}^*\Sigma_c(2D, 4D, 4G), \bar{D}^*\Sigma_c^*(6S, 2D, 4D, 6D, 4G, 6G)$	(16 ch)

- 6 $\bar{D}Y_c$ channels: $\bar{D}\Lambda_c, \bar{D}^*\Lambda_c, \bar{D}\Sigma_c, \bar{D}\Sigma_c^*, \bar{D}^*\Sigma_c, \bar{D}^*\Sigma_c^*$.
- $S - D$ mixing induced by the Tensor force (S_{12})


Heavy hadron- π coupling

HQS and OPEP

- Effective Lagrangians: Heavy hadron and π

R. Casalbuoni *et al.*, Phys.Rept.**281** (1997)145, T. M. Yan, *et al.*, PRD**46**(1992)1148

Y.-R.Liu and M.Oka, PRD**85**(2012)014015



- ▷ Heavy meson: $\bar{D}^{(*)} \bar{D}^{(*)} \pi$ ($DD\pi$: Parity violation)

$$\mathcal{L}_{\pi HH} = -\frac{g_{\pi}}{2f_{\pi}} \text{Tr} [H \gamma_{\mu} \gamma_5 \partial^{\mu} \hat{\pi} \bar{H}], \quad H = \frac{1+\not{\epsilon}}{2} [D_{\mu}^{*} \gamma^{\mu} - D \gamma_5]$$


Heavy hadron- π coupling

HQS and OPEP

- Effective Lagrangians: Heavy hadron and π

R. Casalbuoni *et al.*, Phys.Rept.**281** (1997)145, T. M. Yan, *et al.*, PRD**46**(1992)1148

Y.-R.Liu and M.Oka, PRD**85**(2012)014015



- ▷ Heavy meson: $\bar{D}^{(*)}\bar{D}^{(*)}\pi$ ($DD\pi$: Parity violation)

$$\mathcal{L}_{\pi HH} = -\frac{g_\pi}{2f_\pi} \text{Tr} [H\gamma_\mu\gamma_5\partial^\mu\hat{\pi}\bar{H}], \quad H = \frac{1+\not{v}}{2} [D_\mu^*\gamma^\mu - D\gamma_5]$$


Heavy hadron- π coupling

HQS and OPEP

- Effective Lagrangians: Heavy hadron and π

R. Casalbuoni *et al.*, Phys.Rept.**281** (1997)145, T. M. Yan, *et al.*, PRD**46**(1992)1148

Y.-R.Liu and M.Oka, PRD**85**(2012)014015



- Heavy meson: $\bar{D}^{(*)}\bar{D}^{(*)}\pi$ (**$DD\pi$: Parity violation**)

$$\mathcal{L}_{\pi HH} = -\frac{g_\pi}{2f_\pi} \text{Tr} [H\gamma_\mu\gamma_5\partial^\mu\hat{\pi}\bar{H}], \quad \mathbf{H} = \frac{1+\not{v}}{2} [\mathbf{D}_\mu^*\gamma^\mu - \mathbf{D}\gamma_5]$$

- Heavy baryon: $\Sigma_c^{(*)}\Sigma_c^{(*)}\pi, \Lambda_c\Sigma_c^{(*)}\pi$ ($\Lambda_c\Lambda_c\pi$: Isospin breaking)


$$\mathcal{L}_{\pi BB} = -\frac{3}{4f_\pi} \mathbf{g}_1 (i v_\kappa) \varepsilon^{\mu\nu\lambda\kappa} \text{tr} [\bar{S}_\mu \partial_\nu \hat{\pi} S_\lambda] - \frac{g_4}{2f_\pi} \text{tr} [\bar{S}^\mu \partial_\mu \hat{\pi} \Lambda_c] + \text{H.c.},$$

$$\mathbf{S}_\mu = \Sigma_{c\mu}^* - \frac{1}{\sqrt{3}} (\gamma_\mu + \mathbf{v}_\mu) \gamma_5 \Sigma_c, \quad g_\pi = 0.59, g_1 = 1.00, g_4 = 1.06$$

Spectroscopic factors S_i (Spin structure)

$5q$ potential

- Spin of $5q$ states $\rightarrow S_{c\bar{c}}$ and S_{3q} configuration
e.g. for $J^P = 1/2^-$, (i), (ii), (iii)




$$J^P = 1/2^-$$

	$S_{c\bar{c}}$	S_{3q}
type (i)	0	1/2
(ii)	1	1/2
(iii)	1	3/2

Spectroscopic factors S_i (Spin structure)

$5q$ potential

- Spin of $5q$ states $\rightarrow S_{c\bar{c}}$ and S_{3q} configuration
e.g. for $J^P = 1/2^-$, (i), (ii), (iii)



$$J^P = 1/2^-$$

	$S_{c\bar{c}}$	S_{3q}
type (i)	0	1/2
(ii)	1	1/2
(iii)	1	3/2

- Overlap** of the spin wavefunctions of 5-quark state and $\bar{D}Y_c$

$$S_i = \langle (\bar{D}Y_c)_i | 5q \rangle$$

\Rightarrow Relative strength of couplings to $\bar{D}Y_c$ channel

Volume integrals of the potentials

- Bound and Resonant states appears for $f^{5q} \gtrsim 25$
⇔ Large? Small?

Volume integrals of the potentials

- Bound and Resonant states appears for $f^{5q} \gtrsim 25$
 \Leftrightarrow Large? Small?

▶ Volume integral $V(q=0) = \int V(r) dr^3$

Comparison with the NN interaction (Bonn potential)

R. Machleidt, K. Holinde and C. Elster, Phys. Rept. **149**, 1 (1987).

$$\left| V_{f=25}^{5q}(0) \right| = 1.1 \times 10^{-4} \text{ MeV} \sim 0.03 |C_{NN}^{\sigma}(0)|$$

(C_{NN}^{σ} : Central force of σ exchange)

- $\left| V_{f=25}^{5q}(0) \right|$ is **much smaller** than $|C_{NN}^{\sigma}(0)|$.

However, the bound and resonant states are obtained!