

Quark Matter 2019 - the XXVIIIth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions



Contribution ID: 650

Type: **Poster Presentation**

Beam-energy dependence of spatial and temporal characteristics of shape-selected events in Au+Au collisions at STAR

Monday 4 November 2019 17:40 (20 minutes)

Femtoscopic measurements can be leveraged to gain insight into the expansion dynamics of the hot and dense medium created in heavy-ion collisions. This poster presentation will report and discuss excitation functions for shape-selected two-pion HBT radii (R_{out} , R_{side} and R_{long}) measured for a broad range of collision centrality and average pair transverse momentum (k_T) with the STAR detector. The shape selections were accomplished via cuts on the distributions of the second-order Q_2 vector [1]. The excitation functions, which span the full range of the RHIC beam energy scan ($\sqrt{s_{NN}} = 7.7-200$ GeV), indicate clear sensitivities to the magnitude of the Q_2 vector which give insight into the expansion dynamics. The connection between the magnitude of the Q_2 vector and the spatiotemporal characteristics of the quark-gluon plasma produced in the collisions will be discussed.

[1] J. Schukraft, A. Timmins, and S. A. Voloshin, Phys. Lett. B 719 (2013) 394.

Primary author: SCHWEID, Benjamin (Stony Brook University)

Presenter: SCHWEID, Benjamin (Stony Brook University)

Session Classification: Poster Session

Track Classification: Collective dynamics and final state interaction