

Quark Matter 2019 - the XXVIIIth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions



Contribution ID: 713

Type: **Poster Presentation**

Time evolution of chirality imbalance of the massive fermions under parallel electric and magnetic fields

Monday 4 November 2019 17:40 (20 minutes)

In the past several years there has been increasing interest in the chirality imbalance n_5 , which is the difference between right- and left-handed fermions. The chirality imbalance is expected to be produced from the Adler-Bell-Jackiw anomaly (ABJ-anomaly) and plays a key role to understand anomalous transport phenomena in the hot/dense quark matter or the Dirac/Weyl semimetals under the magnetic field. One of interesting transport phenomena in the presence of chirality imbalance is the chiral magnetic effect (CME), appearance of electric current in direction of the external magnetic field.

The ABJ-anomaly relation implies that the electric field gives a crucial contribution to the emergence of the chirality imbalance in addition to the magnetic field. In the previous work, we have studied the chirality imbalance and the CME using the solution of the Dirac equation in the constant magnetic and general time-dependent electric fields, and found that the time-dependence of the gauge field is essentially important for the production of n_5 . In particular, we found that n_5 and CME at $t \rightarrow \infty$ are solely determined by the probability distributions of the fermion created non-perturbatively by the electric field.

In this talk, we study the emergence of the chirality imbalance and the electric current under spatially-uniform and parallel electromagnetic field in the vacuum of massive fermion. For the time-dependence, we assume the constant magnetic field, but do not impose any specific form for the electric field with boundary conditions $E(t \rightarrow \pm\infty) \rightarrow 0$. We solve the Dirac equation and calculate vacuum expectation values of the currents with the gauge invariant regularization.

As a result, asymptotic forms of n_5 and CME consists of a constant part and an oscillating part independent of details of intermediate time-dependence of the gauge potential. The non-zero constant term is proportional to a relativistic velocity and the momentum distribution of the created particle, and understood as a classical analogue of the electric current. We discuss how the chirality imbalance arises by the time-dependent electric field.

Authors: Mr AOI, Hayato (Department of Physics, Faculty of Science, Tokyo University of Science); Prof. SUZUKI, Katsuhiko (Department of Physics, Faculty of Science, Tokyo University of Science)

Presenter: Mr AOI, Hayato (Department of Physics, Faculty of Science, Tokyo University of Science)

Session Classification: Poster Session

Track Classification: Chirality, vorticity and spin polarization