Quark Matter 2019 - the XXVIIIth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions



Contribution ID: 147

Type: Poster Presentation

Bottomonium production in p+p and Pb+Pb collisions with ATLAS

Monday 4 November 2019 17:40 (20 minutes)

Bottomonium, a bound state of a bottom quark and its antiquark, is an excellent probe of the hot and dense medium created in heavy-ion collisions at LHC. The ATLAS collaboration collected the large datasets of pp and Pb+Pb collisions in 2017 and 2018 corresponding to integrated luminosities of 242 pb⁻¹ and 1.39 nb⁻¹ respectively, at a centre-of-mass energy per nucleon pair of 5.02 TeV. In this poster, bottomonium states are reconstructed via the dimuon decay channel in the rapidity range of |y| < 1, and their production in PbPb collisions is compared to that in pp collisions to extract the nuclear modification factor, R_{AA} , as a function of event centrality and transverse momentum. In addition, the relative suppression of the excited states Y(nS) to the ground state Y(1S) is studied.

Author:LEE, Songkyo (Iowa State University (US))Presenter:LEE, Songkyo (Iowa State University (US))Session Classification:Poster Session

Track Classification: Heavy flavor and quarkonium