Quark Matter 2019 - the XXVIIIth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions



Contribution ID: 759

Type: Poster Presentation

Suppression of elliptic flow without viscosity

Monday 4 November 2019 17:40 (20 minutes)

We investigate fluid-to-particle conversion using the usual Cooper-Frye approach but with more general local equilibrium distributions than the Boltzmann or Bose/Fermi distributions typically used. Even though we study ideal fluids (i.e., shear stress and bulk pressure are zero everywhere), we find a suppression of elliptic flow (v_2) at high transverse momenta $(p_T > 1.5 \text{ GeV/c})$, relative to results obtained with the traditional Boltzmann distributions [1]. The non-viscous suppression shows qualitatively similar features to the well-known shear viscous suppression of v_2 ; for example, it increases with p_T , and it is smaller for heavier species as seen in self-consistent kinetic theory calculations. Our results question whether all of the v_2 suppression seen in the data can be attributed to viscous effects, and indicate that shear viscosities extracted from RHIC and LHC elliptic flow data might be overestimated. For more details, see Ref. [1].

[1] A. Takacs and D. Molnar, arXiv:1906.12311 [nucl-th]

Authors: TAKÁCS, Ádám (Wigner Research Centre for Physics); MOLNAR, Denes (Purdue University)

Presenter: MOLNAR, Denes (Purdue University)

Session Classification: Poster Session

Track Classification: Collective dynamics and final state interaction