

UNIVERSITY OF BERGEN



PARTON PROPAGATION AND ENERGY LOSS: NEW THEORETICAL PROGRESS

Konrad Tywoniuk

Many thanks to: Y. Mehtar-Tani, D. Pablos, A. Takacs, Alba Soto-Ontoso,...

*28th International Conference on Ultrarelativistic Nucleus-Nucleus Collisions (Quark Matter 2019)
3-9 November, 2019, Wuhan, China*


OUTLINE

- the structure of QCD jets
- the evolving story of an observable
- inevitability of jet substructure
- R-dependent jet spectrum in PbPb collisions



JET PHYSICS IN VACUUM

VACUUM SPLITTING




Generic $1 \rightarrow 2$ (on-shell) splitting in QCD:

$$d\Pi_{a \rightarrow bc} = \frac{\alpha_s}{\pi} \frac{d\theta}{\theta} P_{ba}^{(c)}(z) dz \approx \frac{2\alpha_s C_R}{\pi} \frac{d\theta}{\theta} \frac{dz}{z}$$

Diverges for soft & collinear radiation!



VACUUM SPLITTING




Generic $1 \rightarrow 2$ (on-shell) splitting in QCD:

$$d\Pi_{a \rightarrow bc} = \frac{\alpha_s}{\pi} \frac{d\theta}{\theta} P_{ba}^{(c)}(z) dz \approx \frac{2\alpha_s C_R}{\pi} \frac{d\theta}{\theta} \frac{dz}{z}$$

Diverges for soft & collinear radiation!

Large phase space for radiation compensates α_s !




$$\text{Prob} = \frac{\alpha_s C_R}{\pi} \log^2 \frac{p_T R}{\Lambda_{\text{QCD}}} \gg 1$$

Need for resummation of collinear logarithms for final-state radiation.



VACUUM SPLITTING



Observed as "jets": collimated sprays of particles/energy.


Sterman, Weinberg (1977)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent “leaves”
- accessed in experimental data using jet reconstruction (C/A algorithm)



SPACE-TIME PICTURE OF A JET

Andersson, Gustafson, Lönnblad, Pettersson Z.Phys.C (1989)

Andersson, Gustafson, Samuelsson NPB (1996)




hadronization


from $t_f \sim (Q_0 R)^{-1} \sim 2$ fm
to $t_f \sim E/Q_0^2 \sim 300$ fm

- semi-classical picture, angular ordering
- Lund diagram (primary emission plane)
 - secondary branchings located on independent "leaves"
- accessed in experimental data using jet reconstruction (C/A algorithm)







mass/virtuality



transverse momentum



angle



formation time

Freedom to choose ordering variable at leading logarithmic approximation.



Color coherence dictates angular ordering.



LOGARITHMIC RESUMMATION

Konishi, Ukawa, Veneziano Nucl. Phys. B1567 (1979);
 Bassetto, Ciafaloni, Marchesini Phys. Rept. 100 (1983);
 Dokshitzer, Khoze, Mueller, Troyan "Basics of Perturbative QCD" (1991)

Multi-gluon emissions: resummation of leading logarithmic contributions and Monte Carlo development.



Example: small-R resummation in $t = \log 1/R$:

$$\frac{\partial}{\partial t} f_{jet/i}(z, t) = \int_0^1 \frac{dz'}{z'} \frac{\alpha_s}{\pi} P_{ji}(z) f_{jet/j}\left(\frac{z}{z'}, t\right)$$


Evolution from hard scale to jet scale via DGLAP.

Dasgupta, Dreyer, Salam, Soyez 1411.5182, 1602.01110
 Kang, Ringer, Vitev 1606.06732



JET PHYSICS IN MEDIUM


ENERGY-LOSS BASICS



Workhorse of the field: measuring & parameterizing the shift of spectrum to access information about medium interactions.





ENERGY-LOSS BASICS



Workhorse of the field: measuring & parameterizing the shift of spectrum to access information about medium interactions.





ENERGY-LOSS BASICS



Workhorse of the field: measuring & parameterizing the shift of spectrum to access information about medium interactions.



ENERGY-LOSS BASICS





Workhorse of the field: measuring & parameterizing the shift of spectrum to access information about medium interactions.

Significant difference of hadron and jet suppression at high- p_T !



ENERGY-LOSS BASICS



- disclaimer: in my talk, I will focus on effects above the thermal scale - however, close connection to bottom-up thermalization & hydrodynamization should be further investigated.

Tachibana (Tue 10:00), Chen (Tue 10:20); Kumar (Wed 09:40), He (Wed 10:00); Pablos (Wed 12:20)




MEDIUM-INDUCED RADIATION

Baier, Dokshitzer, Mueller, Peigné, Schiff (1996); Zakharov (1996) (Arnold, Moore, Yaffe (2003))

$$z \frac{dI_{ba}}{dz} = \frac{\alpha_s z P_{ba}(z)}{(z(1-z)E)^2} 2\text{Re} \int_0^\infty dt_2 \int_0^{t_2} dt_1$$


$$\partial_{\mathbf{x}} \cdot \partial_{\mathbf{y}} \left[\mathcal{K}_{ba}(\mathbf{x}, t_2; \mathbf{y}, t_1) - \mathcal{K}_0(\mathbf{x}, t_2; \mathbf{y}, t_1) \right]_{\mathbf{x}=\mathbf{y}=0}$$



MEDIUM-INDUCED RADIATION

Baier, Dokshitzer, Mueller, Peigné, Schiff (1996); Zakharov (1996) (Arnold, Moore, Yaffe (2003))

$$z \frac{dI_{ba}}{dz} = \frac{\alpha_s z P_{ba}(z)}{(z(1-z)E)^2} 2\text{Re} \int_0^\infty dt_2 \int_0^{t_2} dt_1 \partial_{\mathbf{x}} \cdot \partial_{\mathbf{y}} \left[\mathcal{K}_{ba}(\mathbf{x}, t_2; \mathbf{y}, t_1) - \mathcal{K}_0(\mathbf{x}, t_2; \mathbf{y}, t_1) \right]_{\mathbf{x}=\mathbf{y}=0}$$



3-body interactions of the medium via Schrödinger equation with potential:

$$v(\mathbf{x}, t) = \int_{\mathbf{q}} \frac{d\sigma_{e1}}{d^2\mathbf{q}} (1 - e^{i\mathbf{q}\cdot\mathbf{x}}) \approx \underbrace{\frac{1}{4} \hat{q} x^2}_{\text{HO}} \log \frac{4}{\mu^2 x^2}$$


Solved numerically in Caron-Huot, Gale 1006.2379, Feal, Vazquez 1811.01591, Ke, Xu and Bass 1810.08177, Feal, Salgado, Vazquez 1911.01309



MEDIUM-INDUCED RADIATION

Baier, Dokshitzer, Mueller, Peigné, Schiff (1996); Zakharov (1996) (Arnold, Moore, Yaffe (2003))

$$z \frac{dI_{ba}}{dz} = \frac{\alpha_s z P_{ba}(z)}{(z(1-z)E)^2} 2\text{Re} \int_0^\infty dt_2 \int_0^{t_2} dt_1 \partial_{\mathbf{x}} \cdot \partial_{\mathbf{y}} \left[\mathcal{K}_{ba}(\mathbf{x}, t_2; \mathbf{y}, t_1) - \mathcal{K}_0(\mathbf{x}, t_2; \mathbf{y}, t_1) \right]_{\mathbf{x}=\mathbf{y}=0}$$



3-body interactions of the medium via Schrödinger equation with potential:

$$v(\mathbf{x}, t) = \int_{\mathbf{q}} \frac{d\sigma_{e1}}{d^2\mathbf{q}} (1 - e^{i\mathbf{q}\cdot\mathbf{x}}) \approx \underbrace{\frac{1}{4} \hat{q} x^2}_{\text{HO}} \log \frac{4}{\mu^2 x^2}$$

Solved numerically in Caron-Huot, Gale 1006.2379, Feal, Vazquez 1811.01591, Ke, Xu and Bass 1810.08177, Feal, Salgado, Vazquez 1911.01309

$$\mathcal{K}(t_2, t_1) = \mathcal{K}_0(t_2 - t_1) + \int dt \mathcal{K}_0(t_2 - t) v(t) \mathcal{K}(t, t_1)$$

opacity expansion


Wiedemann (2000); Gyulassy, Levai, Vitev (2001); Sievert, Vitev, Yoon 1903.06170
Wang, Gu (2001), Majumder 0912.2987
Mehtar-Tani 1903.00506, Mehtar-Tani KT 1910.02032



MEDIUM-INDUCED RADIATION

Baier, Dokshitzer, Mueller, Peigné, Schiff (1996); Zakharov (1996) (Arnold, Moore, Yaffe (2003))

$$z \frac{dI_{ba}}{dz} = \frac{\alpha_s z P_{ba}(z)}{(z(1-z)E)^2} 2\text{Re} \int_0^\infty dt_2 \int_0^{t_2} dt_1 \partial_{\mathbf{x}} \cdot \partial_{\mathbf{y}} \left[\mathcal{K}_{ba}(\mathbf{x}, t_2; \mathbf{y}, t_1) - \mathcal{K}_0(\mathbf{x}, t_2; \mathbf{y}, t_1) \right]_{\mathbf{x}=\mathbf{y}=0}$$



3-body interactions of the medium via Schrödinger equation with potential:

$$v(\mathbf{x}, t) = \int_{\mathbf{q}} \frac{d\sigma_{el}}{d^2\mathbf{q}} (1 - e^{i\mathbf{q}\cdot\mathbf{x}}) \approx \underbrace{\frac{1}{4} \hat{q} x^2}_{\text{HO}} \log \frac{4}{\mu^2 x^2}$$

Solved numerically in Caron-Huot, Gale 1006.2379, Feal, Vazquez 1811.01591, Ke, Xu and Bass 1810.08177, Feal, Salgado, Vazquez 1911.01309

$$\mathcal{K}(t_2, t_1) = \mathcal{K}_0(t_2 - t_1) + \int dt \mathcal{K}_0(t_2 - t) v(t) \mathcal{K}(t, t_1)$$

opacity expansion


Wiedemann (2000); Gyulassy, Levai, Vitev (2001); Sievert, Vitev, Yoon 1903.06170
Wang, Gu (2001), Majumder 0912.2987
Mehtar-Tani 1903.00506, Mehtar-Tani KT 1910.02032

Iqbal (Wed 15:40)
Djordjevic (Wed 10:20)
Hauksson (Tue 15:20)

Unified understanding of in-medium radiative processes!



REGIMES



Momentum broadening $\langle k^2 \rangle \sim \hat{q}t$ leads to modified bremsstrahlung spectrum \rightarrow no collinear divergence!

$$\omega \frac{dI}{d\omega} = \frac{\alpha_s C_R L}{\pi t_f} = \frac{\alpha_s C_R}{\pi} \sqrt{\frac{\hat{q} L^2}{\omega}}$$


Bethe-Heitler regime ($t_f \lesssim \lambda$)

LPM regime ($\lambda < t_f < L$)

N=1 regime ($L < t_f \sim E/\mu^2$)



REGIMES



Momentum broadening $\langle k^2 \rangle \sim \hat{q}t$ leads to modified bremsstrahlung spectrum \rightarrow no collinear divergence!

$$\omega \frac{dI}{d\omega} = \frac{\alpha_s C_R L}{\pi t_f} = \frac{\alpha_s C_R}{\pi} \sqrt{\frac{\hat{q}L^2}{\omega}}$$

Bethe-Heitler regime ($t_f \lesssim \lambda$)

LPM regime ($\lambda < t_f < L$)

N=1 regime ($L < t_f \sim E/\mu^2$)

Soft scale: copious, large angle gluons leading to energy loss & thermalization.



QUENCHING OF PARTONIC HARD SPECTRUM

Baier, Dokshitzer, Mueller, Schiff (2001); Salgado, Wiedemann (2003)

Probability of radiating energy ϵ out of the cone
(1 gluon emission)

$$\mathcal{P}(\epsilon) \simeq \delta(\epsilon) \left[1 - \int_0^\infty d\omega \frac{dI_{>}}{d\omega} \right] + \frac{dI_{>}}{d\epsilon}$$



QUENCHING OF PARTONIC HARD SPECTRUM

Baier, Dokshitzer, Mueller, Schiff (2001); Salgado, Wiedemann (2003)

Probability of radiating energy ϵ out of the cone
(1 gluon emission)

$$\mathcal{P}(\epsilon) \simeq \delta(\epsilon) \left[1 - \int_0^\infty d\omega \frac{dI_{>}}{d\omega} \right] + \frac{dI_{>}}{d\epsilon}$$

$$\frac{d\sigma_{\text{med}}}{dp_T} = \int_0^\infty d\epsilon \mathcal{P}(\epsilon) \left. \frac{d\sigma_{\text{vac}}}{dp'_T} \right|_{p'_T=p_T+\epsilon} \approx \frac{d\sigma_{\text{vac}}}{dp_T} \underbrace{\int_0^\infty d\epsilon \mathcal{P}(\epsilon) e^{-\epsilon \frac{n}{p_T}}}_{Q(p_T)}$$

quenching factor



QUENCHING OF PARTONIC HARD SPECTRUM


Baier, Dokshitzer, Mueller, Schiff (2001); Salgado, Wiedemann (2003)

Probability of radiating energy ϵ out of the cone
(1 gluon emission)

$$\mathcal{P}(\epsilon) \simeq \delta(\epsilon) \left[1 - \int_0^\infty d\omega \frac{dI_{>}}{d\omega} \right] + \frac{dI_{>}}{d\epsilon}$$


$$\frac{d\sigma_{\text{med}}}{dp_T} = \int_0^\infty d\epsilon \mathcal{P}(\epsilon) \frac{d\sigma_{\text{vac}}}{dp'_T} \Big|_{p'_T=p_T+\epsilon} \approx \frac{d\sigma_{\text{vac}}}{dp_T} \underbrace{\int_0^\infty d\epsilon \mathcal{P}(\epsilon) e^{-\epsilon \frac{n}{p_T}}}_{Q(p_T)}$$

quenching factor




- full $\mathcal{P}(\epsilon)$ resums multiple soft gluon emission via rate equation
- generalizes for N partons $Q \rightarrow Q^N$ (incoherent case)





1 emitter (coherent)
factorisation



n emitters (partially incoherent)
factorisation breaking

new element: importance of jet substructure fluctuations!

also seen in MC studies: Milhano, Zapp 1512.08107; Casalderrey-Solana, Milhano, Pablos, Rajagopal 1808.07386

Ringer (Wed 09:40)



VACUUM RADIATION IN MEDIUM

Vacuum radiation at short timescales was considered first in the context of antenna radiation.

Mehtar-Tani, Salgado, KT PRL (2010), PLB (2012), JHEP (2013); Casalderrey, Iancu JHEP (2011)

Dominguez, Milhano, Salgado, KT, Vila 1907.03653



VACUUM RADIATION IN MEDIUM


Vacuum radiation at short timescales was considered first in the context of antenna radiation.

Mehtar-Tani, Salgado, KT PRL (2010), PLB (2012), JHEP (2013); Casalderrey, Iancu JHEP (2011)

Dominguez, Milhano, Salgado, KT, Vila 1907.03653

First step: calculation of decoherence and energy loss of an initially color correlated pair

Mehtar-Tani, KT 1706.06047



VACUUM RADIATION IN MEDIUM


Vacuum radiation at short timescales was considered first in the context of antenna radiation.

Mehtar-Tani, Salgado, KT PRL (2010), PLB (2012), JHEP (2013); Casalderrey, Iancu JHEP (2011)

Dominguez, Milhano, Salgado, KT, Vila 1907.03653

First step: calculation of decoherence and energy loss of an initially color correlated pair

Mehtar-Tani, KT 1706.06047



VACUUM RADIATION IN MEDIUM


Vacuum radiation at short timescales was considered first in the context of antenna radiation.

Mehtar-Tani, Salgado, KT PRL (2010), PLB (2012), JHEP (2013); Casalderrey, Iancu JHEP (2011)

Dominguez, Milhano, Salgado, KT, Vila 1907.03653

First step: calculation of decoherence and energy loss of an initially color correlated pair

Mehtar-Tani, KT 1706.06047



$$t_d \sim (\hat{q}\theta_{12}^2)^{-1/3}$$

$$\theta_c \sim \sqrt{\frac{1}{\hat{q}L^3}}$$

$$Q_2(p_T) = Q(p_T) \times Q_{\text{sing}}(p_T)$$


energy loss of total
color charge

delayed energy loss from
resolved partons



PHASE SPACE ANALYSIS

Y. Mehtar-Tani, KT 1706.06047, 1707.07361
 Caucal, Iancu, Mueller, Soyez 1801.09703




Red area: vacuum emissions (w/ $k_t^2 > \sqrt{\hat{q}\omega}$ and $\theta > \theta_c$) that are not resolved by the medium.

At border: all inside emissions get quenched



PHASE SPACE ANALYSIS

Y. Mehtar-Tani, KT 1706.06047, 1707.07361
 Caucal, Iancu, Mueller, Soyez 1801.09703



Red area: vacuum emissions (w/ $k_t^2 > \sqrt{\hat{q}\omega}$ and $\theta > \theta_c$) that are not resolved by the medium.


At border: all inside emissions get quenched

How many modes are emitted inside?



PHASE SPACE ANALYSIS

Y. Mehtar-Tani, KT 1706.06047, 1707.07361
 Caucal, Iancu, Mueller, Soyez 1801.09703



Red area: vacuum emissions (w/ $k_t^2 > \sqrt{\hat{q}\omega}$ and $\theta > \theta_c$) that are not resolved by the medium.

At border: all inside emissions get quenched

How many modes are emitted inside?

$$\begin{aligned}
 (\text{PS})_{\text{in}} &= \int_0^R \frac{d\theta}{\theta} \int_0^1 dz \frac{\alpha_s}{\pi} P(z) \Theta_{\text{in}} \\
 &\approx 2 \frac{\alpha_s C_R}{\pi} \log \frac{R}{\theta_c} \left(\log \frac{p_T}{\omega_c} + \frac{2}{3} \log \frac{R}{\theta_c} \right)
 \end{aligned}$$


Potentially large and **needs to be resummed.**

No coherence leads to stronger $\sim \log^2 p_T$ dependence!



PHASE SPACE ANALYSIS

Y. Mehtar-Tani, KT 1706.06047, 1707.07361
 Caucal, Iancu, Mueller, Soyez 1801.09703



Red area: vacuum emissions (w/ $k_t^2 > \sqrt{\hat{q}\omega}$ and $\theta > \theta_c$) that are not resolved by the medium.

At border: all inside emissions get quenched

How many modes are emitted inside?

$$\begin{aligned}
 (\text{PS})_{\text{in}} &= \int_0^R \frac{d\theta}{\theta} \int_0^1 dz \frac{\alpha_s}{\pi} P(z) \Theta_{\text{in}} \\
 &\approx 2 \frac{\alpha_s C_R}{\pi} \log \frac{R}{\theta_c} \left(\log \frac{p_T}{\omega_c} + \frac{2}{3} \log \frac{R}{\theta_c} \right)
 \end{aligned}$$

Potentially large and **needs to be resummed.**

No coherence leads to stronger $\sim \log^2 p_T$ dependence!




Caucal (Wed 15:20)

Non-trivial correlation between jet energy loss and substructure!

EFFECTIVE THEORY OF JET QUENCHING

Mehtar-Tani, KT in preparation

A probabilistic picture can be established due to the separation of jet and medium scales.




EFFECTIVE THEORY OF JET QUENCHING

Mehtar-Tani, KT in preparation

A probabilistic picture can be established due to the separation of jet and medium scales.

Outside of the medium, we have AO vacuum (vetoed shower):

$$Z_{\text{out}}(p, R) = u(p) + \int^R d\Pi (1 - \Theta_{\text{in}}) [Z_{\text{out}}(zp, \theta) Z_{\text{out}}((1-z)p, \theta) - Z_{\text{out}}(p, \theta)]$$



EFFECTIVE THEORY OF JET QUENCHING

Mehtar-Tani, KT in preparation


A probabilistic picture can be established due to the separation of jet and medium scales.

Outside of the medium, we have AO vacuum (vetoed shower):

$$Z_{\text{out}}(p, R) = u(p) + \int^R d\Pi (1 - \Theta_{\text{in}}) [Z_{\text{out}}(zp, \theta) Z_{\text{out}}((1-z)p, \theta) - Z_{\text{out}}(p, \theta)]$$

Vacuum-like (vetoed) AO shower inside the medium

$$Z(p, R) = Q(p_T) Z_{\text{out}}(p, 1) + \int^R d\Pi \Theta_{\text{in}} [Z(zp, \theta) Z((1-z)p, \theta) - Z(p, \theta)]$$



EFFECTIVE THEORY OF JET QUENCHING

Mehtar-Tani, KT in preparation

A probabilistic picture can be established due to the separation of jet and medium scales.


Outside of the medium, we have AO vacuum (vetoed shower):

$$Z_{\text{out}}(p, R) = u(p) + \int^R d\Pi (1 - \Theta_{\text{in}}) [Z_{\text{out}}(zp, \theta) Z_{\text{out}}((1-z)p, \theta) - Z_{\text{out}}(p, \theta)]$$

Vacuum-like (vetoed) AO shower inside the medium

$$Z(p, R) = Q(p_T) Z_{\text{out}}(p, 1) + \int^R d\Pi \Theta_{\text{in}} [Z(zp, \theta) Z((1-z)p, \theta) - Z(p, \theta)]$$

out-shower restarted from max angle (AAO)




EFFECTIVE THEORY OF JET QUENCHING

Mehtar-Tani, KT in preparation

A probabilistic picture can be established due to the separation of jet and medium scales.

Outside of the medium, we have AO vacuum (vetoed shower):

$$Z_{\text{out}}(p, R) = u(p) + \int^R d\Pi (1 - \Theta_{\text{in}}) [Z_{\text{out}}(zp, \theta) Z_{\text{out}}((1-z)p, \theta) - Z_{\text{out}}(p, \theta)]$$



Vacuum-like (vetoed) AO shower inside the medium

$$Z(p, R) = Q(p_T) Z_{\text{out}}(p, 1) + \int^R d\Pi \Theta_{\text{in}} [Z(zp, \theta) Z((1-z)p, \theta) - Z(p, \theta)]$$

out-shower restarted from max angle (AAO)

accounts for quenching through iterative use of GF



RESUMMED QUENCHING WEIGHT

Mehtar-Tani, KT 1707.07361
Mehtar-Tani, KT in preparation

Non-trivial normalization of the GF: $Z(p, R; \{u = 1\}) = Q(p, R)!$

Mismatch between real and virtual diagrams!



RESUMMED QUENCHING WEIGHT

Mehtar-Tani, KT 1707.07361
Mehtar-Tani, KT in preparation

Non-trivial normalization of the GF: $Z(p, R; \{u = 1\}) = Q(p, R)$!

Mismatch between real and virtual diagrams!

$$Q(p, R) = \underbrace{Q(p_T)} + \int^R d\Pi \Theta_{\text{in}} [Q(zp, \theta)Q((1-z)p, \theta) - Q(p, \theta)]$$

non-linear evolution of quenching



RESUMMED QUENCHING WEIGHT



Mehtar-Tani, KT 1707.07361
Mehtar-Tani, KT in preparation

Non-trivial normalization of the GF: $Z(p, R; \{u = 1\}) = Q(p, R)$!

Mismatch between real and virtual diagrams!

$$Q(p, R) = \underbrace{Q(p_T)}_{\text{non-linear evolution of quenching}} + \int^R d\Pi \Theta_{\text{in}} [Q(zp, \theta)Q((1-z)p, \theta) - Q(p, \theta)]$$

non-linear evolution of quenching



THE MICROJET SPECTRUM IN MEDIUM

Mehtar-Tani, KT in preparation

R-dependence of the jet spectrum in terms of a jet fragmentation function

$$\frac{d\sigma^{\text{jet}}(R)}{dp_T} = \sum_{i=q,g} \frac{d\sigma_i}{dp_T} \int_0^1 dz z^{n_i(p_T)-1} f_{\text{jet}/i}(z, t)$$



THE MICROJET SPECTRUM IN MEDIUM

Mehtar-Tani, KT in preparation

R-dependence of the jet spectrum in terms of a jet fragmentation function

$$\frac{d\sigma^{\text{jet}}(R)}{dp_T} = \sum_{i=q,g} \frac{d\sigma_i}{dp_T} \int_0^1 dz z^{n_i(p_T)-1} f_{\text{jet}/i}(z, t)$$

$$\begin{aligned} Q_q(p, 1) f_{\text{jet}/q}(z, p, t) &= Q_q(p, R) \delta(1 - z) \\ &+ \int d\Pi \Theta_{\text{in}} \frac{1}{z'} P_{qq}(z') f_{\text{jet}/q}\left(\frac{z}{z'}, z' p, t'\right) Q_q(z' p, \theta') Q_g((1 - z') p, \theta') \\ &- \int d\Pi \Theta_{\text{in}} P_{qq}(z') f_{\text{jet}/q}(z, p, t') Q_q(p, \theta') \end{aligned}$$

Medium modified DGLAP



THE MICROJET SPECTRUM IN MEDIUM

Mehtar-Tani, KT in preparation

R-dependence of the jet spectrum in terms of a jet fragmentation function


$$\frac{d\sigma^{\text{jet}}(R)}{dp_T} = \sum_{i=q,g} \frac{d\sigma_i}{dp_T} \int_0^1 dz z^{n_i(p_T)-1} f_{\text{jet}/i}(z, t)$$

$$Q_q(p, 1) f_{\text{jet}/q}(z, p, t) = Q_q(p, R) \delta(1 - z)$$

$$+ \int d\Pi \Theta_{\text{in}} \frac{1}{z'} P_{qq}(z') f_{\text{jet}/q}\left(\frac{z}{z'}, z'p, t'\right) Q_q(z'p, \theta') Q_g((1-z')p, \theta')$$

$$- \int d\Pi \Theta_{\text{in}} P_{qq}(z') f_{\text{jet}/q}(z, p, t') Q_q(p, \theta')$$

Taylor (Wed 11:20)



Medium modified DGLAP

- non-trivial evolution with jet cone size reproduced in pp and AA!
- sensitive to the of recapturing of energy at large angles and multi-parton quenching



OUTLOOK



OUTLOOK

- improved theoretical control of radiative mechanism of energy loss



OUTLOOK

- improved theoretical control of radiative mechanism of energy loss
- hierarchy of jet vs medium scales established
 - scaling of energy loss with relevant phase space
 - probabilistic interpretation



OUTLOOK

- improved theoretical control of radiative mechanism of energy loss
- hierarchy of jet vs medium scales established
 - scaling of energy loss with relevant phase space
 - probabilistic interpretation
- incoherent energy loss breaks factorization
 - resummed quenching weight
 - medium modified DGLAP




OUTLOOK

- improved theoretical control of radiative mechanism of energy loss
- hierarchy of jet vs medium scales established
 - scaling of energy loss with relevant phase space
 - probabilistic interpretation
- incoherent energy loss breaks factorization
 - resummed quenching weight
 - medium modified DGLAP
- jet R_{AA} spectra well described: promising for future substructure studies



BACKUP

OBSERVABLES FROM PRIMARY EMISSIONS




Soft Drop/mMDT

[Dasgupta, Fregoso, Marzani, Salam1307.0007;
Larkoski, Marzani, Soyez, Thaler 1402.2657]

- find first primary branch in the tree that satisfies $z > z_{\text{cut}}(\theta/R)^\beta$
- many observables: $z_{g'}$, m_g and n_{SD}




OBSERVABLES FROM PRIMARY EMISSIONS



Soft Drop/mMDT

[Dasgupta, Fregoso, Marzani, Salam1307.0007;
Larkoski, Marzani, Soyez, Thaler 1402.2657]

- find first primary branch in the tree that satisfies $z > z_{\text{cut}}(\theta/R)^\beta$
- many observables: $z_{g'}$, m_g and n_{SD}



NEW! Dynamical grooming

[Mehtar-Tani, Soto-Ontoso, KT 1911.00375]

- find hardest branch in the tree,


$$\kappa^{(a)} = \max_{i \in C/A} z_i(1 - z_i)(\theta/R)^a$$
- earlier emissions interpreted as radiation off total color charge.



OBSERVABLES FROM PRIMARY EMISSIONS

Zardoshti (Tue 09:20)


Havener (Wed 09:00)



Soft Drop/mMDT

[Dasgupta, Fregoso, Marzani, Salam1307.0007;
Larkoski, Marzani, Soyez, Thaler 1402.2657]

- find first primary branch in the tree that satisfies $z > z_{\text{cut}}(\theta/R)^\beta$
- many observables: z_g , m_g and n_{SD}




NEW! Dynamical grooming

[Mehtar-Tani, Soto-Ontoso, KT 1911.00375]

- find hardest branch in the tree,
 $\kappa^{(a)} = \max_{i \in C/A} z_i(1 - z_i)(\theta/R)^a$
- earlier emissions interpreted as radiation off total color charge.



EMERGING PROBABILISTIC PICTURE




[See also Caucal, Iancu, Soyez, Mueller 1801.09703]



TWO SEPARATED REGIMES

see also Kurkela, Wiedemann | 407.0293

Energy loss for jets is driven by out-of-cone emissions.



$$\omega \sim \omega_c = \hat{q}L^2$$

$$N \sim \mathcal{O}(\alpha_s)$$

$$\theta_{\text{br}} \sim \theta_c = (\hat{q}L^3)^{-1/2}$$

perturbative: rare, small-angle radiation
intra-jet structure modifications

$$\omega \sim \omega_s = \alpha_s^2 \hat{q}L^2$$

$$N \sim 1$$

$$\theta_{\text{br}} \sim \frac{1}{\alpha_s^2} \theta_c$$


non-perturbative: copious, large-angle
emissions

out-of-cone energy-loss, thermalization

TWO SEPARATED REGIMES

see also Kurkela, Wiedemann | 407.0293

Energy loss for jets is driven by out-of-cone emissions.



$$\omega \sim \omega_c = \hat{q}L^2$$

$$N \sim \mathcal{O}(\alpha_s)$$

$$\theta_{\text{br}} \sim \theta_c = (\hat{q}L^3)^{-1/2}$$

perturbative: rare, small-angle radiation
intra-jet structure modifications

$$\omega \sim \omega_s = \alpha_s^2 \hat{q}L^2$$

$$N \sim 1$$

$$\theta_{\text{br}} \sim \frac{1}{\alpha_s^2} \theta_c$$

non-perturbative: copious, large-angle
emissions

out-of-cone energy-loss, thermalization

Gluons with $\omega_s \sim 5 \text{ GeV}$ melt into the plasma!

Blaizot, Dominguez, Iancu, Mehtar-Tani | 301.6102



SUPPRESSION OF HARD SPECTRUM

Simplified example: consider a $1 \rightarrow N$ process followed by energy loss.

$$\frac{d\sigma_{\text{excl}}}{dk_1 \dots dk_N} = \int dp \left\{ \prod_i \int d\epsilon_i \mathcal{P}(\epsilon_i) \right\} f_{1 \rightarrow N}(k_1 + \epsilon_1, \dots, k_N + \epsilon_N | p) \delta\left(p - \sum_i k_i - \sum_i \epsilon_i\right) \frac{d\sigma_0}{dp}$$

Leading-order effect:
$$\frac{d\sigma_{\text{excl}}}{dk_1 \dots dk_N} \simeq \left(\int d\epsilon \mathcal{P}(\epsilon) e^{-n\epsilon/p} \right)^N \frac{d\sigma_{0,\text{excl}}}{dk_1 \dots dk_N} + \dots$$


systematically improvable...

Provides a framework for studying multi-parton energy-loss effects for inclusive observables !



RECOMBINATION ALGORITHMS


Pairwise clustering of particles that minimize $d_{ij} = \min(p_{T,i}^{2\alpha}, p_{T,j}^{2\alpha}) \Delta R_{ij}^2 / R^2$



1) Cambridge/Aachen (CA)

[Dokshitzer, Leder, Moretti, Webber (1997)]


- only angular measure ($\alpha=0$)
- ideal for substructure measurements



2) k_t algorithm

[Catani, Dokshitzer, Seymour, Webber (1993); Ellis, Soper (1993)]

- k_t weighted metric ($\alpha = 1$)
- sensitive to soft activity



3) anti- k_t algorithm

[Cacciari, Salam, Soyez (2008)]

- anti- k_t weighted metric ($\alpha = -1$)
- resilient to soft activity, ideal for identifying candidate jets

Jets are separated whenever beam distance is shortest $d_{iB} = p_{T,i}^{2\alpha}$

