

Quark Matter 2019 - the XXVIIIth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions



Contribution ID: 564

Type: **Poster Presentation**

Multiplicity and rapidity dependence of v_2 in small system by using two-particle correlations

Monday 4 November 2019 17:40 (20 minutes)

Recent PHENIX measurements indicate that the initial geometry is the cause of the observed positive v_2 and v_3 in high-multiplicity p+Au, d+Au, and $^3\text{He}+\text{Au}$ collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV. These results were obtained using the event-plane method, with the event plane determined in the backward rapidity range and correlated with particles in other sub-events at mid- and forward-rapidity. In this poster, we present the latest PHENIX measurements, in which we employ the two-particle correlation method and investigate the effects of the size of the rapidity gap between particles as well as different non-flow subtraction methods in order to understand possible non-flow contributions to the observed v_2 . We also extend the measurements of v_2 from most-central to peripheral collisions to understand the centrality evolution of v_2 in small systems. In this poster, we will present the current analysis status of v_2 measurement via the two particle correlation analysis with various subtraction methods in a wide centrality range of p+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV.

Author: XU, Qiao (Vanderbilt University (US))

Presenter: XU, Qiao (Vanderbilt University (US))

Session Classification: Poster Session

Track Classification: Small systems