

Highlights from the GlueX experiment

Lubomir Pentchev for the
GlueX Collaboration





*“First measurements of near-threshold J/ψ exclusive photoproduction off the proton” PRL 123, 072001 (2019)
(probing gluonic structure of the proton at high x)*


Introduction




- Overview of Hall D/Gluex apparatus
- Selected results from GlueX/Hall D experiments:
 - searching for hybrid mesons
 - beam asymmetries
 - η decay via Primakoff reaction
- Near-threshold J/ψ photoproduction:
 - the experiment
 - proton gluonic distributions at high-x
 - search for LHCb pentaquarks
 - future expected results

Experiment	Description	PAC days complete
GlueX-I	Spectroscopy of light and hybrid mesons (low intensity)	 80+40 80+40
GlueX-II	Spectroscopy of hadrons with strange quarks (high intensity)	220+
PrimEx-eta	Eta radiative decay width	 79 24
CPP	Charge pion polarizability	25
JEF	Rare eta decays	-
SRC	Short-range correlations with real photon beams	15

Hall D Apparatus




- Photon beam from coherent Bremsstrahlung off thin diamond
- Photon energy tagged by scattered electron $\sim 0.2\%$ resolution
- Beam collimated at 75m, $<35 \mu\text{rad}$
- Intensity: $\sim 2 \cdot 10^7 - 5 \cdot 10^7 \gamma/\text{sec}$ above J/ψ threshold (8.2 GeV) – total $\sim 68 \text{ pb}^{-1}$ in 2016-2017 runs (**25% of total statistics up to date**)
- Photons are linearly polarized $\sim 40\%$ at peak; polarization plane angle alternates from run to run: PERP, PARA



GlueX spectrometer


2T-solenoid, LH target

Tracking (FDC,CDC) , Calorimetry (BCAL,FCAL) , Timing (TOF,SC)



- Hermetic detector: $1 - 120^\circ$ polar and full azimuthal acceptance
- Tracking: $\sigma_p/p \sim 1 - 5\%$
- Calorimetry: $\sigma_E/E \sim 6\%/\sqrt{E} + 2\%$

Searching for light hybrid mesons ($q\bar{q}g$)




Hybrids with exotic quantum numbers
(not allowed in $q\bar{q}$ model)

Experiment	$p_{beam}, \text{ GeV}$	Reaction	Resonance
GAMS	32,38,100	$\pi^- p \rightarrow \pi^0 \eta n$	$\pi_1(1400)$
E852	18	$\pi^- p \rightarrow \pi^- \eta^{(\prime)} p$	$\pi_1(1400/1600)$
Crystal Barrel	Annihilation	$\bar{p} n \rightarrow \pi^- \pi^0 \eta$	$\pi_1(1400)$
VES	37	$\pi^- p \rightarrow \pi^- \eta^{(\prime)} p$	$\pi_1(1600)$
COMPASS	190	$\pi^- p \rightarrow \pi^- \eta' p$	$\pi_1(1600)$
COMPASS	190	$\pi^- p \rightarrow \pi^- \eta p$	$\pi_1(1400)$

COMPASS: $\pi_1 \rightarrow \pi^- \eta^{(\prime)}$ best evidence for exotic hybrid





COMPASS, PLB 740 (2015)





Dudek et al. PRD 88 (2013)

Searching for hybrids: JPAC analysis of COMPASS data

Coupled channel analysis finds **one pole for exotic $\pi_1(1564)$** , describing both $\eta\pi$ and $\eta'\pi$ channels in the P-wave intensities






$a_2(1320)$ $a_2'(1700)$



JPAC, PRL 122 (2019)

$\pi^- \eta^{(\prime)}$ can have J^{PC} 0^{++} 1^{-+} 2^{++}

Searching for hybrids: GlueX $\eta\pi/\eta'\pi$ spectroscopy




- With full statistics up to date, GlueX competitive with COMPASS for one channel
- Multiple channels studied in GlueX
- Different production mechanism, different background
- GlueX linearly polarized photon beam – important tool in understanding the $\eta\pi$ spectrum

Beam Asymmetry

$$Y(\phi) \sim (1 - P_\gamma \Sigma \cos 2(\phi_p - \phi_\gamma))$$


Alternating (run by run) the orientation of the polarization by 90° , the efficiency cancels in the asymmetry ratio:

$$\frac{Y_\perp - F_R Y_\parallel}{Y_\perp + F_R Y_\parallel} = P_\gamma \Sigma \cos 2\phi_p$$




$\Sigma = 1$ for natural exchange ($0^+ 1^- 2^+$)

$\Sigma = -1$ for unnatural exchange ($0^- 1^+ 2^-$)



Beam Asymmetry





$$\gamma p \rightarrow (\pi^0, \eta, \eta') p$$

GlueX, PRC 95 (R) (2017) –
first Jlab 12 GeV paper


Natural exchange dominates if $\Sigma \sim 1$

More papers on asymmetries expected in 2019



η radiative decay via Primakoff reaction

$\eta \rightarrow \gamma\gamma$ yield [$8.0 \text{ GeV} < E_\gamma < 11.2 \text{ GeV}$]




Precision measurement of $\Gamma_{\eta \rightarrow \gamma\gamma}$:

significant impact on fundamental QCD parameters (Goldstone nature of light pseudoscalars, light quark mass ratios, chiral anomaly etc.)


Near-threshold J/ψ photoproduction

Exclusive reaction $\gamma p \rightarrow J/\psi p \rightarrow e^+e^-p$



- Near threshold – all valence quarks participate, corresponding to high-x gluon exchange
- Using VMD ($\gamma \rightarrow J/\psi$) one can study $J/\psi p \rightarrow J/\psi p$
- Look for LHCb P_c : $\gamma p \rightarrow P_c \rightarrow J/\psi p$



Near-threshold J/ψ photoproduction



- Electrons identified by E/p
- Tagged photon beam, 0.2% energy resolution
- Kinematic fit: 13 MeV mass resolution
- Presented results (~ 470 J/ψ): 25% of statistics accumulated up to date
- Preliminary results (ERRORS ONLY!) will be shown based on 70% of data

J/ψ total cross-section

Brodsky et al. PL B498 (2000):
 $\sigma(E_\gamma)$ depending on number of
 hard-gluons exchanged
 using dimensional scaling



GlueX errors: quadratic sums of statistical and systematic ones



SLAC data: $\gamma d(p) \rightarrow J/\psi X$ from $d\sigma/dt(t = t_{min})$

$$F^2(t) \sim (1 - t/m_{2g}^2)^{-4} \quad m_{2g} = 1.14 \text{ GeV}$$

Cornell data: $\gamma Be \rightarrow J/\psi X$

GlueX data falls toward threshold less steeply than 2g exchange model

J/ψ differential cross-section – perturbative calculations




Sibirtsev et al. JP G 30 (2004):

- Cross-section is very sensitive to gPDF at high x
- In case of two gluon exchange:

$$d\sigma/dt(t=0) \sim x^2 g^2(x) \text{ where } x = m_{J/\psi}^2/s$$

- The near-threshold energy dependence is defined by the gPDF as $x \rightarrow 1$
- t -dependence not defined by the pQCD calculations (discussed later)
- Comparison to the old data only


J/ψ total cross-section – perturbative calculations



Kharzeev et al. EPJ C (1999):

- Forward J/ψ - p scattering amplitude calculated using gluonic PDF and dispersion relations
- **Real part of the amplitude dominates near threshold, critically important: contains scale anomaly term related to the mass of the proton arising from gluons.**
- Shape of the theoretical curve similar to GlueX data; absolute normalization within the uncertainties of the model

J/ψ differential cross-section and proton gluonic FF



gluonic form factor
(dipole form in analogy
with the e.-m. FF):

$$F(t) \sim 1/(1 - t/m_0^2)^2$$



Frankfurt and Strikman
PRD66 (2002)

	e.m. FF	gluonic FF
reaction	$ep \rightarrow ep$	$J/\psi p \rightarrow J/\psi p$
transverse size of probe	0	$\ll 1 \text{ fm}$
effective mass scale m_0	0.84 GeV (vector meson)	$\sim 1.1 \text{ GeV}$ (two-gluon mass)

J/ψ total cross-section – non-perturbative (holographic) calculations

Red – maximal contribution from gluons, favored by GlueX data

Blue – no gluonic contribution



Y. Hatta, A. Rajan, and D.-L. Yang, arXiv:1906.00894:

Proton gluonic FF: “..these are nothing but the gravitational form factors A_g, B_g, C_g, \bar{C}_g ”


$$\begin{aligned} \langle P' | (T_g)_{\mu}^{\mu} | P \rangle &= \langle P' | \left(\frac{\beta(g)}{2g} F_{\mu\nu}^a F_a^{\mu\nu} + m \gamma_m \bar{\psi} \psi \right) | P \rangle \\ &= \bar{u}(P') \left[A_g M + \frac{B_g}{4M} \Delta^2 - 3 \frac{\Delta^2}{M} C_g + 4 \bar{C}_g M \right] u(P) \end{aligned}$$

A_g, B_g, C_g were recently calculated on lattice:


P. E. Shanahan and W. Detmold, arXiv:1810.04626

LHCb pentaquarks

$$\Lambda_b^0 \rightarrow J/\psi p K^-$$



Phys. Rev. Lett., 115, 072001 (2015)





Phys. Rev. Lett. 122, 222001 (2019)

- J^P of P_c states not determined yet
- Molecules (most likely), but compact states or rescattering effects not excluded


State	M [MeV]	Γ [MeV]	(95% CL)	\mathcal{R} [%]
$P_c(4312)^+$	$4311.9 \pm 0.7_{-0.6}^{+6.8}$	$9.8 \pm 2.7_{-4.5}^{+3.7}$	(< 27)	$0.30 \pm 0.07_{-0.09}^{+0.34}$
$P_c(4440)^+$	$4440.3 \pm 1.3_{-4.7}^{+4.1}$	$20.6 \pm 4.9_{-10.1}^{+8.7}$	(< 49)	$1.11 \pm 0.33_{-0.10}^{+0.22}$
$P_c(4457)^+$	$4457.3 \pm 0.6_{-1.7}^{+4.1}$	$6.4 \pm 2.0_{-1.9}^{+5.7}$	(< 20)	$0.53 \pm 0.16_{-0.13}^{+0.15}$

LHCb pentaquarks and J/ψ photo-production

- If LHCb pentaquarks exist they should be seen in s-channel photoproduction (free of rescattering effects in the final state):




LHCb P_c related to $\gamma p \rightarrow P_c \rightarrow J/\psi p$



- V.Kubarovsky and M.B.Voloshin, PRD 92.031502 (2015).*
- M.Karliner and J.Rosner, arXiv: PLB 752, 329 (2016).*
- A.Blin, C.Fernandez-Ramirez, A.Jackura, V.Mathieu, V.Mokeev, A.Pilloni, and A.Szczepaniak, PRD 94,034002 (2016).*

J/ψ cross-section: model-dependent upper limits

PRL 123, 072001 (2019)



Assuming:

- all P_c independent $J^P = 3/2^-$
- s-channel model:


$$\sigma(\gamma p \rightarrow P_c \rightarrow J/\psi p) \approx 0.35 \mu\text{b} \text{ Br}^2(P_c \rightarrow J/\psi p) (2J+1)$$
- JPAC model for t-channel:
 Pomeron and tensor part extracted at high energies

	$\mathcal{B}(P_c^+ \rightarrow J/\psi p)$ Upper Limits, %		$\sigma_{\text{max}} \times \mathcal{B}(P_c^+ \rightarrow J/\psi p)$ Upper Limits, nb	
	p.t.p. only	total	p.t.p. only	total
$P_c^+(4312)$	2.9	4.6	3.7	4.6
$P_c^+(4440)$	1.6	2.3	1.2	1.8
$P_c^+(4457)$	2.7	3.8	2.9	3.9

Upper limits at 90% confidence level

J/ψ cross-section: model-dependent upper limits

PRL 123, 072001 (2019)



Assuming:

- all P_c independent $J^P = 3/2^-$
- s-channel model:


$$\sigma(\gamma p \rightarrow P_c \rightarrow J/\psi p) \approx 0.35 \mu\text{b} \text{ Br}^2(P_c \rightarrow J/\psi p) (2J+1)$$
- JPAC model for t-channel: Pomeron and tensor part extracted at high energies

	$\mathcal{B}(P_c^+ \rightarrow J/\psi p)$ Upper Limits, %		$\sigma_{\text{max}} \times \mathcal{B}(P_c^+ \rightarrow J/\psi p)$ Upper Limits, nb	
	p.t.p. only	total	p.t.p. only	total
$P_c^+(4312)$	2.9	4.6	3.7	4.6
$P_c^+(4440)$	1.6	2.3	1.2	1.8
$P_c^+(4457)$	2.7	3.8	2.9	3.9

Upper limits at 90% confidence level

J/ψ cross-section: model-dependent upper limits

PRL 123, 072001 (2019)



Assuming:


- all P_c independent $J^P = 3/2^-$
- s-channel model:

$$\sigma(\gamma p \rightarrow P_c \rightarrow J/\psi p) \approx 0.35 \mu\text{b} \text{ Br}^2(P_c \rightarrow J/\psi p) (2J+1)$$
- JPAC model for t-channel: Pomeron and tensor part extracted at high energies

	$\mathcal{B}(P_c^+ \rightarrow J/\psi p)$ Upper Limits, %		$\sigma_{\text{max}} \times \mathcal{B}(P_c^+ \rightarrow J/\psi p)$ Upper Limits, nb	
	p.t.p. only	total	p.t.p. only	total
$P_c^+(4312)$	2.9	4.6	3.7	4.6
$P_c^+(4440)$	1.6	2.3	1.2	1.8
$P_c^+(4457)$	2.7	3.8	2.9	3.9

Upper limits at 90% confidence level

Lower limits on $\text{Br}(P_c \rightarrow J/\psi p)$ from data?



X. Cao, J-P. Dai *arXiv:1904.06015*


$$\mathcal{R} = \frac{\mathcal{B}(\Lambda_b^0 \rightarrow P_c^+ K^-) \mathcal{B}(P_c^+ \rightarrow J/\psi p)}{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-)}$$

$$\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-) = (3.2^{+0.6}_{-0.5}) \times 10^{-4}$$

$\mathcal{B}(\Lambda_b \rightarrow P_c^+ K^-) < 10^{-3}$ at the level of $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)$ and $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-)$

(model dependent 2-4%) $\mathcal{B}(P_c^+ \rightarrow J/\psi p) > 0.05\%$
 GlueX

Lower limits on $\text{Br}(P_c \rightarrow J/\psi p)$ from data?



X. Cao, J-P. Dai *arXiv:1904.06015*

$$\mathcal{R} = \frac{\mathcal{B}(\Lambda_b^0 \rightarrow P_c^+ K^-) \mathcal{B}(P_c^+ \rightarrow J/\psi p)}{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-)}$$

$$\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-) = (3.2^{+0.6}_{-0.5}) \times 10^{-4}$$

$\mathcal{B}(\Lambda_b \rightarrow P_c^+ K^-) < 10^{-3}$ at the level of $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)$ and $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-)$


(model dependent 2-4%) $> \mathcal{B}(P_c^+ \rightarrow J/\psi p) > 0.05\%$
 GlueX

Outlook

- First phase of GlueX experiment finished and already producing results.
- Next phase – strangeness program and higher intensity.
- First measurement of near-threshold J/ψ exclusive photoproduction – important input to models of the gluonic structure of the proton at high X .
- Do not see evidence for LHCb pentaquarks and set model-dependent limits on $\text{Br}(P_c \rightarrow J/\psi p)$ at several percent level. This allows us to discriminate between different pentaquark models.
- Expect results by the end of the year with 4 times more statistics



Back-ups

J/ψ total cross-section – future results



Preliminary results with 70% of statistics up to date –
only errors (stat.) shown


J/ ψ differential cross-section – perturbative calculations



Kharzeev et al. EPJ C (1999):

- Forward J/ ψ -p scattering amplitude calculated using gluonic PDF and dispersion relations.
- Very sensitive to gPDF at high x
- real part of the amplitude dominates, contains scale anomaly term related to the mass of the proton arising from gluons.

Searching for hybrids: light mesons lattice QCD



Br($P_c \rightarrow J/\psi p$) calculations: pentaquark models

model	Γ_{P_c} , MeV	$\Gamma_{J/\psi p}$, MeV	$\mathcal{B}(P_c \rightarrow J/\psi p)$	J^P	reference
molecular (OPE) $\Sigma_c \bar{D}^{(*)}$	21.7 (4450)	0.03 (4450)	0.14% (4450)	1/2 ⁻ (4312)	M.Eides and V.Petrov
				1/2 ⁻ (4440)	Phys.Rev.D98, 114037
				3/2 ⁻ (4457)	
hadro- charmonium	– (4312)	suppr.(4312)	suppr. (4312)	1/2 ⁺ (4312)	same as above
	44.8 (4440)	11 (4440)	25% (4440)	1/2 ⁻ (4440)	and M.Eides, V.Petrov
	16.2 (4457)	11 (4457)	68% (4457)	3/2 ⁻ (4457)	M.Polyakov,arXiv:1904.1161
compact diquark	–	suppressed	suppressed	3/2 ⁻ (4312)	A.Ali, A.Parkhomenko
				3/2 ⁺ (4440)	Phys.Lett.B793, 365
				5/2 ⁺ (4457)	
molecular (ERE) $\Sigma_c \bar{D}^{(*)}$	9.8* (4312)	6.5	66%	1/2 ⁻ (4312)	Z.-H. Guo and J.Oller
	20.6* (4440)	16.3	79%	1(3)/2 ⁻ (4440)	Phys.Lett.B793, 144
	6.4* (4457)	3.5	55%	1(3)/2 ⁻ (4457)	
molecular (DSE) $\Sigma_c \bar{D}^{(*)}$	15.2 (4306)	4**	26%	1/2 ⁻ (4306)	C.Xiao, J.Nieves, E.Oset,
	23.4 (4453)	18**	77%	1/2 ⁻ (4453)	arxiv:1904.01296
	3.0(4453)	2**	67%	3/2 ⁻ (4453)	Phys.Rev.D88, 056012


* The total width measured by LHCb has been used.

** The width calculated from coupling constants.


Br($P_c \rightarrow J/\psi p$) calculations: molecular vs hadrocharmonium

model	Γ_{P_c} , MeV	$\Gamma_{J/\psi p}$, MeV	$\mathcal{B}(P_c \rightarrow J/\psi p)$	J^P	reference
molecular (OPE) $\Sigma_c \bar{D}^*$	21.7 (4450)	0.03 (4450)	0.14% (4450)	$1/2^-$ (4312)	M.Eides and V.Petrov
				$1/2^-$ (4440)	Phys.Rev.D98, 114037
				$3/2^-$ (4457)	
hadro- charmonium	– (4312)	suppr.(4312)	suppr. (4312)	$1/2^+$ (4312)	same as above
	44.8 (4440)	11 (4440)	25% (4440)	$1/2^-$ (4440)	and M.Eides, V.Petrov
	16.2 (4457)	11 (4457)	68% (4457)	$3/2^-$ (4457)	M.Polyakov,arXiv:1904.1161

all subsystems in color singlet states




Lower limits on pentaquark BR



Br($P_c \rightarrow J/\psi p$) calculations: compact diquark

model	Γ_{P_c} , MeV	$\Gamma_{J/\psi p}$, MeV	$\mathcal{B}(P_c \rightarrow J/\psi p)$	J^P	reference
compact diquark	–	suppressed	suppressed	$3/2^-$ (4312)	A.Ali, A.Parkhomenko
				$3/2^+$ (4440)	Phys.Lett.B793, 365
				$5/2^+$ (4457)	





diquarks in color anti-triplet states

The bound-state effect in (u \bar{c})-diquark reduces the probability to form $c\bar{c}$ -state

Attempts to suppress VMD coupling

J.-J. Wu, T.-S. Lee, B.-S. Zou arXiv:1906.05375




$$F_V(q) = \frac{\Lambda_V^4}{\Lambda_V^4 + (q^2 - m_V^2)^2}$$


$$\Lambda = 0.55 \text{ GeV}$$

- J/ψ is suppressed by 10^{-3} , VMD coupling dominated by ρ and ω
- How to explain J/ψ photoproduction at high energies with such suppression???
- Other papers (*J. Phys. G4* (1978) 989, *Phys. Rev. Lett.* 38 (1977) 263) suggest some moderate suppression (factor of 2-3)

Proton Gluonic Form Factors: A,B,C (lattice calculations)



(a)



(b)

Fits in dipole form:

	m (GeV)	α
A_g	1.13(6)	0.58(5)
D_g	0.48(5)	-10(3)

