

Explaining the ANITA events by a $L_e - L_\tau$ gauge model

Yasaman Farzan

Physics school, IPM, Tehran

ICTP



Multi-messenger era

- Photon detectors in different electromagnetic spectrum: INTEGRAL, FERMI-LAT, HESS, SKA and etc
- Cosmic ray: AMS-02, AUGER, VOYAGER, etc
- Gravitational Wave (GW) detectors: LIGO, VIRGO, etc
- Neutrino Telescopes: ICECUBE, ANTARES, KM3NET, [ANITA](#)

Unprecedented opportunity

Explore cosmos as well as exotic properties of particles

Unprecedented opportunity

Explore cosmos as well as exotic properties of particles

We crave for anomalies.

Unprecedented opportunity

Explore cosmos as well as exotic properties of particles


We crave for anomalies.

ANITA has provided us with anomalies.

ANITA

ANtarctic Impulse Transient Antenna (ANITA)


Flying over an altitude of 37 km



Radio detectors



ANITA's signatures



Two anomalous events

TABLE I: *ANITA-I,-III anomalous upward air showers.*

| event, flight | 3985267, ANITA-I | 15717147, ANITA-III |
|---------------------------|--|---|
| date, time | 2006-12-28,00:33:20UTC | 2014-12-20,08:33:22.5UTC |
| Lat., Lon. ⁽¹⁾ | -82.6559, 17.2842 | -81.39856, 129.01626 |
| Altitude | 2.56 km | 2.75 km |
| Ice depth | 3.53 km | 3.22 km |
| El., Az. | $-27.4 \pm 0.3^\circ$, $159.62 \pm 0.7^\circ$ | $-35.0 \pm 0.3^\circ$, $61.41 \pm 0.7^\circ$ |
| RA, Dec ⁽²⁾ | 282.14064, +20.33043 | 50.78203, +38.65498 |
| $E_{shower}^{(3)}$ | 0.6 ± 0.4 EeV | $0.56_{-0.2}^{+0.3}$ EeV |


ANITA collaboration, PRL 117 (2017) 071101

Extensive Air Shower
(EAS)

ANITA collaboration,
PRL 121 (2018), no 16, 161102

Event characteristics

- Zenith angles: $117.4^\circ \pm 0.3^\circ$ and $125^\circ \pm 0.3^\circ$
- Chord sizes: 5800 km and 7300 km
- Energies 0.6 ± 0.4 EeV and $0.56_{-0.2}^{+0.4}$ EeV.



Gandhi et al,
Astropart phys 5
(1996) 81-110

m



$$s = m^2 + m_\nu^2 + 2mE_\nu$$


$$E_\nu = 0.6 \text{ EeV}$$

$$m = m_e \quad \sqrt{s} \sim TeV \gg m_Z$$

For partons

$$\sqrt{s} \sim \sqrt{2xm_p E_\nu}$$

$$\sigma \propto \frac{1}{s} \quad \text{????}$$



Gandhi et al,
Astropart phys 5
(1996) 81-110

Mean free path for neutrino

- The mean free path in mantle for neutrinos of energy of EeV
500-800 km << chord size  Neutrinos will be absorbed.
The probability of neutrinos passing the Earth is $\sim 10^{-9}$.


Neutrino recreation

$$\nu_\tau \rightarrow \tau \rightarrow \nu_\tau \rightarrow \dots \rightarrow \nu_\tau$$

$$P \sim 10^{-7}$$

See however, Cumming et al, arXiv:1910.00992

Bounds from AUGER and ICECUBE




Transient source
ICECUBE collaboration,
arXiv: 1908.08060

See also, Safa, Pizzuto, Arguelles, Halzen,
Hussain, Kheirandish, Vandenbroucke,
arXiv: 1909.10487

ANITA's signatures

Slides by Jiwoo Nam for ANITA collaboration, ICRC 2017



| Origin | RF production | Polarization | RF Direction | Polarity |
|------------------|-----------------|--------------|---------------|----------|
| 1) Neutrinos | Askaryan | V-pol | Below Horizon | Normal |
| 2) CR-reflected | Geo-synchrotron | H-pol | Below Horizon | Inverted |
| 3) CR-direct | Geo-synchrotron | H-pol | Above Horizon | Normal |
| 4) Tau Neutrino? | Geo-synchrotron | H-pol | Below Horizon | Normal |


Explanation within standard model

- Shoemaker et al. The Antarctic Subsurface as a possible explanation,
[arXiv:1905.02846](https://arxiv.org/abs/1905.02846)
- Sub-surface reflection  no phase inversion
- Could be tested by radar surveying the Antarctic region

Another explanation within standard model

- Coherent transition radiation (CTR) Geomagnetically induced current

From downward going cosmic ray



K. D. de Vries and S. Prohira, PRL 123 (2019) 091102

Downward or Upward

Optical Cherenkov instrument

EUSO-SPB2 (**balloon-based**) next run

POEMMA (**sattelite based**) proposed

Cumming et al., [arXiv:1910.00992](https://arxiv.org/abs/1910.00992)

Beyond SM explanations

For a review, see Anchordoqui et al, “The pros and cons of beyond standard model interpretations of ANITA events, [arXiv:1907.06308](https://arxiv.org/abs/1907.06308)


Three classes of models

- 1) Messenger lives inside the Earth
- 2) Originate in the neutrino-nucleon collisions inside the Earth
- 3) Messenger comes from cosmological distances.


Some nice ideas

- I. Esteban et al, Reflection of radio pulse produced by axion-photon conversion in ionosphere, [arXiv:1905.10372](#);
- Cherry and Shoemaker, A sterile neutrino origin for the upward directed cosmic ray shower detected by ANITA, [PRD99 \(2019\) 063016](#)

$$m_4 \leq 2 \text{ MeV} \times (.01 / \sin^2 2\theta_{\tau 4})^{1/6}$$



Our scenario



Very high energy coming from cosmos

A. Esmaili and YF, arXiv:1909.07995

Probability of producing tau neutrino or tau

γ is the inverse of the mean free path of N_1 (as well as that of N_2): $\gamma(x) = n_e(x)\sigma$.
mean free path of ν_τ , $\tau_\nu = (\sigma_{SM} \rho/m_p)^{-1}$

Γ_{tot} The total decay rate of N_2 including the time dilation

$$P = \text{Min}[1, \Gamma_{tot}\tau_\nu](2B) \int_0^L e^{-\int_0^x \gamma(y)dy} e^{-\int_x^L (\gamma(z)+\Gamma_{tot})dz} \gamma dx$$


Branching ratio Survival probability of N_1 Interaction probability

For constant density


$$P \simeq \text{Min}[1, \Gamma_{tot}\tau_\nu] \frac{\gamma}{\Gamma_{tot}} \left[e^{-L\gamma} - e^{-L(\gamma+\Gamma_{tot})} \right]$$

P is maximal for $\Gamma_{tot} \sim 1/\tau_\nu \sim (500 \text{ km})^{-1}$ and $\gamma \sim 1/L \sim (5000 \text{ km})^{-1}$.

Energy of final neutrinos





↓
Electrons in the Earth
Very high energy coming from cosmos



$$E_\nu \sim \frac{M_2 - M_1}{M_1} \frac{E_{N_1}}{4}$$

Scattering and decay



New gauge interaction

$L_e - L_\tau$ with a gauge coupling of $g_{e-\tau}$

$$g_N Z'_\mu \bar{N}_2 \sigma^\mu N_1 + \text{H.c.}$$

$$\sigma(N_1 + e \rightarrow N_2 + e) = \frac{g_N^2 g_{e-\tau}^2}{8\pi} s \int_{-1}^1 \frac{4 + (1 + \cos\theta)^2}{(s(1 - \cos\theta)/2 + m_{Z'}^2)^2} d\cos\theta$$

$$s \gg m_{Z'}^2 \quad \longrightarrow \quad \sigma \simeq (2g_N^2 g_{e-\tau}^2 / \pi m_{Z'}^2)$$

LEP bounds

$$e^- e^+ \rightarrow \gamma Z'$$

$$\frac{g_{e-\tau}}{m_{Z'}} < 2.0 \times 10^{-4} \text{GeV}^{-1} \quad \text{for} \quad 200 \text{ GeV} < m_{Z'}$$

$$\frac{g_{e-\tau}}{m_{Z'}} < 6.9 \times 10^{-4} \text{GeV}^{-1} \quad \text{for} \quad 100 \text{ GeV} < m_{Z'} < 200 \text{ GeV}.$$

Freitas and Westhoff, JHEP 1410 (2014) 116

Saturating LEP bounds

$$g_{e-\tau} = 6.9 \times 10^{-2} (m_{Z'}/100 \text{ GeV})$$



$$\sigma(e + N_1 \rightarrow e + N_2) = \sigma(e + N_2 \rightarrow e + N_1) = 10^{-34} g_N^2 \text{ cm}^2$$

$$\boxed{\gamma^{-1} = (n_e \sigma)^{-1} = 8 \times 10^4 \text{ km}/g_N^2}$$

Decay rate of N_2

$$\Gamma_{tot} = \frac{g_N^2 g_{e-\tau}^2}{10\pi^3} \frac{(M_2 - M_1)^5}{m_{Z'}^4} \left(\frac{M_2}{E_{N_2}} \right)$$

For given g_N and $\frac{g_{e-\tau}}{m_{Z'}}$, Γ_{tot} can be fixed by tuning $(M_2 - M_1)$.

The stability of N_1

Z_2 symmetry

$$N_1 \rightarrow -N_1 \quad N_2 \rightarrow -N_2 \quad SM \rightarrow SM$$

Lightest one is stable and travels long distances

N_1 in early universe

$$\langle \sigma(N_1 \bar{N}_2 \rightarrow \text{lepton pairs}) v \rangle \sim 3 \frac{g_N^2 g_{e-\tau}^2 M_1^2}{\pi m_{Z'}^4} = 1.6 \times 10^{-35} \left(\frac{M_1/m_{Z'}}{0.1} \right)^2 \text{cm}^2 \gtrsim 1 \text{ pb}$$

A proper dark matter component

How to obtain

$$g_N Z'_\mu \bar{N}_2 \sigma^\mu N_1 + \text{H.c.}$$

Within

$$L_e - L_\tau$$

The $L_e - L_\tau$ model

- Two Weyl fermions with opposite U(1) charge

$$\psi_1 = (N_1 + N_2)/\sqrt{2} \text{ and } \psi_2 = (N_1 - N_2)/\sqrt{2}.$$

Anomaly cancelation

$$g_N(\bar{\psi}_1 \sigma^\mu \psi_1 - \bar{\psi}_2 \sigma^\mu \psi_2) Z'_\mu = g_N(\bar{N}_1 \sigma^\mu N_2 + \bar{N}_2 \sigma^\mu N_1) Z'_\mu$$

Mass terms

$$m(\psi_1^T c \psi_2 + \psi_2^T c \psi_1) = m(N_1^T c N_1 - N_2^T c N_2)$$

$$c = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

$\psi_1 \leftrightarrow \psi_2$ and $\Phi \leftrightarrow \Phi^*$ (and $Z' \leftrightarrow -Z'$ and $e \leftrightarrow \tau$)

$$Y\langle\Phi\rangle(N_1^T c N_1 + N_2^T c N_2)$$

$$M_1 = |m + Y\langle\Phi\rangle| \quad \text{and} \quad M_2 = |m - Y\langle\Phi\rangle|$$

Quasi-degeneracy

$$M_1 = |m + Y\langle\Phi\rangle| \quad \text{and} \quad M_2 = |m - Y\langle\Phi\rangle|$$

It is natural to have

$$m \ll |Y\langle\Phi\rangle| \quad \text{Or} \quad m \gg |Y\langle\Phi\rangle|$$

Other possibilities to gauge

$B - 3L_\tau$

some other anomaly free combination of B , L_τ , L_μ and L_e

Maybe light Z'

Typical values giving rise to large P

$$\frac{g_{e-\tau}}{m_{Z'}} \sim 7 \times 10^{-4} \text{ GeV}^{-1}$$

$$g_N \sim 3$$

$$M_2 \simeq M_1 \sim 10 \text{ GeV}$$

$$M_2 - M_1 \sim \text{GeV} \frac{10 \text{ GeV}}{M_2}$$



$$\Gamma_{tot} \sim \tau_\nu^{-1} \sim (500 \text{ km})^{-1}$$

$$\sigma \sim L^{-1}$$

Prediction for the ILC

$$e^- e^+ \rightarrow \gamma Z'$$

$$(P_{e^-} + P_{e^+} - P_\gamma)^2 = m_{Z'}^2$$


$$Z' \rightarrow N_1 \bar{N}_2, N_2 \bar{N}_1$$
$$M_2 - M_1 > 2m_\tau$$

In 1/3 of cases:

$$N_2 \rightarrow N_1 \nu_e \bar{\nu}_e, N_1 \nu_\tau \bar{\nu}_\tau$$



In 1/3 of cases:

$$N_2 \rightarrow N_1 e^- e^+$$

In 1/3 of cases:

$$N_2 \rightarrow N_1 \tau \bar{\tau}$$

Displaced vertex

$$e^- e^+ \rightarrow \gamma Z' \quad Z' \rightarrow N_1 \bar{N}_2, N_2 \bar{N}_1 \quad N_2 \rightarrow N_1 e^- e^+$$

The displacement is given by $(M_2 - M_1)^5 g_N^2$

$$E_{e^-} \sim E_{e^+} \sim \frac{s + m_{Z'}^2}{8\sqrt{s}} \left(1 - \frac{M_1}{M_2}\right)$$

$\sqrt{s} = 500$ GeV and $1 - M_1/M_2 \sim 0.1$



$$E_{e^-} \sim E_{e^+} \sim 6 \text{ GeV}$$

Non-Standard Interactions

$$2\sqrt{2}\epsilon^e G_F (\bar{\nu}_e \gamma^\mu P_L \nu_e - \bar{\nu}_\tau \gamma^\mu P_L \nu_\tau)(\bar{e} \gamma_\mu e)$$

$$\epsilon^e \simeq 0.01$$

5 times improvement is required.

Agarwalla, Borexino collaboration, 1905.03512; Choubey and Ohlsson, 1410.0410

Possible mechanism for N_1 production

- Very heavy dark matter


$$\Phi(\psi_1^T c \psi_2 + \psi_2^T c \psi_1) = \Phi(N_1^T c N_1 - N_2^T c N_2)$$


Similar fluxes for N_1 and N_2 . $N_2 \rightarrow N_1 \nu_e \bar{\nu}_e, N_1 \nu_\tau \bar{\nu}_\tau, N_1 \tau \bar{\tau}$

| Benchmark \ Parameter | g_N | $g_{e-\tau}$ | $m_{Z'}$ [GeV] | M_2 [GeV] | M_1 [GeV] | τ_{N_2} [km] | γ^{-1} [km] |
|-----------------------|-------|----------------------|----------------|-------------|-------------|-------------------|--------------------|
| (A) | 3.0 | 6.9×10^{-2} | 100 | 10 | 9.3 | 42 | 7,730 |
| (B) | 2.0 | 6.9×10^{-2} | 100 | 10 | 9.3 | 94 | 17,390 |
| (C) | 3.0 | 1.4×10^{-1} | 200 | 10 | 9.3 | 168 | 7,730 |
| (D) | 3.0 | 6.9×10^{-2} | 100 | 10 | 8.7 | 2 | 7,730 |

$$E_\nu \sim E_{N_1} (M_2 - M_1) / (4M_1)$$

$$E_{N_1} = (3.2, 32, 320) \text{ EeV}$$







$$\theta_z \simeq (100^\circ - 110^\circ).$$

$$P \simeq 0.012$$

Exposure and Flux

- ANITA exposure: $2.7 \text{ km}^2 \text{ yr sr}$
- Incident N_1 flux $\simeq 40 \text{ km}^{-2} \text{ yr}^{-1} \text{ sr}^{-1}$






Cherry and Shoemaker, PRD99 (2019) 63016


ICECUBE data taking ~ 8 years

ANITA data taking ~ 1 month



Cherry and Shoemaker, PRD99 (2019) 63016

| Benchmark | Parameter | g_N | $g_{e-\tau}$ | $m_{Z'} \text{ [GeV]}$ | $M_2 \text{ [GeV]}$ | $M_1 \text{ [GeV]}$ | $\tau_{N_2} \text{ [km]}$ | $\gamma^{-1} \text{ [km]}$ |
|-----------|-----------|-------|----------------------|------------------------|---------------------|---------------------|---------------------------|----------------------------|
| (A) | | 3.0 | 6.9×10^{-2} | 100 | 10 | 9.3 | 42 | 7,730 |



$$E_\nu \simeq E_{N_1} (M_2 - M_1) / (4M_1).$$

Summary and conclusions

- 2 anomalous **ANITA** events (tested by radar surveying of Antarctic region and by **POEMMA** and **EUSO-SPB2**)
- Our scenario: $N_1 + e \rightarrow N_2 + e$ $N_2 \rightarrow N_1 \nu_\tau \bar{\nu}_\tau$
- Our model: $L_e - L_\tau$ with Z' to be discovered by ILC
- Induce NSI to be tested by solar neutrinos or by PINGU
- Bonus: A candidate for dark matter

Some possible signature at LHC

$$\lambda_{H\Phi} |H|^2 |\Phi|^2$$

$$H \rightarrow \Phi \bar{\Phi} \rightarrow N_1 \bar{N}_1 N_2 \bar{N}_2 \rightarrow 2N_1 2\bar{N}_1 l \bar{l} l' \bar{l}'$$

$$\lambda_{H\Phi} \ll m_\tau / \langle H \rangle$$