

# Anisotropic flow measurements with MPD at NICA\*

A. TARANENKO, P.PARFENOV, E. KASHIRIN, O. GOLOSOV, A.  
DEMANOV, A. TRUTTSE, E. VOLODIKHIN

National Research Nuclear University (Moscow Engineering Physics Institute)  
Moscow, Russia

The anisotropic transverse flow is one of the key observables to study the properties of dense matter created in heavy-ion collisions. The performance of MPD detector for directed and elliptic flow measurements is studied with Monte-Carlo simulations of gold ions at NICA energies  $\sqrt{s_{NN}} = 4 - 11$  GeV using different heavy-ion event generators.

PACS numbers: PACS numbers come here

## 1. Introduction

The Multi-Purpose Detector (MPD) at NICA collider has a substantial discovery potential concerning the exploration of the QCD phase diagram in the region of high net-baryon densities and moderate temperatures [1]. The anisotropic collective flow, as manifested by the anisotropic emission of particles in the plane transverse to the beam direction, is one of the important observable sensitive to the transport properties of the strongly interacting matter: the equation of state (EOS), the specific shear and bulk viscosity [2]. The azimuthal anisotropy of produced particles can be quantified by the Fourier coefficients  $v_n$  in the expansion of the particles azimuthal distribution as:  $dN/d\phi \propto 1 + \sum_{n=1}^{\infty} 2v_n \cos(n(\phi - \Psi_n))$ , where  $n$  is the order of the harmonic,  $\phi$  is the azimuthal angle of particles of a given type, and  $\Psi_n$  is the azimuthal angle of the  $n$ th-order event plane. In this work, we briefly review the available experimental results for the collision energy dependence of directed ( $v_1$ ) and elliptic ( $v_2$ ) flow and discuss the anticipated performance of MPD detector for flow measurements at NICA energies.

---

\* Presented at NICA days 2019 and IVth MPD Collaboration meeting in Warsaw

## 2. Collision energy dependence of directed and elliptic flow

The slope of the rapidity dependence  $dv_1/dy$  near mid-rapidity ( $y \sim 0$ ) is a convenient way to characterize the overall magnitude of directed flow signal [3]. A minimum in  $dv_1/dy$  for baryons could be related to the softening of equation-of-state due to the first order phase transition between hadronic matter and sQGP [3]. The recent results from the STAR experiment at RHIC ( $\sqrt{s_{NN}} = 7.7 - 200$  GeV) seem to support this prediction, both protons and  $\Lambda$  hyperons  $dv_1/dy$  show a minimum around  $\sqrt{s_{NN}}=10-20$  GeV [4], see left panel of Fig. 1. The green curves show the approximate upper and lower bounds for centrality dependence of  $dv_1/dy$  for protons.




Fig. 1. Left: Excitation function of  $dv_1/dy$  for protons from Au+Au collisions based on results from STAR [4]. Excitation function of  $v_2(p_T)$  of protons from mid-central Au+Au collisions at energies from  $\sqrt{s_{NN}} = 2$  to 62.4 GeV [2].

In general, the assumption of purely hadronic physics is disfavored by the comparison with predictions from current state-of-the-art models [3]. However, all current models are not able to reproduce the  $\sqrt{s_{NN}}$  dependence of baryon  $dv_1/dy$ -slope reported by STAR. Thus, further progress in the area of model calculations and high-statistics differential measurements of  $v_1$  is needed.

The right panel of Fig. 1 shows the excitation function for differential elliptic flow  $v_2(p_T)$  of protons from mid-central Au+Au collisions at  $\sqrt{s_{NN}} = 2$  to 62.4 GeV [2]. Figure shows that the  $v_2(p_T)$  for protons changes relatively little as a function of beam energy in the range  $\sqrt{s_{NN}} = 11.5 - 62.4$  GeV [5] and this may result from the interplay of the hydrodynamic and hadronic transport phase [6]. In the energy range  $\sqrt{s_{NN}} = 11 - 2$  GeV, the passage time  $t_{pass}$  increases from 2 fm/c to 16 fm/c and the shadowing effects by the spectator matter start to play an important role for the generation of elliptic flow [7].

### 3. Performance for flow measurements with MPD detector

The anticipated performance of the MPD detector for the directed flow measurements of charged pions, kaons and protons is demonstrated in the Figure 2. A sample of 5M minimum-bias Au+Au collisions at  $\sqrt{s_{NN}} = 5$  (left panel) and 11 GeV (right panel) simulated with UrQMD event generator [6] was used for the analysis. The MPD detector response was simulated using GEANT4 toolkit and the resulting signals from the detector subsystems were used as input information for the full reconstruction procedure, which includes the realistic particle identification in TOF and TPC detectors ( $|\eta| < 1.5$ ) of MPD. The first order event plane was reconstructed using the energy deposition of particles detected in the forward hadronic calorimeters (FHCaL), located at ( $2 < |\eta| < 5$ ).




Fig. 2. Directed flow  $v_1$  for charged pions, kaons and protons as a function of rapidity  $y$  for Au+Au collisions at  $\sqrt{s_{NN}} = 5$  (left) and 11 GeV (right). The results from the UrQMD model are marked as true, and the ones from the full MPD reconstruction procedure are marked as reco.

Figure 3 shows the elliptic flow signal  $v_2$  of protons from 10-40% mid-central Au+Au collisions at  $\sqrt{s_{NN}} = 7.7$  GeV obtained using the event plane analysis of 5M events from the current state of the art transport models: UrQMD [8], JAM [7], PHSD [9], AMPT [10] and hybrid vHLL+UrQMD model[6]. A second-order event plane angle ( $\Psi_2$ ) is reconstructed from the charged particles reconstructed in TPC at mid-rapidity ( $|\eta| < 1.0$ ). For comparison we plotted the published results from STAR experiment [5] obtained by the same method of analysis. Pure hadronic transport system (as described by the UrQMD or JAM models) does not appear to explain the relatively large flow of the particles at NICA energies [6]. The hybrid models: vHLL+UrQMD [6] and string melting version of AMPT [10] provides

much better description of the data.




Fig. 3. The comparison of  $v_2(p_T)$  of protons from 10-40% mid-central Au+Au collisions from event plane analysis using events UrQMD, PHSD, JAM, AMPT and hybrid vHLE+UrQMD models with published results from STAR [5].

*In summary*, the high-statistics differential measurements of  $v_n$  anticipated from the MPD experiment expected to provide the valuable information about the properties of strongly interacting matter at NICA energies.

#### 4. Acknowledgements

The reported study was funded by RFBR according to the research project No 18-02-00086.

#### REFERENCES

- [1] V. D. Kekelidze, Phys. Part. Nucl. **49** (2018) no.4, 457.
- [2] A. Taranenko, EPJ Web Conf. **204**, 03009 (2019).
- [3] S. Singha et al. Adv. High Energy Phys. **2016** (2016) 2836989
- [4] L. Adamczyk *et al.* [STAR], Phys. Rev. Lett. **120** (2018) no.6, 062301
- [5] L. Adamczyk *et al.* [STAR], Phys. Rev. C **93** (2016) no.1, 014907
- [6] I. A. Karpenko, P. Huovinen, H. Petersen and M. Bleicher, Phys. Rev. C **91**, no. 6, 064901 (2015)
- [7] C. Zhang, J. Chen, X. Luo, F. Liu and Y. Nara, Phys. Rev. C **97** (2018) no.6, 064913
- [8] S. A. Bass *et al.*, Prog. Part. Nucl. Phys. **41**, 255 (1998)
- [9] W. Cassing and E. L. Bratkovskaya, Phys. Rev. C **78**, 034919 (2008)
- [10] Z. W. Lin, C. M. Ko, B. A. Li, B. Zhang and S. Pal, Phys. Rev. C **72**, 064901 (2005)