

Partikeldagarna 2019

Wednesday, 2 October 2019 - Thursday, 3 October 2019

Linköping University

Book of Abstracts

Contents

| | |
|--|---|
| CP violation 53 | 1 |
| High-energy gamma astronomy 54 | 1 |
| Prospects and Progress of the Light Dark Matter eXperiment (LDMX) – 15+5 14 | 1 |
| Physics opportunities with the CERN SPS Beam Dump Facility (15+5) 17 | 1 |
| The HIBEAM/NNBAR Experiment at the ESS (15+5) 23 | 1 |
| Observing the sky continuously at extreme photon energies with ALTO/COMET (11+3) 32 | 1 |
| The helicity-flow method (11+3) 12 | 2 |
| Blazars as astroparticle physics laboratories: the case of KUV00311-1938 (11+3) 35 | 2 |
| Finite temperature effects on particle decays (11+3) 26 | 3 |
| Anomaly-free model building: algebraic geometry and the Froggatt-Nielsen mechanism (11+3) 21 | 3 |
| Non-Standard Decays of Vector-Like Quarks 16 | 3 |
| Search for a Dark Jet Resonances Using Substructure with ATLAS 11 | 4 |
| Synergies between different disciplines for the European Particle Physics Strategy Update 57 | 4 |
| Impact of different compilers and build types on Geant4 simulation execution time 38 | 5 |
| IceCube search for neutrinos from precursors and afterglows of gamma ray bursts 15 | 5 |
| Searches for dark matter mediators decaying to jets in ATLAS 27 | 6 |
| Unfolding of multivariate tools and statistical analysis for Higgs boson pair production searches in the ATLAS detector at the Large Hadron Collider 20 | 6 |
| Exploring selections across channels in Dark Matter searches with top quarks at the ATLAS experiment of the LHC 19 | 6 |
| Displaced vertices and material map in ATLAS 56 | 7 |
| CERN Council including update of the European strategy for particle physics (15+5) 41 | 7 |
| IPPOG (12+3) 43 | 7 |

| | |
|---|----|
| IUPAP (12+3) 45 | 8 |
| Election of a new board (15) 47 | 8 |
| (R)ECFA (12+3) 42 | 8 |
| High-school project with open data (8+2) 44 | 8 |
| ACCU (12+3) 46 | 8 |
| VR+RFI (12+3) 40 | 8 |
| Coffee break 3 | 8 |
| Top-related SM and BSM physics in ATLAS (15+5) 51 | 9 |
| ATLAS upgrade (15+5) 49 | 9 |
| ATLAS detector operation, data-quality and physics object performance (15+5) 48 | 9 |
| Higgs physics in ATLAS (15+5) 50 | 9 |
| Other BSM physics in ATLAS (15+5) 52 | 9 |
| ALICE (15+5) 13 | 9 |
| MAX IV Status (15+5) 55 | 10 |
| Direct detection of low-mass dark matter with strong matter interactions – 15+5 31 | 10 |
| Local dark matter density from the Milky Way’s rotation curve using Gaia DR2 data – 15+5 9 | 10 |
| The Hyper-Kamiokande Experiment – 15+5 33 | 10 |
| ESSnuSB: physics reach and instrumentation development – 15+5 36 | 10 |
| Measuring the local dark matter density at direct detection experiments – 15+5 30 | 11 |
| Update on the Neutrino Observatory in ice with radio technique – 15+5 37 | 11 |
| Results from IceCube and the path to the Upgrade and Gen2 – 15+5 34 | 11 |
| Registration with coffee 1 | 11 |
| Coffee break 5 | 11 |
| Thu morning session 2 6 | 11 |
| Lunch 7 | 12 |
| Short walk 8 | 12 |
| Lunch 2 | 12 |
| Coffee break 4 | 12 |

Plenary talk 1 / 53

CP violation

Corresponding Author: yasmine.sara.amhis@cern.ch

Plenary talk 2 / 54

High-energy gamma astronomy

Corresponding Author: stefano.vercellone@inaf.it

Thu afternoon session 2 / 14

Prospects and Progress of the Light Dark Matter eXperiment (LDMX) – 15+5

Author: Ruth Pottgen¹

Co-authors: Torsten Akesson¹; Lene Kristian Bryngemark¹

¹ *Lund University (SE)*

Corresponding Authors: torsten.akesson@cern.ch, ruth.poettgen@cern.ch, lene.bryngemark@cern.ch

Thu afternoon session 2 / 17

Physics opportunities with the CERN SPS Beam Dump Facility (15+5)

Author: Richard Jacobsson¹

¹ *CERN*

Corresponding Author: richard.jacobsson@cern.ch

Thu afternoon session 2 / 23

The HIBEAM/NNBAR Experiment at the ESS (15+5)

Author: David Anthony Milstead¹

¹ *Stockholm University (SE)*

Corresponding Author: milstead@fysik.su.se

Thu morning session 1 / 32

Observing the sky continuously at extreme photon energies with ALTO/COMET (11+3)

Author: Mohanraj Senniappan¹

Co-authors: Satyendra Thoudam²; Yvonne Becherini¹; Michael Punch ; Jean-Pierre Ernenwein³; Tomas Bylund¹

¹ *Linnaeus University*

² *Khalifa University*

³ *Aix-Marseille Universite*

Corresponding Authors: yvonne.becherini@lnu.se, satyendra.thoudam@ku.ac.ae, tomas.bylund@lnu.se, mpunch.in2p3@gmail.com, mohanraj.senniappan@lnu.se, jeanpierre.ernenwein@lnu.se

ALTO is a future very-high-energy (VHE) gamma-ray observatory, which is currently in the prototyping phase. The proposed design of the array consists of more than a thousand Water Cherenkov Detectors (WCDs) each coupled with a liquid scintillator. The observatory will be installed at an altitude of 4 to 5 km above the sea level in the Southern Hemisphere. WCDs sample the secondary particles in the extensive air showers generated by VHE gamma-rays and cosmic rays while the scintillators help in tagging the presence of muons. Preliminary studies using the simulation shows that the scintillators help in improving the signal over background discrimination by 15 to 30% depending on the energy range. The first phase of the prototype, comprised of two WCDs and scintillators, has been taking data since February 2019. In the poster, I will present the simulation performance and some results from the prototype activities. Finally I will also present the future steps of the project.

Thu morning session 1 / 12

The helicity-flow method (11+3)

Author: Andrew Lifson¹

Co-authors: Christian Reuschle¹; Malin Sjordahl¹

¹ *Lund University*

Corresponding Authors: malin.sjordahl@thep.lu.se, andrew.lifson@thep.lu.se, christian.reuschle@thep.lu.se

It is well-known that the spinor-helicity method can significantly simplify the calculations of both Feynman diagrams and scattering amplitudes. In this work, we attempt to further simplify the Feynman-diagram calculation by converting the spinor-helicity method into a flow method, analogous to allowing a one-line journey from Feynman diagram to inner products. The cases of massless QED and QCD will be discussed.

Thu morning session 1 / 35

Blazars as astroparticle physics laboratories: the case of KUV00311-1938 (11+3)

Author: Tomas Bylund^{None}

Blazars are very bright gamma ray sources in the sky that are visible out to very large redshifts. Powered by material falling into a supermassive black hole, blazars have jets that can emit gamma rays up to TeV energies. As these TeV photons propagate through space they are expected to encounter ambient magnetic fields and extragalactic background light (EBL). The presence of magnetic fields

can convert them to Axion like particles (ALPs), while the presence of the EBL absorbs the initial spectrum via pair production. We studied the multi-wavelength behaviour of KUV 00311-1938, a blazar with an unknown redshift, finding that a simple EBL model can explain the observed emission. We subsequently used the EBL model to constrain the redshift of the source.

Thu morning session 1 / 26

Finite temperature effects on particle decays (11+3)

Author: Torbjörn Lundberg¹

Co-author: Roman Pasechnik²

¹ *Lund University*

² *Lund university*

Corresponding Authors: roman.pasechnik@thep.lu.se, torbjorn.lundberg@thep.lu.se

At high densities and temperatures the standard quantum field theoretical approach to particle physics should be modified. Temperature enters explicitly in observables, for instance in decay rates, and under certain conditions expected results deviate significantly from the case of zero-temperature. I have put together a collection of thermal decay rates covering scalars, pseudo scalars and fermions consequently expanding the existing literature. I aim to lay out the procedure of thermal calculations detailing the explicit appearance of temperature in the two-point correlation function. I also highlight the interpretation of the thermal decay rate in comparison with the zero-temperature dito.

Thu morning session 1 / 21

Anomaly-free model building: algebraic geometry and the Froggatt-Nielsen mechanism (11+3)

Authors: Felix Tellander¹; Johan Rathsman^{None}

¹ *Lund University*

Corresponding Authors: johan.rathsman@thep.lu.se, felix.tellander.355@student.lu.se

We present methods to find anomaly-free gauged U(1) Froggatt-Nielsen type models using results from real algebraic geometry. These methods should be of general interest for model building beyond the Standard Model when rational charges are required. We consider models with a gauged U(1) flavour symmetry with one flavon and provide several model examples based on different physical assumptions. Necessary conditions for these models to be free from gauge anomalies are derived and we show that the field content of the Standard Model is not sufficient and thus additional fields are needed. Two such extensions are considered; two Higgs doublets and right-handed neutrinos providing Dirac masses. With these extensions, the fermion masses and mixings in the Standard Model, including neutrinos, can in great part be explained.

Moreover, we show that the UV-behaviour of these models are in general plagued by Landau poles. Two different UV-completions are considered; through vector-like fermions and through Higgs doublets. In the fermion completion, the gauge couplings are in general plagued with Landau poles while in a scalar completion this may be avoided, but instead the quartic couplings generally blow up. Thus, the generic case is that neither completion works, but the scalar completion might be saved by appropriate choice of parameters in the scalar potential. This conclusion does not change if we allow U(1) to be anomalous or global.

Thu morning session 2 / 16

Non-Standard Decays of Vector-Like Quarks

Author: Thomas Mathisen¹

¹ *Uppsala University (SE)*

Corresponding Author: thomas.mathisen@cern.ch

With strong constraints on the mass of vector-like quarks (VLQ) from the top partner (T) decay $T \rightarrow SM$, it is necessary to consider non-standard decays of such partners. This talk considers models where the VLQ decays to a BSM (pseudo)scalar (S) and a top-quark. The scalar is assumed to be fermiophobic, and dominantly decays into two SM bosons. With dedicated analyses, we realistically quantify the sensitivity of the LHC to both the T and S masses, assuming both current and foreseen luminosities.

Thu morning session 2 / 11

Search for a Dark Jet Resonances Using Substructure with ATLAS

Author: Eva Brottmann Hansen¹

¹ *Lund University (SE)*

Corresponding Author: eva.hansen@cern.ch

The possibility of dark matter being a composite particle, such as a hadron under a gauge group, is becoming increasingly appealing as the non-excluded part of parameter space for the WIMP paradigm is shrinking. In this talk I will present an ongoing search for a resonance originating from a QCD-like dark sector, that decays to two dark partons which then hadronise to form "dark jets". The search targets a selection of model parameters where a fraction of the dark mesons are stable and the rest decay promptly to visible particles. The signature is a pair of jets with potentially different substructure than normal QCD-jets. I will introduce the model and the initial signal studies that have been made and then go into more detail with the mass-decorrelated tagger implemented for the signal region definition.

Thu morning session 2 / 57

Synergies between different disciplines for the European Particle Physics Strategy Update

Author: Caterina Doglioni¹

¹ *Lund University (SE)*

Corresponding Author: caterina.doglioni@cern.ch

One overarching objective of science is to further our understanding of the universe, from its early stages to its current state and future evolution. This depends on gaining insight on the universe's most macroscopic components, for example galaxies and stars, as well as describing its smallest components, namely elementary particles and nuclei and their interactions. It is clear that this endeavor requires combined expertise from the fields of astroparticle physics, particle physics and nuclear physics. Pursuing common scientific drivers also require mastering challenges related to

instrumentation (e.g. beams and detectors), data acquisition, selection and analysis, and making data and results available to the broader science communities. Joint work and recognition of these “foundational” topics will help all communities grow towards their individual and common scientific goals. This contribution presents the work that various communities and experiments are doing to solve one of the many common challenges faced by particle physics and astrophysics: the necessity of dealing with large, sometimes heterogeneous datasets and derive insight from them in short periods of time.

Thu morning session 2 / 38

Impact of different compilers and build types on Geant4 simulation execution time

Author: Einar Alfred Elen¹

¹ *Lund University (SE)*

Corresponding Author: einar.alfred.elen@cern.ch

Experimental observations and advanced computer simulations in High Energy Physics (HEP) paved way for the recent discoveries at the Large Hadron Collider (LHC) at CERN. Currently, Monte Carlo simulations account for a very significant amount of computational resources of the Worldwide LHC Computing Grid (WLCG).

In looking at the recent trends in modern computer architectures we see a significant deficit in expected growth in performance. Coupled with the increasing compute demand for High Luminosity (HL-LHC) run, it becomes vital to address this shortfall with more efficient simulation.

The simulation software for particle tracking algorithms of the LHC experiments predominantly relies on the Geant4 simulation toolkit. The Geant4 framework can be built either as a dynamic or a static library, the former being the more widely used approach. This study focuses on evaluating the impact of both having libraries statically vs dynamically linked and compiler optimization levels, on the simulation software’s execution time.

Multiple versions of the more widely used versions of compilers for UNIX-like systems have been used for these investigations. Both compiler optimization levels (e.g. O3, O2 on GCC) and link-time optimization (LTO) have been studied. Initial results indicate that significant execution time reductions can be obtained by switching from static to dynamic linking.

Thu morning session 2 / 15

IceCube search for neutrinos from precursors and afterglows of gamma ray bursts

Authors: Kunal Deoskar¹; Chad Finley^{None}

¹ *Stockholm University*

Corresponding Authors: cfinley@fysik.su.se, kunal.deoskar@fysik.su.se

The sources and accelerating mechanisms of cosmic rays, with energies as high as 10^{20} eV, are not completely understood. Gamma ray bursts (GRBs) have long been considered as promising source candidates, yet so far don’t show evidence for a correlated neutrino signal to prove this hypothesis. Previous analyses by IceCube have searched for neutrino in coincidence with the prompt phase of the GRBs, typically lasting for 100s or less. Here I will describe a search for neutrino correlations before and after this prompt phase using an extended time window. Presently I am calculating the spatial and temporal coincidence of a list of GRBs with the neutrino detections made by IceCube

and comparing the results for different precursor and afterglow models hypothesised for short and long duration GRBs. These searches build on a similar methodology as was used for the detection of longer time-scale transients such as blazar flares, applying it for the first time to short transients like GRBs.

Thu morning session 2 / 27

Searches for dark matter mediators decaying to jets in ATLAS

Author: Eric Edward Corrigan¹

¹ *Lund University (SE)*

Corresponding Author: eric.edward.corrigan@cern.ch

An important part of the ATLAS dark matter (DM) search programme is comprised of searches for new resonances (dark matter mediators) decaying to hadronic final states. This talk will give an overview of one such analysis: the recently-published search for low-mass dark-matter mediators decaying to jets, with an associated high-pT photon. This search triggers on an associated photon in order to significantly extend the sensitive region of ATLAS for dark matter mediators and generic resonances. Also covered is the reinterpretation of the resulting limits in the case of non-discovery.

Thu morning session 2 / 20

Unfolding of multivariate tools and statistical analysis for Higgs boson pair production searches in the ATLAS detector at the Large Hadron Collider

Author: Christina Dimitriadi¹

¹ *Uppsala University*

Corresponding Author: christinedimitriadi@gmail.com

Recently, searches for pair production of Higgs bosons in several final states have been carried out by the ATLAS experiment at the Large Hadron Collider (LHC). This study focuses on the search for non-resonant di-Higgs production decaying to a final state with two b -jets and two τ -leptons using 36.1 fb^{-1} of data recorded by the ATLAS detector. The analysis for this process has already been performed. Boosted decision trees (BDTs) are used in the analysis to improve the separation of the signal from background processes and several variables that provide good discrimination between signal and background are used as inputs to the BDT. This study aims to unfold the BDT of the analysis and optimize a cut-based analysis so that the gain from using the BDT can be estimated. Two variables, related to the invariant masses and angular distances of the Higgs boson decay products, are defined and the optimal cuts are found to be $X_{m_{\tau\tau}m_{bb}} > 1.8$ and $X_{\Delta R_{\tau\tau}\Delta R_{bb}} > 4.0$. Then, the upper limits on the SM HH production cross section are set when fitting m_{HH} with the cut-based analysis. An expected limit of 0.78 pb, 23 times the SM prediction is obtained when neglecting systematic uncertainties, compared to the limit of 15 times the SM as recomputed when using the BDT. Comparing the two results, the sensitivity is worsened by 50% when not using the BDT.

Thu morning session 2 / 19

Exploring selections across channels in Dark Matter searches with top quarks at the ATLAS experiment of the LHC

Author: Olga Sunneborn Gudnadottir¹

¹ *Uppsala University (SE)*

Corresponding Author: olga.sunneborn.gudnadottir@cern.ch

Current estimates put Dark Matter to 26.8% of the energy-matter content of the universe, but very little is known about it other than its gravitational interactions. Efforts to learn more about Dark Matter include searching for it at high energy particle colliders. The lack of information about the nature of Dark Matter makes this a complicated task, and many searches are performed in different channels, and considering different theoretical models. In my master thesis, I explore two such analyses, performed in the ATLAS collaboration using data from the ATLAS detector at the Large Hadron Collider at CERN: the tW +MET final state and the $t\bar{t}$ +MET final state. I have made a generation-level study of the overlap between the signal regions used, and come to the conclusion that there is some. I have also compared the models used in these analyses, the 2HDM+a and the simplified spin-0 pseudoscalar model. Given the simplifications made in my study, however, more sophisticated approaches should be used before anything conclusive can be said.

Thu morning session 2 / 56

Displaced vertices and material map in ATLAS

Author: Karl Filip Backman¹

¹ *Stockholm University (SE)*

Corresponding Author: filip.backman@cern.ch

A key observable for new physics at the Large Hadron Collider is the displaced vertex signature. Heavy and hitherto unseen particles can be produced and travel a short distance prior to decaying into Standard Model particles. A search for a such signature associated with a muon and charged particles from the ATLAS collaboration is shown in this poster. The search is based on a data sample of proton-proton collisions at a centre-of-mass energy of 13 TeV, corresponding to an integrated luminosity 136 fb⁻¹. The observed event yields are compatible with those expected from background processes. The results are presented as limits at 95% confidence level on model-independent cross sections for processes beyond the Standard Model, and interpreted as exclusion limits on scenarios with pair-production of long-lived top squarks that decay via a small R-parity-violating coupling to a quark and a muon. A key background to the search arises from interactions of particles from the proton-proton collision with detector and accelerator-related material. The poster also includes details of a study undertaken to estimate and minimise this background.

Wed afternoon session 1 / 41

CERN Council including update of the European strategy for particle physics (15+5)

Corresponding Author: ker@fysik.su.se

Wed afternoon session 1 / 43

IPPOG (12+3)

Corresponding Author: jonas.strandberg@cern.ch

Wed afternoon session 1 / 45

IUPAP (12+3)

Corresponding Author: johan.rathsman@thep.lu.se

Wed afternoon session 1 / 47

Election of a new board (15)

Corresponding Authors: sara.strandberg@cern.ch, johan.rathsman@thep.lu.se

Wed afternoon session 1 / 42

(R)ECFA (12+3)

Corresponding Author: milstead@fysik.su.se

Wed afternoon session 1 / 44

High-school project with open data (8+2)

Corresponding Author: elise.mariam@gmail.com

Wed afternoon session 1 / 46

ACCU (12+3)

Corresponding Author: sara.strandberg@cern.ch

Wed afternoon session 1 / 40

VR+RFI (12+3)

Corresponding Author: mathias.hamberg@vr.se

Wed afternoon session 2 / 3

Coffee break

Wed afternoon session 2 / 51

Top-related SM and BSM physics in ATLAS (15+5)

Corresponding Author: venugopal.ellajosyula@cern.ch

Wed afternoon session 2 / 49

ATLAS upgrade (15+5)

Corresponding Author: geoffrey.mullier@cern.ch

Wed afternoon session 2 / 48

ATLAS detector operation, data-quality and physics object performance (15+5)

Corresponding Author: laura.pereira.sanchez@cern.ch

Wed afternoon session 2 / 50

Higgs physics in ATLAS (15+5)

Corresponding Author: michael.edward.nelson@cern.ch

Wed afternoon session 2 / 52

Other BSM physics in ATLAS (15+5)

Corresponding Author: eva.hansen@cern.ch

Wed afternoon session 2 / 13

ALICE (15+5)

Authors: Alice Ohlson¹; Peter Christiansen¹; David Silvermyr¹

¹ Lund University (SE)

Corresponding Authors: peter.christiansen@hep.lu.se, david.silvermyr@gmail.com, alice.ohlson@cern.ch

Wed morning session / 55

MAX IV Status (15+5)

Corresponding Author: bmeirose.cern@gmail.com

Wed morning session / 31

Direct detection of low-mass dark matter with strong matter interactions – 15+5

Author: Timon Emken¹

¹ *Chalmers University of Technology*

Corresponding Author: emken@chalmers.se

Wed morning session / 9

Local dark matter density from the Milky Way's rotation curve using Gaia DR2 data – 15+5

Author: Pablo Fernández de Salas¹

¹ *Stockholm University*

Corresponding Author: pablo.fernandez@fysik.su.se

Wed morning session / 33

The Hyper-Kamiokande Experiment – 15+5

Corresponding Authors: eosullivan.queens@gmail.com, eosullivan@owl.phy.queensu.ca

Wed morning session / 36

ESSnuSB: physics reach and instrumentation development – 15+5

Author: Mattias Blennow¹

Co-author: Tord Johan Carl Ekelof²

¹ *KTH Royal Institute of Technology*

² *Uppsala University (SE)*

Corresponding Authors: tord.johan.carl.ekelof@cern.ch, emb@kth.se

Wed morning session / 30

Measuring the local dark matter density at direct detection experiments – 15+5

Authors: Riccardo Catena¹; Riccardo Catena¹

¹ *Chalmers University of Technology*

Corresponding Authors: riccardo.catena@chalmers.se, catena@chalmers.se

Wed morning session / 37

Update on the Neutrino Observatory in ice with radio technique – 15+5

Author: Allan Hallgren¹

¹ *Uppsala University*

Corresponding Author: allan.hallgren@physics.uu.se

Wed morning session / 34

Results from IceCube and the path to the Upgrade and Gen2 – 15+5

Author: Chad Finley^{None}

Corresponding Author: cfinley@fysik.su.se

1

Registration with coffee

5

Coffee break

6

Thu morning session 2

7

Lunch

8

Short walk

2

Lunch

4

Coffee break