

# CLIC sensitivity to measure $\sigma_{H\nu\nu} \times \text{BR} (H \rightarrow \gamma\gamma)$ at 3 TeV center-of-mass energy


ALPS2019, 22-27 April 2019

Goran Kacarevic,  
kacarevicgoran@vin.bg.ac.rs

# Compact Linear Collider - CLIC


- CLIC is a TeV scale  $e^+e^-$  linear collider
- Several energy stages: 380 GeV ( $1\text{ab}^{-1}$ ), 1.5 TeV ( $2.5\text{ab}^{-1}$ ) and 3 TeV ( $5\text{ab}^{-1}$ )
- Modular, upgradeable, site length 11 – 50 km

- All-silicon low mass vertex and tracking detectors
- High-granularity calorimeters
- 4T superconducting solenoid
- Muon detectors within return yoke
- **Optimized for Particle Flow reconstruction**



# Higgs Production at CLIC

- CLIC as a Higgs factory:
  - $1.6 \cdot 10^5$  Higgs at 380 GeV
  - $\sim 10^6$  Higgs per higher energy stage
- Top threshold scan, model-independent Higgs measurements in the Higgsstrahlung
- Excellent sensitivity at higher energies to BSM physics, top and Higgs measurements
- WW-fusion dominates Higgs production at the energies above 500 GeV
- Production cross-section can be increased by a factor of 1.48, by electron polarization -80%:+80%, where runtime will be 4:1 with -80% electron polarisation




Production cross section as a function of centre-of-mass energy for the main Higgs boson production processes at an  $e^+e^-$  collider

$$H \rightarrow \gamma\gamma$$


- One of the channels at LHC where Higgs boson was confirmed (discovered)
- High efficiency of photon reconstruction, excellent ECAL performance
- Higgs boson is coupled to photons via loop diagram and thus it is sensitive to eventual contribution of BSM physics through **precision measurements**

- WW fusion dominant H production process at 3 TeV
- $\sigma(H_{\nu\nu})$  at 3 TeV is 415 fb
- $BR(H \rightarrow \gamma\gamma)$  is 0.23%
- $\sigma(H_{\nu\nu}) \times BR(H \rightarrow \gamma\gamma) \approx 0.95$  fb
- $N_{\text{signal}} \approx 4750 \text{ evt} / 5 \text{ ab}^{-1}$



10 GeV photons

Silicon-tungsten ECAL energy resolution in ILD detector for 10 GeV photons in the barrel



# Simulation Details

- Assuming  $m_H = 126 \text{ GeV}$ ,  $5\text{ab}^{-1}$
- Polarization is conservatively taken as a scaling factor for signal and background.
- Signal and background simulation: WHIZARD v1.95, including ISR and BS and realistic luminosity spectrum.
- Particle interaction with the CLIC\_ILD detector is fully simulated.
- Hadronic background from BS is overlaid in the digitalization phase.
- Photons are reconstructed using PandoraPFA v02-04-00 photon processor.

## Generator Level Cuts:

- Applied to reduce CPU time
- On high cross-section backgrounds ( $ee \rightarrow \gamma\gamma$ ,  $ee \rightarrow e e \gamma$ ,  $ee \rightarrow e e \gamma \gamma$ ,  $ee \rightarrow \nu \nu \gamma$ ,  $ee \rightarrow \nu \nu \gamma \gamma$ ):
  - At least two photons per event with event with  $p_T > 10 \text{ GeV}$  and  $5^\circ < \theta_\gamma < 175^\circ$
  - At least one Higgs candidate with  $100 \text{ GeV} < M_{\gamma\gamma} < 150 \text{ GeV}$

# Signal and background processes


Process	$\sigma$ (fb)	Expected events at $5\text{ab}^{-1}$	Events Simulated
$\sigma(h\nu\nu) \times \text{BR}(h \rightarrow \gamma\gamma)$	0.95	4750	24 550
$ee \rightarrow \gamma\gamma$	867	$4.25 \cdot 10^6$	29 900
$ee \rightarrow ee\gamma$	185 392	$927 \cdot 10^6$	$3 \cdot 10^6$
$ee \rightarrow ee\gamma\gamma$	4 245	$21 \cdot 10^6$	144 900
$ee \rightarrow \nu\nu\gamma$	16 806	$82.5 \cdot 10^6$	194 900
$ee \rightarrow \nu\nu\gamma\gamma$	2 616	$13 \cdot 10^6$	155 500
$ee \rightarrow qq\gamma^*$	584	$3 \cdot 10^6$	$1.2 \cdot 10^6$
$ee \rightarrow qq\gamma\gamma^*$	72	360 000	299 600

\*Process without stdhep cut

- **Higgs candidate definition**
  - Identify two Higgs photons without variables to be used later in MVA
- **Preselection**
  - Reduction of the high cross-section backgrounds
- **Separation of signal with MVA**
  - Signal to background separation with the maximal statistical significance  $S$
- Observable to measure:  $\sigma (H\nu\nu) \times \text{BR}(H \rightarrow \gamma\gamma)$  ;  $g_{H\gamma\gamma}$  will be derived from a global fit
- Relative statistical uncertainty of the measurement:  $\delta (\sigma (H\nu\nu) \times \text{BR}(H \rightarrow \gamma\gamma)) = 1/S$

# Higgs Candidate Definition


- Exactly 2 isolated photons with  $p_T > 15$  GeV
  - Removing photons from ISR and machine background photons.
  - Signal loss: 14.8%
- Photon isolation -  $E_{\text{PFO}} < 20$  GeV within 14 mrad cone around the photon.
  - Reduces effective cross-section of background processes with FSR
  - 23% background rejection. Signal loss < 0.1%





# Higgs Candidate Definition: Signal Efficiency




- 14.8 % events with less than 2 photons
- 7.5 % events with 3 or more photons.
- Signal efficiency is 77.7 % by definition of the Higgs candidate.



# Preselection

Preselection:

- Two isolated photons (Higgs candidate)
- Candidate energy:  $100 \text{ GeV} < E(\gamma\gamma) < 1000 \text{ GeV}$
- Candidate transverse momentum:  $20 \text{ GeV} < p_T(\gamma\gamma) < 600 \text{ GeV}$
- Candidate invariant mass:  $110 \text{ GeV} < M(\gamma\gamma) < 140 \text{ GeV}$
- Preselection efficiency: 70%.
- Background is reduced by a factor of 1000.
- Signal to background ratio ( $N_s/N_b$ ) is  $7.8 \cdot 10^{-3}$  after preselection.



# MVA Variables

- All background processes used for MVA training

TMVA is optimised with thirteen sensitive observables:


- Higgs candidate mass :  $M(\gamma\gamma)$
- Higgs candidate energy:  $E(\gamma\gamma)$
- Higgs candidate transverse momentum:  $p_T(\gamma\gamma)$
- Higgs candidate polar angle:  $\theta(\gamma\gamma)$
- Cosine of the helicity angle:  $\cos \theta_{\text{hel}}$
- \*Photons transverse momenta:  $p_T(\gamma_1)$  and  $p_T(\gamma_2)$
- \*Photons polar angle:  $\theta(\gamma_1)$ ,  $\theta(\gamma_2)$
- \*Photons energy:  $E(\gamma_1)$ ,  $E(\gamma_2)$
- ECAL energy per event :  $E_{\text{ECAL}}$
- HCAL energy per event:  $E_{\text{HCAL}}$

\* Photons are sorted by higher value, where  $p_T(\gamma_1) > p_T(\gamma_2)$

# MVA Variables II


The most sensitive observables ranked by TMVA:

- Higgs candidate mass
- $p_T(\gamma_2)$
- Higgs candidate energy




# Correlations matrices

Correlation Matrix (signal)





Correlation Matrix (background)



Variables are sufficiently uncorrelated for MVA to perform


# BDTG Performance



- Best significance for  $\text{BDTG} > 0.68$
- BDTG efficiency: 54%
- Total signal efficiency (BDTG + preselection): 38%
- Signal events remaining after preselection and MVA:  $1790/5\text{ab}^{-1}$



# Higgs invariant mass distributions for signal and backgrounds before and after MVA



Remain backgrounds after MVA are either from photon +  $E_{\text{miss}}$  processes ( $ee \rightarrow \nu\bar{\nu}\gamma$ ,  $ee \rightarrow \nu\bar{\nu}\gamma\gamma$ ) or from the high cross-section  $ee \rightarrow ee\gamma$

$$\delta (\sigma(H\nu\nu) \times \text{BR}(H\gamma\gamma)) = 1/S = \frac{\sqrt{N_s+N_b}}{N_s} = 7.9\%$$

# Discussion

Process ( $5\text{ab}^{-1}$ )	Events after preselection	Preselection efficiency (%)	Events after MVA	Total selection efficiency (%)
$ee \rightarrow \gamma\gamma$	4 540	0.1	173	0.004
$ee \rightarrow ee\gamma$	272 570	0.03	7 718	0.00008
$ee \rightarrow ee\gamma\gamma$	10 605	0.05	241	0.001
$ee \rightarrow \nu\nu\gamma$	73 155	0.09	5 847	0.007
$ee \rightarrow \nu\nu\gamma\gamma$	47 283	0.04	3 636	0.03
$ee \rightarrow qq\gamma$	14 538	0.5	444	0.015
$ee \rightarrow qq\gamma\gamma$	3 873	1	175	0.05
<b>H<math>\nu\nu</math></b>	<b>3 320</b>	<b>70</b>	<b>1790</b>	<b>38</b>

- Statistical precision of the result is limited by the signal statistics and by the presence of irreducible or high cross-section backgrounds



# Impact of Polarization

5 ab <sup>-1</sup>	Signal Events	Background events	$\delta (\sigma(H\nu\nu) \times BR(H\gamma\gamma))$
Without polarization	1 790	18 236	7.9 %
With polarization	2 634	26 989	6.5 %

- Conservatively taken to increase both signal and background cross-sections by a factor of 1.48, polarization improves the statistical precision to 6.5%

# List of systematic uncertainties

- Integral luminosity uncertainty ( $10^{-3}$ )→negligible
- Uncertainty of the luminosity spectrum (0.15%)
- Relative uncertainty of the electron beam polarization (0.2%)
- Uncertainty of the photon identification efficiency (1%)
- Photon energy resolution uncertainty (2% relative uncertainty of the sampling term) → negligible

Effect	Systematic Uncertainty
Luminosity spectrum	0.15%
Beam polarization	0.1%
Photon identification	0.11%
<b>Total uncertainty</b>	<b>0.21%</b>

# Concluding remarks

- The highest energy operation at CLIC enables  $\sigma(H\nu\nu) \times BR(H \rightarrow \gamma\gamma)$  measurement with the relative statistical uncertainties of 7.9% and ultimately with 6.5% including electron-beam polarization.
- All relevant physics and background processes are fully simulated, as well as the response of the CLIC\_ILD detector.
- Systematic uncertainties (luminosity spectrum, uncertainty of the photon identification efficiency, uncertainty of the photon energy resolution, etc.) are estimated to be an order of magnitude smaller than the statistical one.

*This analysis was part of my PhD thesis and the corresponding paper is in preparation.*

# ХВАЛА НА ПАЖЊИ




# Back Up




# Signal and background selection efficiencies

5 ab <sup>-1</sup>	Signal Events	Background events	$\delta (\sigma(H\nu\nu) \times BR(H\gamma\gamma))$
Without polarization	1 790	18 236	7.9 %
With polarization	2 634	26 989	6.5 %

# Distributions of MVA variables




# Distributions of MVA variables





# Dependence on ECAL Resolution in Fast Simulation at CLIC at 1.4 TeV

- Differences due to non-Gaussian tails and non-linearity in full simulation
- Improvement of sampling term 20%  $\rightarrow$  5%:  
rel. stat. uncertainty: 13.7%  $\rightarrow$  12.1% (with constant term)  
rel. stat. uncertainty: 12.9%  $\rightarrow$  10.5% (no constant term)



Credit: C. Grefe