

Cross sections for Cosmic Rays @ CERN

November 13th to 15th 2019

The COMPASS++/AMBER program for cross-sections measurements

P. Zuccon

Trento University and INFN TIFPA




UNIVERSITÀ DEGLI STUDI
DI TRENTO



Introduction

- anti-p production cross section from p-p and p-He interactions is poorly measured and cannot simply be constrained from available measurements.
- an accurate prediction of the expected anti-p flux in cosmic rays in the rigidity range from few GeV to several hundreds of GeVs, is interesting to understand cosmic ray and possibly search for signals of new physics
- LHC-b collaboration reported a measurement of the anti-p XS from 8 TeV p-He, and foresee a similar measurement with 4 TeV protons.
- NA61 published p-p to anti-p at 20, 31, 40, 80, and 158 GeV/c
- we want to investigate the possibility to perform a measurement with the SPS protons between 50 and 280 GeV/c on fixed LH2 and LHe targets, and a magnetic spectrometer

Fraction origin of anti-p from CR interaction with ISM




Martin W. Winkler (Stockholm University)

LHCb-CONF-2017-002
 Measurement performed at 7 TeV
 $p\text{-He} \rightarrow \bar{p} + X$

NA61 p+p data beam momenta of
 20, 31, 40, 80, and 158 GeV/c
 Eur. Phys. J. C 77, 671 (2017)




AMS DATA on \bar{p}/p



Two major uncertainties limit the prediction of the anti-p flux from CR interaction with ISM


- production cross sections $p-p \rightarrow \bar{p} + X$ $p\text{-He} \rightarrow \bar{p} + X$
- CR propagation in the galaxy

AMS Data vs Predictions



Overall Prediction Uncertainties

Nicolò Masi, INFN-Bo



COMPASS++/AMBER

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH



CERN-SPSC-2019-003
SPSC-I-250
January 28, 2019

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH



CERN-SPSC-2019-022
SPSC-P-360
October 14, 2019

Letter of Intent

Letter of Intent:

A New QCD facility at the M2 beam line of the CERN SPS*

COMPASS++[†]/AMBER[†]

B. Adams^{13,12}, C.A. Aidala¹, R. Akhunzyanov¹⁴, G.D. Alexeev¹⁴, M.G. Alexeev⁴¹, A. Amoroso^{41,42},
V. Andrieux⁴⁴, N.V. Anfimov¹⁴, V. Anosov¹⁴, A. Antoshkin¹⁴, K. Augsten^{14,32}, W. Augustyniak⁴⁶,
-----⁴-----²-----⁴⁷-----^{41,42}-----⁸-----⁹-----⁸

Proposal for Measurements at the M2 beam line of the CERN SPS

- Phase-1 -

COMPASS++^{*}/AMBER[†]

Proposal to SPSC ->



[hep-ex] 25 Jan 2019

COMPASS++/ AMBER

Proposal for Measurements at the M2 beam line of the CERN SPS

- Phase-1 -

COMPASS++^{*}/AMBER[†]

Year	Activity	Duration	Beam
2021	Proton radius test measurement	20 days	μ
2022	Proton radius measurement	120 (+40) days	μ
	Antiproton production test measurement 	10 days	p
2023	Antiproton production measurement 	20(+10) days	p
	Proton radius measurement	140 (+10) days	μ
2024 2024+	Drell-Yan: pion PDFs and charmonium production mechanism	$\lesssim 2$ years	$p, K^+, \pi^+, \bar{p}, K^-, \pi^-$

People involved

- UniTN and TIFPA, P. Zuccon, F. Nozzoli
- UniBO and INFN, N. Masi, L. Quadrani, A. Contin
- UniTO and INFN, M. Chiosso, O. Denisov, F. Donato, M. Kosmeier
- Nagoya University, N. Horikawa (cryo – targets)

- Support from the COMPASS++/AMBER community at large

When ?

Autumn 2022: commissioning run 10 days

Spring 2023: data taking run 20 days

COMPASS @ CERN






Fig. 4. Side view of the target region with the liquid hydrogen target system.

Acceptance:
 ± 180 mrad
 ± 10 deg
 $2.4 < \eta < 6$





New anti-p cross sections with COMPASS++/AMBER

- Use secondary proton beam from SPS
50, 100, 190, 280 GeV/c
- Use **Liquid Hydrogen** target and **Liquid He** target
- Use the COMPASS spectrometer to reconstruct the **inelastic events** -> Negative tracks and their momentum
- Use the COMPASS RICH detector to identify **anti-p** and reject π and **K**
- Provide a measurement the **anti-p** production cross section for p-p and p-He.

Expected performances





p-LH2 event features @ 190 GeV/c

3.05 10^5 interacting events




p-LHe event features @ 190 GeV/c

2.25 10^5 interacting events




COMPASS Rec accuracy


Tracking Efficiency



Tracking Efficiency




Momentum resolution



COMPASS simulation (p-p 190GeV)

Antiproton tracking efficiency
Pseudo-rapidity vs $\log_{10}(\text{momentum})$




Estimate RICH particle identification performance
from p-p data already collected at 190 GeV

Select pure samples of ρ , ρ bar, π^+ and π^-


DATA SET collected in 2009 (p-LH2)

Select a sample of V0

- two tracks forming a vertex
- $\rho = p_1 + p_2$ points to the primary vertex




Armenteros plot




$$\alpha = \frac{p_{l1} - p_{l2}}{p_{l1} + p_{l2}}$$


Mass selection




Mass selection




Lambda Mass





Selected Tracks from K_0 and Λ_0




RICH angle vs $\beta\gamma$



π^+




p






Gaussian Fit of the RICH angle


MPV



MPV



Universal RICH response function



anti-p and p are identified above threshold (~ 18 GeV/c) with an efficiency $>95\%$ up to ~ 60 (GeV/c)

From the Kaon threshold (~ 9 GeV/c) to the p threshold, p and anti-p are identified using RICH in veto mode.

Combine the elements to obtain a cross
section measurement

Cross section measurement

- Strategy

- Count all the p-p (or p-He) interaction in the target (R_i)
- Identify events with one (or multiple) anti-p vs reconstructed momentum and angle ($R_s(p, \theta)$)
- Calculate the double differential cross section as

$$\frac{d\sigma_{\bar{p}}}{dp d\theta} = \frac{R_s(p, \theta)}{R_i} \sigma_{pp}$$

- Several possible pitfalls and sources of systematic errors!

Summary of the expected errors

Systematic

	efficiency	est sys error
Track Recon	95%	~1%
Rich Efficiency	~ 99%	~0.5 %
RICH PID	99 to 75 %	0.1 to 4%
Trigger		1%
Vertex error	98%	0.5%
Beam Purity	99.9%	0.5%
TOTAL		4 to 6 %

Statistical Errors when considering:

- 2 targets (LHe and LH2, each 50 cm long),
- The estimated acceptance and efficiency from the full COMPASS MC.
- 20 bins in momentum from [10, 50] GeV/c and 20 bin in P_t
- 75% beam purity at 5×10^5 p/s beam intensity

For a single combination p-momentum and target type

4(12) hours of beam time → expect typical **1.7(1.0) % statistical** error


Relevance of the measurement

p-p source term coverage



M. Korsmeier

Red AMBER


Black NA61




p-He He-p Source term coverage



Comparison of $p\text{-}p \rightarrow \bar{p} + X$ with NA61




Eur.Phys.J. C77 (2017) no.10, 671



Expected cross section measurement $p\text{-He} \rightarrow \bar{p} + X$

p 190 GeV/c



Summary

- CR antimatter data sensitivity to exotic sources is degraded by the poor knowledge of the $p\bar{p}$ production cross sections
- We have shown that a measurement of $p\text{-}p\text{-}\rightarrow p\bar{p} + X$ and $p\text{-He}\text{-}\rightarrow p\bar{p} + X$ can be performed at CERN with the COMPASS++/AMBER spectrometer at momenta ranging from 50 GeV/c to 280 GeV/c
- A new collaboration is being set up and a proposal to SPSC has been submitted
- The new COMPASS++/AMBER data might have an impact on the DM sensitivity on the CR $p\text{-}\bar{p}$ channel



GO LUCA!
Long Life to AMS





Backup Slides



Coverage of 190 GeV/c p on LHe



Particle range at 50 GeV/c momentum

MC_PID	Name	range	Mass	Charge	Lifetime	Energy	Beta	gamma
		m	GeV/c ²		s	GeV		
20	SIGMA 0	9.31E-10	1.19E+00	0	7.40E-20	5.00E+01	1.00E+00	42
28	ANTISIGMA 0	9.31E-10	1.19E+00	0	7.40E-20	5.00E+01	1.00E+00	42
17	ETA	1.50E-08	5.48E-01	0	5.49E-19	5.00E+01	1.00E+00	91
7	PION 0	9.33E-06	1.35E-01	0	8.40E-17	5.00E+01	1.00E+00	370
24	OMEGA -	7.38E-01	1.67E+00	-1	8.22E-11	5.00E+01	9.99E-01	30
32	ANTIOMEGA +	7.38E-01	1.67E+00	1	8.22E-11	5.00E+01	9.99E-01	30
27	ANTISIGMA -	1.01E+00	1.19E+00	-1	7.99E-11	5.00E+01	1.00E+00	42
19	SIGMA +	1.01E+00	1.19E+00	1	7.99E-11	5.00E+01	1.00E+00	42
21	SIGMA -	1.85E+00	1.20E+00	-1	1.48E-10	5.00E+01	1.00E+00	42
29	ANTISIGMA +	1.85E+00	1.20E+00	1	1.48E-10	5.00E+01	1.00E+00	42
23	XI -	1.86E+00	1.32E+00	-1	1.64E-10	5.00E+01	1.00E+00	38
31	ANTIXI +	1.86E+00	1.32E+00	1	1.64E-10	5.00E+01	1.00E+00	38
16	KAON 0 SHORT	2.69E+00	4.98E-01	0	8.93E-11	5.00E+01	1.00E+00	100
22	XI 0	3.31E+00	1.32E+00	0	2.90E-10	5.00E+01	1.00E+00	38
30	ANTIXI 0	3.31E+00	1.32E+00	0	2.90E-10	5.00E+01	1.00E+00	38
18	LAMBDA	3.54E+00	1.12E+00	0	2.63E-10	5.00E+01	1.00E+00	45
26	ANTILAMBDA	3.54E+00	1.12E+00	0	2.63E-10	5.00E+01	1.00E+00	45
12	KAON -	3.76E+02	4.94E-01	-1	1.24E-08	5.00E+01	1.00E+00	101
11	KAON +	3.76E+02	4.94E-01	1	1.24E-08	5.00E+01	1.00E+00	101
10	KAON 0 LONG	1.56E+03	4.98E-01	0	5.17E-08	5.00E+01	1.00E+00	100
9	PION -	2.80E+03	1.40E-01	-1	2.60E-08	5.00E+01	1.00E+00	358
8	PION +	2.80E+03	1.40E-01	1	2.60E-08	5.00E+01	1.00E+00	358
6	MUON -	3.12E+05	1.06E-01	-1	2.20E-06	5.00E+01	1.00E+00	473
5	MUON +	3.12E+05	1.06E-01	1	2.20E-06	5.00E+01	1.00E+00	473
13	NEUTRON	1.42E+13	9.40E-01	0	8.87E+02	5.00E+01	1.00E+00	53
25	ANTINEUTRON	1.42E+13	9.40E-01	0	8.87E+02	5.00E+01	1.00E+00	53
46	TRITON	5.35E+24	2.81E+00	1	1.00E+15	5.01E+01	9.98E-01	18
45	DEUTERON	8.00E+24	1.88E+00	1	1.00E+15	5.00E+01	9.99E-01	27
15	ANTIPROTON	1.60E+25	9.38E-01	-1	1.00E+15	5.00E+01	1.00E+00	53
14	PROTON	1.60E+25	9.38E-01	1	1.00E+15	5.00E+01	1.00E+00	53
3	ELECTRON	2.94E+28	5.11E-04	-1	1.00E+15	5.00E+01	1.00E+00	97847
2	POSITRON	2.94E+28	5.11E-04	1	1.00E+15	5.00E+01	1.00E+00	97847
49	HE3	5.35E+24	2.81E+00	2	1.00E+15	5.01E+01	9.98E-01	18
47	ALPHA	4.04E+24	3.73E+00	2	1.00E+15	5.01E+01	9.97E-01	13

p-p 190 GeV/c LH2



Compass Trigger system




Fig. 54. Allowed combinations for target pointing in the RPD part of the proton trigger.




Fig. 4. Side view of the target region with the liquid hydrogen target system.




Fig. 51. Arrangement of trigger elements in the spectrometer (schematic side view, not to scale).





Fig. 53. Time residual of the beam trigger.

Rate statistics and pileup

- Typical beam intensity is $5 \cdot 10^7$ p for a 9.8s spill
- We expect $\sim 5.4\%$ of the protons to interact with the 40cm LH2 target $\rightarrow \sim 270\text{k}$ interaction/s
- Compass standard trigger DT0 is *BT & BK & Sandwich Veto & RPD*
- This reduce the trigger rate to 33 kHz which can be handled by the COMPASS DAQ
- For the future measurements we will ask to reduce the beam intensity to $5 \cdot 10^6$ and plan to use the trigger *BT & BK & Sandwich Veto*
- This will provide an expected rate of 25 kHz


Upstream Threshold
Cherenkov counter



Beam and Trigger error
 $\sim 0.5\%$

Lost Interaction events

- Select a fiducial volume on the target [-68,-30] cm
- Look how many events have a reconstructed vertex within the fiducial volume



MC events: 288312

No Vertex: 2753 (0.95%)

Vtx outside: 2856 (0.99%)

Thanks to the Recoil Detector
no-vertex events can be cross-
checked with data

Vertex error 0.5%