

Polarization studies with MadGraph5_aMC@NLO

Diogo Buarque Franzosi,
Olivier Mattelaer, Richard Ruiz, Sujay Shil

MCNet meeting

Outline

- Introduction
- Implementation and syntax in `MadGraph5_aMC@NLO`
- Applications:
 - ① High energy longitudinal VBS
 - ② Decaying weak bosons
- Conclusion

Motivation

- Longitudinal weak bosons (W_0) are part of EW symmetry breaking (EWSB) sector
- Left-handed weak bosons (W_L) are enhanced in $q\bar{q} \rightarrow W^+$ production due to the left nature of EW gauge
- Top-quark mass related to EW sector $y_t \sim 1$, helicity flipping sensitive
- BSM effects can be highly polarization dependent, e.g. Right-handed current in extended gauge theories (Beg, Budny, Mohapatra, Sirlin 77), Dynamical EWSB and Composite Higgs couples strongly to W_0, \dots
- In the EFT framework different operators contribute to different polarizations, e.g. in VV polarizations e.g. Liu, Wang, 18 and top-quark physics (Hartland, Maltoni, Nocera, Rojo, Slade, Vryonidou, Zhang 19)
- Huge interest from experimental collaborations

Having each polarization separately described is crucial to understand their contributions to physical processes

Polarization in MadGraph5_aMC@NLO

- MadGraph5_aMC@NLO offers a very flexible and powerful framework to simulate collision events,
 - for “arbitrary” multiparticles in the final state,
 - at NLO in QCD accuracy (via e.g. MadLoop and MadFKS),
 - matched to PS,
 - with merged jet multiplicities,
 - for several types of models (UFO format),
 - spin correlated decay via MadSpin or decay chain,
 - interfaced to several analysis frameworks, etc...

In this release: particle polarizations can be specified.

MadGraph5_aMC@NLO implementation

MODE I: Initial/final state particles

- The syntax $\mathcal{P}\{X\}$ is used to define polarization X of particle \mathcal{P} , e.g.

```
generate p p > t t~{L}
generate e+{L} e- > w+{L} w-{T}
generate p p > w+ w-{T} [QCD]
```

- Implemented for particles with spin 1/2, 1, 3/2 and 2
- For LO generation we allow the user to choose the *frame* where polarizations are defined in run_card.dat

```
[1,2] = me_frame
      ! list of particles to sum-up to define the frame
```

MODE II: Unstable particles in the spin correlated narrow width approximation

- Both decay chain, e.g.

generate $p\ p \rightarrow t\ t^{\sim}\{L\}$, $t^{\sim} \rightarrow b^{\sim}\ w^-$

generate $e^+\{L\}\ e^- \rightarrow w^+\{L\}\ w^-\{T\}$, $w^+ \rightarrow e^+\ ve$, $w^- \rightarrow e^- ve^+$

- and MadSpin on a sample with polarized particles, e.g.

decay $w^+ \rightarrow ta^+\ vt$

- Replace propagator numerator with the subset of helicity sum

$$\text{spin 1: } \eta^{\mu\nu} + \frac{p^\mu p^\nu}{m^2} \rightarrow \sum_\lambda \epsilon_\lambda^{\mu^*} \epsilon_\lambda^\nu$$

$$\text{spin 1/2: } \not{p} - m \rightarrow \sum_\lambda u_\lambda \bar{u}_\lambda$$

- Support spin 1/2 and 1
- Implementation does not support interference between different polarizations (this has to be generated with the standard unpolarized ME)

Feature	Unpolarized	Polarized
LO Parton Shower	✓	✓
Merging LO	✓	✓
PS Matching NLO QCD	✓	QCD neutral particles
Merging NLO ($F_x F_x$)	✓	QCD neutral particles
Polarized beams	✓	✓
NLO Fixed order	✓	✓
BSM via UFO	✓	✓
MadSpin and decay chain	✓	Spin 1/2, 1

Application I: High energy VBS

- If the Higgs has modified couplings to weak bosons (e.g. if it is composite)

$$\mathcal{L} \supset m_W^2 W_\mu^+ W^{-\mu} \left(1 + 2a \frac{h}{v} + \dots \right)$$

the miscancellation of diagrams gives energy growing of longitudinal boson scattering amplitudes at high energies

$$A(W_L W_L \rightarrow W_L W_L) \sim (1 - a^2) \frac{s}{v^2}$$

- and affects the prediction in “more physical” observables like VBS, e.g.

$$pp \rightarrow jj W^+ W^-, \quad W \rightarrow f\bar{f}$$

- Our framework can be used to predict each polarization component and in particular enrich the longitudinal sample where effects are expected

- We use the Higgs Characterization model (replace X,Y=L,T)


```
import model HC_UFO-CH
generate p p > j j w+{X} w-{Y} / x1 x2 QCD=0
output VBSCH_pp-wpXwmY
```

- Consider $a = 1$ (SM) and $a = 0.8$, partonic C.M. (pRF) and WW C.M. (WWRF)
- Cuts: $M(WW) > 300 \text{ GeV}$, $\Delta\eta(jj) > 3.6$, $M(jj) > 600 \text{ GeV}$, $p_T(j) > 20 \text{ GeV}$


Process	pRF, SM		pRF, $a = 0.8$		BSM/SM
	σ [pb]	ε [%]	σ [pb]	ε [%]	
$jjW^+ W^-$	0.1526	100	0.1563	100	1.02
$jjW_T^+ W_T^-$	0.09552	62.6	0.09641	61.7	1.01
$jjW_L^+ W_T^-$	0.02411	15.8	0.02421	15.5	1.00
$jjW_T^+ W_L^-$	0.02569	16.8	0.02575	16.5	1.00
$jjW_L^+ W_L^-$	0.007912	5.18	0.009539	6.10	1.20

Process	WWRF, SM		WWRF, $a = 0.8$		BSM/SM
	σ [pb]	ε [%]	σ [pb]	ε [%]	
$jjW^+ W^-$	0.1526	100	0.1563	100	1.02
$jjW_T^+ W_T^-$	0.09449	61.9	0.09525	60.9	1.01
$jjW_L^+ W_T^-$	0.02366	15.5	0.02289	14.6	0.967
$jjW_T^+ W_L^-$	0.02569	16.8	0.02556	16.3	0.995
$jjW_L^+ W_L^-$	0.00980	6.4	0.01196	7.6	1.22

pRF



WWRF



Application II: Decaying the W



- For one W decay, in the W rest frame, the cross section can be written as polarization fractions f_X and the interference terms g_{XY}
e.g. Mirkes 92, Stirling, Vryonidou 12, Bern, Diana, Dixon 11

$$\begin{aligned}\sigma^{-1} \frac{d^2\sigma}{d \cos \theta^* d\phi^*} = \\ \frac{1}{4\pi} \left\{ f_+ \frac{(1 - \cos \theta^*)^2}{2} + f_- \frac{(1 + \cos \theta^*)^2}{2} + f_0 (1 - \cos \theta^*)^2 \right. \\ \left. + \sqrt{2} g_{+0} \sin \theta^* (1 - \cos \theta^*) \cos \phi^* - \sqrt{2} g_{-0} \sin \theta^* (1 + \cos \theta^*) \cos \phi^* \right. \\ \left. - g_{+-} \sin^2 \theta^* \cos(2\phi^*) \right\}\end{aligned}$$

- Interference vanishes for ϕ integrated over all phase space.
- Typically measurements of pol. ratios assume integration over ϕ e.g. CMS 1104.3829, but selection cuts “resurrect” them, e.g. for $W+jets$ Belyaev, Ross 13 and VBS Ballestrero, Maina, Pellicoli 18

- The polarized samples can be decayed with MadSpin
- decay $w+ > \mu^+ \nu_\mu$
- decay $w- > e^- \bar{\nu}_e$
- Cut: $p_T(j) > 20 \text{ GeV}$
 - Independent implementation to PHANTOM Ballestrero, Maina, Pelliccioli

18



- Loop induced processes $gg \rightarrow W_\lambda^+ W_{\lambda'}^-$, $W^\pm \rightarrow l^\pm \nu_\ell$

generate g g > w+{L} w-{L} [QCD]

\sqrt{s} Process	13 TeV		14 TeV		27 TeV		100 TeV	
	σ [pb]	ε [%]	σ [pb]	ε [%]	σ [pb]	ε [%]	σ [pb]	ε [%]
Inclusive $gg \rightarrow W_\lambda^+ W_{\lambda'}^-$ at LO								
WW	3.031	...	3.504	...	11.19	...	78.28	...
$W_T W_T$	2.794	(92%)	3.2	(91%)	10.16	(91%)	71.76	(92%)
$W_L W_T$	0.1519	(5%)	0.1742	(5%)	0.5589	(5%)	3.81	(5%)
$W_L W_L$	0.1224	(4%)	0.1412	(4%)	0.4292	(4%)	2.906	(4%)
$p_T(W^\pm) > 250 \text{ GeV}$								
WW	14.01×10^{-3}	...	17.19×10^{-3}	...	89.86×10^{-3}	...	1.193	...
$W_T W_T$	10.84×10^{-3}	(77%)	13.35×10^{-3}	(77%)	70.06×10^{-3}	(78%)	0.9297	(78%)
$W_L W_T$	0.5661×10^{-3}	(4%)	0.6945×10^{-3}	(4%)	3.549×10^{-3}	(4%)	0.04597	(4%)
$W_L W_L$	2.538×10^{-3}	(18%)	3.107×10^{-3}	(18%)	16.71×10^{-3}	(19%)	0.22	(18%)

Conclusion

- In the MadGraph5_aMC@NLO framework we provide the possibility to generate events at LO or NLO accuracy matched to PS for polarized particles
- Having the different polarization contributions separately is useful, for example:
 - in order to estimate distributions more precisely,
 - estimate the effect of new physics in a particular polarization
 - devise selection cuts to enhance a particular polarization

Backup

spin $\frac{1}{2}$	syntax	HELAS equivalent	propagtor
	{L} {+}	+1 (Left)	Yes
	{R} {-}	-1 (Right)	Yes
spin 1	syntax	HELAS equivalent	propagtor
	{L} {0}	0 (Longitudinal)	Yes
	{T}	1 and -1 (Transverse)	Yes
	{+}	+1	No
	{-}	-1	No
	{A}	Only for propagators	
spin 3/2	syntax	HELAS equivalent	propagtor
	{-1}	-1	No
	{1}	1	No
	{3}	3	No
	{-3}	-3	No
spin 2	syntax	HELAS equivalent	propagtor
	{-2}	-2	No
	{-1}	-1	No
	{0}	0	No
	{1}	1	No
	{2}	2	No

Beam polarization

process	polbeam1	cross-section	expected
$e^+e^- \rightarrow tt$	0	0.1664	
$e^+e^- \rightarrow tt$	100	0.2296	
$e^+e^- \rightarrow tt$	-100	0.1033	
$e^+e^- \rightarrow tt$	25	0.182	0.1822375
$e^+e^- \rightarrow tt$	50	0.1983	0.198025
$e^+e^- \rightarrow tt$	75	0.2137	0.2138125
$e_L^+e^- \rightarrow tt$	0	0.1033	0.1033
$e_L^+e^- \rightarrow tt$	100	0	0
$e_L^+e^- \rightarrow tt$	-100	0.1036	0.1033
$e_R^+e^- \rightarrow tt$	0	0.2293	0.2296
$e_R^+e^- \rightarrow tt$	100	0.2296	0.2296
$e_R^+e^- \rightarrow tt$	-100	0	0
$e_R^+e^- \rightarrow tt$	25	0.143	0.1435
$e_R^+e^- \rightarrow tt$	50	0.1719	0.1722
$e_R^+e^- \rightarrow tt$	75	0.2008	0.2009