

Workshop on Forward Physics and QCD at the LHC, the future Electron Ion Collider and Cosmic Ray Physics



Contribution ID: 54

Type: **not specified**

The use of QCD evolution to detect gluon saturation in exclusive photo-production of vector mesons

Wednesday, 20 November 2019 15:35 (35 minutes)

We investigate photo-production of vector mesons J/Ψ and Upsilon measured both at HERA and LHC, using 2 particular fits of inclusive unintegrated gluon distributions. The fits are based on non-linear Balitsky-Kovchegov evolution (Kutak-Sapeta gluon; KS) and next-to-leading order Balitsky-Fadin-Kuraev-Lipatov evolution (Hentschinski-Sabio Vera-Salas gluon; HSS). We find that linear next-to-leading order evolution can only describe production at highest energies, if perturbative corrections are increased to unnaturally large values; rendering this corrections to a perturbative size, the growth with energy is too strong and the description fails. At the same time, the KS gluon, which we explore both with and without non-linear corrections, requires the latter to achieve an accurate description of the energy dependence of data. We interpret this observation as a clear signal for the presence of high gluon densities in the proton, characteristic for the onset of gluon saturation.

Presenter: HENTSCHINSKI, Martin (Universidad de las Americas, Puebla)

Session Classification: Saturation session