

Elliptic and Kummer-elliptic Integrals in higher order calculations in QFT

Johannes Blümlein

DESY, Zeuthen



J. Blümlein et al. Phys.Lett. B791 (2019) 206; arXiv:1903.06155;
J. Ablinger et al. J.Math.Phys. 59 (2018) no.6, 062305; PoS LL2018 (2018) 017.
J. Blümlein, A. De Freitas, C. Raab, and K. Schönwald, ' The $O(\alpha^2)$ Initial State QED Corrections to $e^+e^- \rightarrow \gamma^*/Z_0^*$ ', DESY 18–196.

Elliptics 19, KMPB, AEI Potsdam

September 20, 2019

Introduction

One of the main and difficult issues in high energy physics is the calculation of involved multi-dimensional integrals.

In the following our attitude will be their **analytic integration**.

For quite some classes of integrals, particularly at lower order in the coupling constant, quite a series of analytic computational methods exist. cf. e.g. CPC 202 (2016) 33 [arXiv:1509.08324] for the algorithms.

- ▶ Hypergeometric functions.
- ▶ Mellin-Barnes representations.
- ▶ In the case of **convergent** massive 3-loop Feynman integrals, they can be performed in terms of **Hyperlogarithms** [Generalization of a method by F. Brown, 2008, to non-vanishing masses and local operators].

Introduction

- ▶ Summation methods based on difference fields, implemented in the Mathematica program **Sigma** [C. Schneider, 2005–].
 - ▶ Reduction of the sums to a small number of key sums.
 - ▶ Expansion of the summands in ε .
 - ▶ Simplification by symbolic summation algorithms based on $\Pi\Sigma$ -fields [Karr 1981 J. ACM, Schneider 2005–].
 - ▶ Harmonic sums, polylogarithms and their various generalizations are algebraically reduced using the package HarmonicSums [Ablinger 2010, 2013, Ablinger, Blümlein, Schneider 2011, 2013].
- ▶ Systems of Differential Equations. Nucl.Phys. B939 (2019) 253 [arXiv:1810.12261]
- ▶ Almkvist-Zeilberger Theorem as Integration Method. [Multi-Integration]

In the following we will concentrate on the method of **Differential Equations** since these are automatically obtained from the **integration-by-parts identities** representing all integrals by the so-called master integrals.

These may either be considered directly or in terms of difference equations obtained through a formal power-series ansatz or a Mellin transform.

Introduction

Starting from the most simple cases and moving to gradually more and more involved (massive) topologies one observes:

- ▶ The lower order topologies correspond to **differential or difference equation systems** which are **first order factorizable**.
- ▶ Here, a wider class of **solution methods** exists. There are methods in both cases to **constructively find** all letters of the alphabet needed to express the solutions in terms of **indefinitely nested sums or iterative integrals**.
- ▶ Later also **differential or difference equations** occur which contain **genuine higher than 1st order factors**.
- ▶ The **first example** are **${}_2F_1$ solutions**. In special cases these are also **elliptic solutions**.
- ▶ In the latter case one may represent the solutions in terms of **modular functions** and in **more special cases** in terms of **modular forms** and therefore in **polynomials of Lambert-Eisenstein series** (elliptic polylogarithms).

Nested Sums & Iterative Integrals

Indefinitely nested sums:

$$S(N) = \sum_{k_1=1}^N s(k_1) \sum_{k_2=1}^{k_1} s(k_2) \dots \sum_{k_m=1}^{k_{m-1}} s(k_m)$$

Iterated integrals:

$$F(x) = \int_0^x dy_1 f_1(y_1) \int_0^{y_1} dy_2 f_2(y_2) \dots \int_0^{y_{l-1}} dy_l f_l(y_l)$$

Mellin transform:

$$\sum_{\alpha} c_{\alpha} S_{\alpha}(N) = \int_0^1 dx x^{N-1} F(x)$$

... much more to say about the historic development, cf. e.g. J. Ablinger, JB, C. Schneider, 1304.7071, 1310.5645

Function Spaces

Sums

Harmonic Sums

$$\sum_{k=1}^N \frac{1}{k} \sum_{l=1}^k \frac{(-1)^l}{l^3}$$

gen. Harmonic Sums

$$\sum_{k=1}^N \frac{(1/2)^k}{k} \sum_{l=1}^k \frac{(-1)^l}{l^3}$$

Cycl. Harmonic Sums

$$\sum_{k=1}^N \frac{1}{(2k+1)} \sum_{l=1}^k \frac{(-1)^l}{l^3}$$

Binomial Sums

$$\sum_{k=1}^N \frac{1}{k^2} \binom{2k}{k} (-1)^k$$

Integrals

Harmonic Polylogarithms

$$\int_0^x \frac{dy}{y} \int_0^y \frac{dz}{1+z}$$

gen. Harmonic Polylogarithms

$$\int_0^x \frac{dy}{y} \int_0^y \frac{dz}{z-3}$$

Cycl. Harmonic Polylogarithms

$$\int_0^x \frac{dy}{1+y^2} \int_0^y \frac{dz}{1-z+z^2}$$

root-valued iterated integrals

$$\int_0^x \frac{dy}{y} \int_0^y \frac{dz}{z\sqrt{1+z}}$$

iterated integrals on CIS fct.

$$\int_0^z dx \frac{\ln(x)}{1+x} {}_2F_1 \left[\begin{matrix} \frac{4}{3}, \frac{5}{3} \\ 2 \end{matrix}; \frac{x^2(x^2-9)^2}{(x^2+3)^3} \right]$$

Special Numbers

multiple zeta values

$$\int_0^1 dx \frac{\text{Li}_3(x)}{1+x} = -2\text{Li}_4(1/2) + \dots$$

gen. multiple zeta values

$$\int_0^1 dx \frac{\ln(x+2)}{x-3/2} = \text{Li}_2(1/3) + \dots$$

cycl. multiple zeta values

$$\mathbf{C} = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2}$$

associated numbers

$$H_{8,w_3} = 2\text{arccot}(\sqrt{7})^2$$

associated numbers

$$\int_0^1 dx {}_2F_1 \left[\begin{matrix} \frac{4}{3}, \frac{5}{3} \\ 2 \end{matrix}; \frac{x^2(x^2-9)^2}{(x^2+3)^3} \right]$$

shuffle, stuffle, and various structural relations \implies algebras

Except the last line integrals, all other ones stem from 1st order factorizable equations.

The world until ~ 1997

Calculations up to ~ 2 Loops massless and single mass:

- ▶ Express all results in terms of $\text{Li}_n(z) = \int_0^z dx \text{Li}_{n-1}(x)/x$, $\text{Li}_0(x) = x/(1-x)$.
- ▶ and possibly $S_{p,n}(z) = (-1)^{n+p-1}/(p!(n-1)!) \int_0^1 dx \ln^{p-1}(x) \ln^n(1-xz)/x$.
- ▶ The argument $z = z(x)$ becomes a more and more complicated function.
- ▶ **covering algebras** of wider function spaces were widely unknown in physics, **despite they were known in mathematics** ...
- ▶ **The complexity of expressions grew significantly**, calling urgently for mathematical extensions.
- ▶ More complex argument structures **do not easily allow an analytic Mellin inversion**.
- ▶ Extremely long expressions are obtained, which would be much more compact, **using adequate mathematical functions**.
- ▶ Somewhen, new functions appeared: $\int_0^z dx \text{Li}_3(x)/(1+x)$ not fitting into this frame.

Spill-Off: New Function Classes and Algebras

- ▶ 1998: Harmonic Sums [Vermaseren; JB]
- ▶ 1999: Harmonic Polylogarithms [Remiddi, Vermaseren]
- ▶ 2000,2003, 2009: Analytic continuation of harmonic sums, systematic algebraic reduction; structural relations [JB]
- ▶ 2001: Generalized Harmonic Sums [Moch, Uwer, Weinzierl]
- ▶ 2004: Infinite harmonic (inverse) binomial sums [Davydychev, Kalmykov; Weinzierl]
- ▶ 2011: (generalized) Cyclotomic Harmonic Sums, polylogarithms and numbers [Ablinger, JB, Schneider]
- ▶ 2013: Systematic Theory of Generalized Harmonic Sums, polylogarithms and numbers [Ablinger, JB, Schneider]
- ▶ 2014: Finite nested Generalized Cyclotomic Harmonic Sums with (inverse) Binomial Weights [Ablinger, JB, Raab, Schneider]
- ▶ 2014-: Elliptic integrals with (involved) rational arguments.
- ▶ now: More-scale problem: Kummer-elliptic integrals

Particle Physics Generates **NEW** Mathematics.

Decoupling of Systems

- ▶ We consider linear systems of N inhomogeneous differential equations and decouple them into a single scalar equation + $(N - 1)$ other determining equations.
- ▶ Usually one may use a series ansatz (+ $\ln^k(x)$ modulation)

$$f(x) = \sum_{k=1}^{\infty} a(k)x^k$$

and obtain

$$\sum_{k=0}^m p_k(N)F(N+k) = G(N)$$

- ▶ The latter equation is now tried to be solved using [difference-field techniques](#).
- ▶ If the equation has successive 1st order solutions one ends up with a nested sums solution. All these cases have been algorithmized. [\[arXiv:1509.08324 \[hep-ph\]\]](#).
- ▶ This even applies for some cases ending up elliptic in x -space [\[arXiv:1310.5645 \[math-ph\]\]](#).

Master integrals for the ρ -parameter @ $O(\alpha_s^3)$

Example : One usually has **no** Gaussian differential equation, but something like Heun or more general, i.e. with more than 3 singularities.

$$\frac{d^2}{dx^2} f_{8a}(x) + \frac{9 - 30x^2 + 5x^4}{x(x^2 - 1)(9 - x^2)} \frac{d}{dx} f_{8a}(x) - \frac{8(-3 + x^2)}{(9 - x^2)(x^2 - 1)} f_{8a}(x) = l_{8a}(x)$$

Homogeneous solutions:

$$\begin{aligned}\psi_{1a}^{(0)}(x) &= \sqrt{2\sqrt{3}\pi} \frac{x^2(x^2 - 1)^2(x^2 - 9)^2}{(x^2 + 3)^4} {}_2F_1\left[\frac{4}{3}, \frac{5}{3}; z\right] \\ \psi_{2a}^{(0)}(x) &= \sqrt{2\sqrt{3}\pi} \frac{x^2(x^2 - 1)^2(x^2 - 9)^2}{(x^2 + 3)^4} {}_2F_1\left[\frac{4}{3}, \frac{5}{3}; 1 - z\right],\end{aligned}$$

with

$$z = z(x) = \frac{x^2(x^2 - 9)^2}{(x^2 + 3)^3}.$$

Use contiguous relations first to get into the ball-park. \implies at least two differently indexed ${}_2F_1$'s are going to appear. All classical ${}_2F_1$ wisdom is always applied first.

When can ${}_2F_1$ -Solutions be mapped to Complete Elliptic Integrals?




Figure 1: The transformation of special ${}_2F_1$ functions under the triangle group.

l	d	R	f
A	2	1	$4x(1-x)$
B	2	$(1-x)^{-1/6}$	$\frac{1}{4}x^2/(x-1)$
C	2	$(1-x)^{-1/8}$	$\frac{1}{4}x^2/(x-1)$
D	2	$(1-x)^{-1/12}$	$\frac{1}{4}x^2/(x-1)$
E	2	$(1-x/2)^{-1/2}$	$x^2/(x-2)^2$
F	3	$(1+3x)^{-1/4}$	$27x(1-x)^2/(1+3x)^3$
G	3	$(1+\omega x)^{-1/2}$	$1-(x+\omega)^3/(x+\bar{\omega})^3$
H	4	$(1-8x/9)^{-1/4}$	$64x^3(1-x)/(9-8x)^3$

Table: The functions R and f for the different hypergeometric transformations of degree d ; $\omega^2 + \omega + 1 = 0$.

$${}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix}; x \right] = R(x) {}_2F_1 \left[\begin{matrix} a', b' \\ c' \end{matrix}; f(x) \right]$$

Master integrals for the ρ -parameter @ $O(\alpha_s^3)$

$$\frac{d^2}{dx^2} f_{8a}(x) + \frac{9 - 30x^2 + 5x^4}{x(x^2 - 1)(9 - x^2)} \frac{d}{dx} f_{8a}(x) - \frac{8(-3 + x^2)}{(9 - x^2)(x^2 - 1)} f_{8a}(x) = l_{8a}(x)$$

Homogeneous solutions:

$$\psi_3^{(0)}(x) = -\frac{\sqrt{1 - 3x}\sqrt{x+1}}{2\sqrt{2\pi}} \left[(x+1)(3x^2+1)\mathbf{E}(z) - (x-1)^2(3x+1)\mathbf{K}(z) \right]$$

$$\psi_4^{(0)}(x) = -\frac{\sqrt{1 - 3x}\sqrt{x+1}}{2\sqrt{2\pi}} \left[8x^2\mathbf{K}(1-z) - (x+1)(3x^2+1)\mathbf{E}(1-z) \right],$$

$$z = \frac{16x^3}{(x+1)^3(3x-1)} \quad \text{[This function is not at all random! (see later).]}$$


\mathbf{K} , \mathbf{E} are the complete elliptic integrals of the 1st and 2nd kind.

$$\mathbf{K}(z) = \frac{2}{\pi} {}_2F_1 \left[\frac{1}{2}, \frac{1}{2}; 1; z \right], \quad \mathbf{E}(z) = \frac{2}{\pi} {}_2F_1 \left[\frac{1}{2}, -\frac{1}{2}; 1; z \right]$$

l_{8a} contains rational functions of x and HPLs.

Solutions with a Singularity

Homogeneous Solution:



Inhomogeneous Solution

$$\psi(x) = \psi_3^{(0)}(x) \left[C_1 - \int dx \psi_4^{(0)}(x) \frac{N(x)}{W(x)} \right] + \psi_4^{(0)}(x) \left[C_2 - \int dx \psi_3^{(0)}(x) \frac{N(x)}{W(x)} \right]$$



$C_{1,2}$: from physical boundary conditions.

Series Solution

$$\begin{aligned}
 f_{8a}(x) = & -\sqrt{3} \left[\pi^3 \left(\frac{35x^2}{108} - \frac{35x^4}{486} - \frac{35x^6}{4374} - \frac{35x^8}{13122} - \frac{70x^{10}}{59049} - \frac{665x^{12}}{1062882} \right) + \left(12x^2 - \frac{8x^4}{3} \right. \right. \\
 & \left. \left. - \frac{8x^6}{27} - \frac{8x^8}{81} - \frac{32x^{10}}{729} - \frac{152x^{12}}{6561} \right) \operatorname{Im} \left[\operatorname{Li}_3 \left(\frac{e^{-i\pi/6}}{\sqrt{3}} \right) \right] \right] - \pi^2 \left(1 + \frac{x^4}{9} - \frac{4x^6}{243} - \frac{46x^8}{6561} \right. \\
 & \left. - \frac{214x^{10}}{59049} - \frac{5546x^{12}}{2657205} \right) - \left(-\frac{3}{2} - \frac{x^4}{6} + \frac{2x^6}{81} + \frac{23x^8}{2187} + \frac{107x^{10}}{19683} + \frac{2773x^{12}}{885735} \right) \psi^{(1)} \left(\frac{1}{3} \right) \\
 & - \sqrt{3} \pi \left(\frac{x^2}{4} - \frac{x^4}{18} - \frac{x^6}{162} - \frac{x^8}{486} - \frac{2x^{10}}{2187} - \frac{19x^{12}}{39366} \right) \ln^2(3) - \left[33x^2 - \frac{5x^4}{4} - \frac{11x^6}{54} \right. \\
 & \left. - \frac{19x^8}{324} - \frac{751x^{10}}{29160} - \frac{2227x^{12}}{164025} + \pi^2 \left(\frac{4x^2}{3} - \frac{8x^4}{27} - \frac{8x^6}{243} - \frac{8x^8}{729} - \frac{32x^{10}}{6561} - \frac{152x^{12}}{59049} \right) \right. \\
 & \left. + \left(-2x^2 + \frac{4x^4}{9} + \frac{4x^6}{81} + \frac{4x^8}{243} + \frac{16x^{10}}{2187} + \frac{76x^{12}}{19683} \right) \psi^{(1)} \left(\frac{1}{3} \right) \right] \ln(x) + \frac{135}{16} + 19x^2 \\
 & - \frac{43x^4}{48} - \frac{89x^6}{324} - \frac{1493x^8}{23328} - \frac{132503x^{10}}{5248800} - \frac{2924131x^{12}}{236196000} - \left(\frac{x^4}{2} - 12x^2 \right) \ln^2(x) \\
 & - 2x^2 \ln^3(x) + O(x^{14} \ln(x))
 \end{aligned}$$

The solution can be easily extended to accuracies of $O(10^{-30})$ using Mathematica or Maple.

Solutions with a Singularity



Non-iterative Iterative Integrals

A New Class of Integrals in QFT:

$$\mathbb{H}_{a_1, \dots, a_{m-1}; \{a_m; F_m(r(y_m))\}, a_{m+1}, \dots, a_q}(x) = \int_0^x dy_1 f_{a_1}(y_1) \int_0^{y_1} dy_2 \dots \int_0^{y_{m-1}} dy_m f_{a_m}(y_m) \\ \times F_m[r(y_m)] H_{a_{m+1}, \dots, a_q}(y_{m+1}), \\ F[r(y)] = \int_0^1 dz g(z, r(y)), \quad r(y) \in \mathbb{Q}[y],$$

In general, this spans all solutions and the story would end here.

May be, most of the practical physicists, would led it end here anyway.

This type of solution applies to many more cases beyond ${}_2F_1$ -solutions (if being properly generalized).

[JB, ICMS 2016, July 2016, Berlin]

If one has no elliptic solution, one has to see, what else one has, and whether these cases are known mathematically as closed form solutions, with which properties etc. etc.

In the elliptic case we proceed as follows.

η -Ratios

Map:

$$x \rightarrow q : q = \exp[-\pi\mathbf{K}(1 - z(x))/\mathbf{K}(z(x))] := \exp[i\pi\tau], \quad |q| < 1$$

$$\eta(\tau) = q^{\frac{1}{12}} \prod_{k=1}^{\infty} (1 - q^{2k})$$

$$\prod_{l=1}^m \eta^{n_l}(l\tau) = \frac{1}{\eta^k(\tau)} \mathcal{M}, \quad n_l \in \mathbb{Z}$$

- ▶ Every η -ratio can be separated into a modular form \mathcal{M} and a factor $\eta^{-k}(\tau)$. [Algorithm 1]
- ▶ For the η -ratio \mathcal{M} is given as a polynomial of (generalized) Lambert-Eisenstein series. [Algorithm 2]
- ▶ All \mathcal{M} can be mapped into polynomials out of $\ln(q)$, $\text{Li}_0(q^j)$, and elliptic polylogarithms (of higher weight and also with indices depending on q).

Elliptic Polylogarithms as a Frame

$$\text{ELi}_{n,m}(x; y; q) = \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \frac{x^k}{k^n} \frac{y^l}{l^m} q^{kl}.$$

Weinzierl et al.:

$$\bar{E}_{n,m}(x; y; q) = \begin{cases} \frac{1}{i} [\text{ELi}_{n,m}(x; y; q) - \text{ELi}_{n,m}(x^{-1}; y^{-1}; q)], & n+m \text{ even} \\ \text{ELi}_{n,m}(x; y; q) + \text{ELi}_{n,m}(x^{-1}; y^{-1}; q), & n+m \text{ odd.} \end{cases}$$

Multiplication:

$$\text{ELi}_{n_1, \dots, n_l; m_1, \dots, m_l; 0, 2\sigma_2, \dots, 2\sigma_{l-1}}(x_1, \dots, x_l; y_1, \dots, y_l; q) = \text{ELi}_{n_1; m_1}(x_1; y_1; q)$$

$$\text{ELi}_{n_2, \dots, n_l; m_2, \dots, m_l; 2\sigma_2, \dots, 2\sigma_{l-1}}(x_2, \dots, x_l; y_2, \dots, y_l; q),$$

$$\begin{aligned} \text{ELi}_{n_1, \dots, n_l; m_1, \dots, m_l; 2\sigma_1, \dots, 2\sigma_{l-1}}(x_1, \dots, x_l; y_1, \dots, y_l; q) &= \sum_{j_1=1}^{\infty} \dots \sum_{j_l=1}^{\infty} \sum_{k_1=1}^{\infty} \dots \sum_{k_l=1}^{\infty} \frac{x_1^{j_1}}{j_1^{n_1}} \dots \frac{x_l^{j_l}}{j_l^{n_l}} \frac{y_1^{k_1}}{k_1^{m_1}} \frac{y_l^{k_l}}{k_l^{m_l}} \\ &\times \frac{q^{j_1 k_1 + \dots + j_l k_l}}{\prod_{i=1}^{l-1} (j_i k_i + \dots + j_l k_l)^{\sigma_i}}, \quad l > 0. \end{aligned}$$

Synchronization: performed for Lambert-Eisenstein series $q^m \rightarrow q$.
Re-translate after this.

Elliptic Solutions and Analytic q -Series

Map:

$$x \rightarrow q : q = \exp[-\pi \mathbf{K}(1 - z(x)) / \mathbf{K}(z(x))], \quad |q| < 1$$

- ▶ One attempts to calculate the integrals of the inhomogeneous solution in terms of q -series analytically.
- ▶ It is expected to write it in terms of products (and integrals over) elliptic polylogarithms [and possibly other functions].
- ▶ Note that the corresponding results are rather deep multi-series!
- ▶ Inspiration from algebraic geometry.

Elliptic polylogarithm (as a partly suitable frame):

$$\text{ELi}_{n,m}(x, y, q) = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \frac{x^j}{j^n} \frac{y^k}{k^m} q^{jk}$$

Is it (and its generalizations) a modular form ?

⇒ The central functions turn out to be more special ones.

The Individual Steps: from IBPs to Closed Form q -Series

- ▶ Generate the master integrals, determine their hierarchy, and look whether you have only 1st order factorization or also **2nd order terms**
- ▶ The latter can be trivial in case; check whether they persist in **Mellin space**
- ▶ If yes, analyze the **2nd order differential equation**
- ▶ One usually finds a ${}_2F_1$ -solution with rational argument $r(z)$, where $r(z)$ has additional singularities, i.e. the problem is of 2nd order, but has **more than 3 singularities**.
- ▶ Triangle group relations may be used to map the ${}_2F_1$ depending on the rational parameters a, b, c to the complete elliptic integrals **or not**.
- ▶ In the latter case return to the formalism on slide 13 and stop.
- ▶ If yes, one may walk along the **q -series avenue**.
- ▶ Different Levels of Complexity:
 - ▶ 1st order factorization in Mellin space:

$$M[K(1-z)](N) = \frac{2^{4N+1}}{(1+2N)^2 \binom{2N}{N}^2}$$

$$M[E(1-z)](N) = \frac{2^{4N+2}}{(1+2N)^2(3+2N) \binom{2N}{N}^2}$$

The Individual Steps: from IBPs to Closed Form q -Series

- ▶ Criteria by Herfurtnner (1991), Movasati et al. (2009) are obeyed.
 \implies 2-loop sunrise and kite diagrams, cf. Weinzierl et al. 2014-17.
Only $\mathbf{K}(r(z))$ and $\mathbf{K}'(r(z))$ contribute as elliptic integrals.
- ▶ Also $\mathbf{E}(r(z))$ and $\mathbf{E}'(r(z))$, square roots of quadratic forms etc. contribute (present case)
- ▶ Transform now: $x \rightarrow q$.
- ▶ The kinematic variable x :

$$k^2 = \frac{-x^3}{(1+x)^3(1-3x)} = \frac{\vartheta_2^4(q)}{\vartheta_3^4(q)}$$
$$x = \frac{\vartheta_2^2(q)}{3\vartheta_2^2(q^3)}, \quad \text{i.e. } x \in [1, +\infty[$$

by a cubic transformation (Legendre-Jacobi).

[see also Borwein, Borwein: AGM; and Broadhurst (2008).]

$$x = \frac{1}{3} \frac{\eta^2(2\tau)\eta^2(3\tau)}{\eta^2(\tau)\eta^4(6\tau)}, \quad \text{singular, } \propto \frac{1}{q}$$

The Individual Steps: from IBPs to Closed Form q -Series

- ▶ Map to a Modular Form, which can be represented by Lambert Series
 - ▶ How to find the η -ratio ? \implies Many are listed as sequences in Sloan's OEIS.
 - ▶ To find a modular form, situated in a corresponding finite-dimensional vector space M_k one has to meet a series of conditions and usually split off a factor $1/\eta^k(\tau)$, $k > 0$.
 - ▶ The remainder modular form is now a polynomial over \mathbb{Q} of Lambert-Eisenstein series

$$\sum_{n=0}^{\infty} \frac{m^n q^{an+b}}{1 - q^{an+b}} \cdot$$

Example:

$$\mathbf{K}(z(x)) = \frac{\pi}{2} \sum_{k=1}^{\infty} \frac{q^k}{1 + q^{2k}}$$

- ▶ In this case, two q series are equal, if both are modular forms, and agree in a series of k first terms, where k is predicted for each congruence sub-group of $\Gamma(N)$.

The Individual Steps: from IBPs to Closed Form q -Series

- ▶ Map Lambert-Eisenstein Series into the frame of Elliptic Polylogarithms
- ▶ Examples:

$$\begin{aligned}\mathbf{K}(z) &= \frac{\pi}{2} \sum_{k=1}^{\infty} \frac{q^k}{1+q^{2k}} = \frac{\pi}{i} \sum_{k=1}^{\infty} [\text{Li}_0(iq^k) - \text{Li}_0(-iq^k)] \\ &= \frac{\pi}{4} \bar{E}_{0,0}(i, 1, q), \\ q \frac{\vartheta'_4(q)}{\vartheta_4(q)} &= -\frac{1}{2} [\text{ELi}_{-1;0}(1; 1; q) + \text{ELi}_{-1;0}(-1; 1; q)] \\ &\quad + [\text{ELi}_{0;0}(1; q^{-1}; q) + \text{ELi}_{0;0}(-1; q^{-1}; q)] \\ &\quad - [\text{ELi}_{-1;0}(1; q^{-1}; q) + \text{ELi}_{-1;0}(-1; q^{-1}; q)].\end{aligned}$$

- ▶ New type of elliptic polylogarithm, e.g.:
 $\text{ELi}_{-1;0}(-1; q^{-1}; q)$, $y = y(q)$!
- ▶ Argument synchronization necessary: $-q \rightarrow q$, $q^k \rightarrow q$ (cyclotomic).

Elliptic Solutions and Analytic q -Series

- ▶ Terms to be translated:
 - ▶ rational functions in x
 - ▶ \mathbf{K}, \mathbf{E}
 - ▶ $\sqrt{(1-3x)(1+x)}$
 - ▶ $H_{\bar{a}}(x)$

Examples:

$$\mathbf{E}(k^2) = \mathbf{K}(k^2) + \frac{\pi^2 q}{\mathbf{K}(k^2)} \frac{d}{dq} \ln [\vartheta_4(q)]$$

$$\mathbf{E}'(k^2) = \frac{\pi}{2\mathbf{K}(k^2)} \left[1 + 2 \ln(q) q \frac{d}{dq} \ln [\vartheta_4(q)] \right].$$

$$\begin{aligned} \frac{1}{\mathbf{K}(k^2)} &= \frac{2}{\pi \eta^{12}(\tau)} \left\{ \frac{5}{48} \left\{ 1 - 24 \text{ELi}_{0,-1}(1; 1; q) - 4 \left[1 - \frac{3}{2} \left[\text{ELi}_{0,-1}(1; 1; q) + \text{ELi}_{0,-1}(1; i; q) \right. \right. \right. \right. \\ &+ \left. \left. \left. \text{ELi}_{0,-1}(1; -1; q) + \text{ELi}_{0,-1}(1; -i; q) \right] \right\} \left\{ -1 + 4 \left[-\frac{1}{2} \left[\text{ELi}_{-2,0}(i; 1/q; q) \right. \right. \right. \right. \\ &+ \left. \left. \left. \text{ELi}_{-2,0}(-i; 1/q; q) \right] + \left[\text{ELi}_{-1,0}(i; 1/q; q) + \text{ELi}_{-1,0}(-i; 1/q; q) \right] - \frac{1}{2} \left[\text{ELi}_{0,0}(i; 1/q; q) \right. \right. \right. \end{aligned}$$

Elliptic Solutions and Analytic q -Series

$$\begin{aligned}
 & \left. \left. \left. + \text{ELi}_{0,0}(-i; 1/q; q) \right] \right] \right\} - \frac{1}{16} \left\{ 5 + 4 \left[-\frac{1}{2} \left[\text{ELi}_{-4,0}(i; 1/q; q) + \text{ELi}_{-4,0}(-i; 1/q; q) \right] \right. \right. \\
 & + 2 \left[\text{ELi}_{-3,0}(i; 1/q; q) + \text{ELi}_{-3,0}(-i; 1/q; q) \right] - 3 \left[\text{ELi}_{-2,0}(i; 1/q; q) \right. \\
 & \left. \left. + \text{ELi}_{-2,0}(-i; 1/q; q) \right] + 2 \left[\text{ELi}_{-1,0}(i; 1/q; q) + \text{ELi}_{-1,0}(-i; 1/q; q) \right] \right. \\
 & \left. \left. - \frac{1}{2} \left[\text{ELi}_{0,0}(i; 1/q; q) + \text{ELi}_{0,0}(-i; 1/q; q) \right] \right] \right\}
 \end{aligned}$$

$$\begin{aligned}
 H_{-1}(x) = \ln(1+x) &= -\ln(3q) - \bar{E}_{0,-1;2}(-1, -1; q) + \bar{E}_{0,-1;2}(\rho_6; -1; q) \\
 &\quad - \bar{E}_{0,-1;2}(\rho_3; -i; q) - \bar{E}_{0,-1;2}(\rho_3; i; q)
 \end{aligned}$$

$$H_1(x) = -H_{-1}(x)|_{q \rightarrow -q} + 2\pi i, \text{ etc.}; \quad \rho_m = \exp(2\pi i/m)$$

$$I(q) = \frac{1}{\eta^k(\tau)} \cdot \mathbf{P} \left[\ln(q), \text{Li}_0(q^m), \text{ELi}_{k,l}(x, y, q), \text{ELi}_{k',l'}(x, q^{-1}, q) \right]$$


$$\int \frac{dq}{q} I(q)$$

is usually not an elliptic polylogarithm, due to the η -factor, but a higher transcendental function in q .

We are still in the unphysical region and have to map back to $x \in [0, 1]$.

$$\Delta\rho = \frac{3G_F m_t^2}{8\pi^2\sqrt{2}} \left(\delta^{(0)}(x) + \frac{\alpha_s}{\pi} \delta^{(1)}(x) + \left(\frac{\alpha_s}{\pi} \right)^2 \delta^{(2)}(x) + \mathcal{O}(\alpha_s^3) \right)$$

$$\begin{aligned} \delta^{(2)}(x) = & \dots + C_F \left(C_F - \frac{C_A}{2} \right) \left[\frac{11-x^2}{12(1-x^2)^2} f_{8a}(x) + \frac{9-x^2}{3(1-x^2)^2} f_{9a}(x) + \frac{1}{12} f_{10a}(x) \right. \\ & \left. + \frac{5-39x^2}{36(1-x^2)^2} f_{8b}(x) + \frac{1-9x^2}{9(1-x^2)^2} f_{9b}(x) + \frac{x^2}{12} f_{10b}(x) \right] \\ & + \frac{C_F T_F}{9(1-x^2)^3} \left[(5x^4 - 28x^2 - 9) f_{8a}(x) + \frac{1-3x^2}{3x^2} (9x^4 + 9x^2 - 2) f_{8b}(x) \right. \\ & \left. + (9-x^2)(x^4 - 6x^2 - 3) f_{9a}(x) + \frac{1-9x^2}{3x^2} (3x^4 + 6x^2 - 1) f_{9b}(x) \right] \end{aligned}$$



For $x = 0$, this agrees with the result by Chetyrkin *et al* (and Avdeev *et al*), $\delta^{(2)}(0) = -3.9696$.

Problems related to more scales at 2 loops

Phase space integrals for:

- ▶ massive 2-loop pure-singlet (PS) Wilson coefficient in DIS:
scales: Q^2, m_c^2
- ▶ Drell-Yan QED initial state corrections to e^+e^- annihilation:
scales: $s' = sz, m_e^2$

An alphabet (PS case):

$$f_{W1(2)}(z) = \frac{1}{1 \mp kz},$$

$$f_{W3(4)}(z) = \frac{1}{\beta \pm z},$$

$$f_{W5(6)}(z) = \frac{1}{k \mp x - (1-x)kz},$$

$$f_{W7(8)}(z) = \frac{1}{k \mp x + (1-x)kz},$$

$$f_{W9}(z) = \frac{z}{k^2 (1 - z^2 (1 - x^2)) - x^2},$$

$$f_{W10}(z) = \frac{1}{z\sqrt{1-z^2}\sqrt{1-k^2z^2}},$$

$$f_{W11}(z) = \frac{z}{\sqrt{1-z^2}\sqrt{1-k^2z^2}},$$

$$f_{W12}(z) = \frac{z}{\sqrt{1-z^2}\sqrt{1-k^2z^2} (k^2 (1 - z^2 (1 - x^2)) - x^2)}.$$

Kummer-elliptic Integrals

Definition:

Iterated integrals of the alphabet \mathfrak{A}'

$$\mathfrak{A}' = \mathfrak{A} \cup \left\{ \frac{1}{\sqrt{1-t^2}\sqrt{1-k^2t^2}}, \frac{t}{\sqrt{1-t^2}\sqrt{1-k^2t^2}}, \frac{\sqrt{1-k^2t^2}}{\sqrt{1-t^2}}, \right. \\ \left. \cup \left\{ \frac{1}{(t-a)\sqrt{1-t^2}\sqrt{1-k^2t^2}} \mid a \in \mathbb{C} \setminus \{\pm 1, \pm \frac{1}{k}\} \right\} \right\}.$$

with $1/(x-a) \in \mathfrak{A}$, $a \in \mathbb{C}$, are called letters for **Kummer-elliptic** integrals.

The first integrals of the above letters are all logarithmic; further integrals from some level on are not. E.g.:

$$H_{12}(t) = \frac{\operatorname{arctanh} \left[\frac{k\sqrt{1-t^2}}{\sqrt{1-k^2t^2}} \right] - \operatorname{arctanh} \left[\frac{k}{z} \right]}{(1-k^2)kz}$$

The spanning basis is found

- ▶ by rationalizing all occurring integrals as widely as possible
- ▶ and by using a Risch-like algorithm afterwards.

The phase space integrals are now fully available in terms of iterative integrals.

The root-valued alphabet for $e^+e^- \rightarrow \gamma^*/z^*$ at $O(\alpha^2)$

$$d_1 = \frac{1}{\sqrt{1-t}\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_2 = \frac{1}{\sqrt{1-t}\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_3 = \frac{1}{t\sqrt{1-t}\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_4 = \frac{1}{(16\rho^2 + (4z - 8\rho(1+z))t + (1-z)^2t^2)\sqrt{1-t}\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_5 = \frac{1}{(16\rho^2 + (4z - 8\rho(1+z))t + (1-z)^2t^2)\sqrt{1-t}\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_6 = \frac{1}{(16\rho^2 + (4z - 8\rho(1+z))t + (1-z)^2t^2)\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_7 = \frac{1}{(16\rho^2 + (4z - 8\rho(1+z))t + (1-z)^2t^2)\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_8 = \frac{1-z}{(4\rho - (1-z)t)\sqrt{1-t}}$$

$$d_9 = \frac{1}{(16\rho^2 + 4(z - 2\rho(1+z))t + (1-z)^2t^2)\sqrt{1-t}}$$

$$d_{10} = \frac{1}{(16\rho^2 + 4(z - 2\rho(1+z))t + (1-z)^2t^2)\sqrt{1-t}}$$

$$d_{11} = \frac{1}{t\sqrt{16\rho^2 - 8\rho(1+z)t + (1-z)^2t^2}}$$

$$d_{12} = \frac{1}{16\rho^2 + 4(z - 2\rho(1+z))t + (1-z)^2t^2}$$

$$d_{13} = \frac{1}{16\rho^2 + 4(z - 2\rho(1+z))t + (1-z)^2t^2}$$

$$d_{14} = \frac{1}{t(1-z) - 4\rho}$$

$$d_{15} = \frac{1}{\sqrt{1-t}(t(1-z) - 4\rho)}$$

$$d_{16} = \frac{1}{\sqrt{t(1-t)}\sqrt{t(1-z)^2 - 16\rho^2}}$$

$$d_{17} = \frac{1}{\sqrt{t(1-t)}(t(1-z) - 4\rho)\sqrt{t(1-z)^2 - 16\rho^2}}$$

$$d_{18} = \frac{1}{\sqrt{t}\sqrt{t(1-z)^2 - 16\rho^2}}$$

$$d_{19} = \frac{1}{\sqrt{t}(t(1-z) - 4\rho)\sqrt{t(1-z)^2 - 16\rho^2}}$$

$$d_{20} = \frac{1}{\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{21} = \frac{1}{\sqrt{1-t}\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{22} = \frac{\sqrt{t}}{\sqrt{t(1-z)^2 - 16\rho^2}\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{23} = \frac{\sqrt{t}}{\sqrt{t(1-z)^2 - 16\rho^2}(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)}$$

$$d_{24} = \frac{1}{(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{25} = \frac{1}{(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{26} = \frac{1}{\sqrt{1-t}(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{27} = \frac{1}{\sqrt{1-t}(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{28} = \frac{1}{\sqrt{t}\sqrt{t(-1+z)^2 - 16\rho^2}\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{29} = \frac{1}{\sqrt{t}\sqrt{t(1-z)^2 - 16\rho^2}(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)}$$

$$d_{30} = \frac{1}{\sqrt{t}\sqrt{t(1-z)^2 - 16\rho^2}(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{31} = \frac{\sqrt{t}}{\sqrt{t(1-z)^2 - 16\rho^2}(t^2(1-z)^2 - 8\rho(1+z)t + 4zt + 16\rho^2)\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{32} = \frac{1}{t\sqrt{1-t}\sqrt{t(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$d_{33} = \frac{1}{\sqrt{1-t}\sqrt{t^2(1-z)^2 - 8\rho t(1+z) + 16\rho^2}}$$

$$v_1 = \frac{1}{\sqrt{1-4t}\sqrt{16t^2 - 8(1+z)t + (1-z)^2}}$$

$$v_2 = \frac{1}{t\sqrt{1-4t}\sqrt{16t^2 - 8(1+z)t + (1-z)^2}}$$

$$v_3 = \frac{1}{\sqrt{1-4t}(4t - (1+z))\sqrt{16t^2 - 8(1+z)t + (1-z)^2}}$$

$$v_4 = \frac{1}{t\sqrt{1-t}}$$

Kummer-elliptic integrals

Why are these representations important ?

- ▶ Complete and irreducible analytic results.
- ▶ Stable starting point for expansions in small ratios of invariants.
- ▶ Led to a clarification of the ISR $O(\alpha^2)$ corrections in e^+e^- annihilation recently. [Want to have this in correct form for new high-luminosity acceleration projects (ILC, Fcc_ee etc.).]

Expansions :

- ▶ Expansions in m_e^2/s or m_c^2/Q^2 lead to prefactors which are usual HPLs
- ▶ Possibility to map to Mellin- N space (e.g. for fast evolution programs).
- ▶ In case of more scales \rightarrow lower complexity (iterative) integrals

ISR corrections to e^+e^- annihilation




Figure: Relative deviations of the results of Berends et al. 1987 from the exact result in % for the $O(\alpha^2)$ corrections. The non-singlet contribution (process II): dash-dotted line; the pure singlet contribution (process III): dashed; the interference term between both contributions (process IV): dots; for $s = M_Z^2$, $M_Z = 91.1879$ GeV.

[1] Berends, Burgers, van Neerven, Nucl. Phys. B 297 (1988) 429 [deviations in all channels](#).

[2] [now confirmed for all channels](#): J. Blümlein, A. De Freitas and W.L. van Neerven, Nucl. Phys. B 855 (2012) 508.

Conclusions

- ▶ We have automated the chain from IBPs to 2nd order solutions within the theory of differential equations [Before we had solved the 1st order factorizing cases for whatsoever basis of MIs.]
- ▶ General solution in the case not 1st order factorizing: Non-iterative iterative integrals III.
- ▶ These solutions might be sufficient and are very precise numerically and the result has a compact representation.
- ▶ In the elliptic cases we were enforced to generalize to structures not yet appearing in the case of the sunrise/kite integrals.
- ▶ Modular forms need to become a manifest part of knowledge for particle physicist working on fundamental QFTs.
- ▶ Any η ratio. can be solved.

Conclusions

- ▶ The general solution is given in terms of polynomials of elliptic polylogarithms, **more precisely: Lambert-Eisenstein series** and a few simpler functions in q -space
- ▶ **What comes next ?** Abel integrals ? K3 surfaces (Kummer, Kähler, Kodaira), Calabi-Yau structures...?
- ▶ Phase space integrals usually lead to **incomplete elliptic integrals** (and related more complex functions). Their occurrence is triggered by **more scales**.
- ▶ These structures form iterative integrals.
- ▶ Expanding in appropriate scale-ratios leads to simpler structures down to HPLs and may be used for numerical representations (in some cases).

Publications: Physics

JB, A. De Freitas, S. Klein, W.L. van Neerven, Nucl. Phys. B755 (2006) 272
I. Bierenbaum, JB, S. Klein, Nucl. Phys. B780 (2007) 40; Nucl.Phys. B820 (2009) 417; Phys.Lett. B672 (2009) 401
JB, S. Klein, B. Tödttli, Phys. Rev. D80 (2009) 094010
I. Bierenbaum, JB, S. Klein, C. Schneider, Nucl. Phys. B803 (2008) 1
J. Ablinger, JB, S. Klein, C. Schneider, F. Wißbrock, Nucl. Phys. B844 (2011) 26
JB, A. Hasselhuhn, S. Klein, C. Schneider, Nucl. Phys. B866 (2013) 196
J. Ablinger et al., Nucl. Phys. B864 (2012) 52; Nucl. Phys. B882 (2014) 263; Nucl. Phys. B885 (2014) 409; Nucl. Phys. B885 (2014) 280; Nucl. Phys. B886 (2014) 733; Nucl.Phys. B890 (2014) 48
A. Behring et al., Eur.Phys.J. C74 (2014) 9, 3033; Nucl. Phys. B897 (2015) 612; Phys. Rev. D92 (2015) 11405
JB, G. Falcioni, A. De Freitas Nucl. Phys. B910 (2016) 568.
J. Ablinger et al., Nucl.Phys. B922 (2017) 1
J. Ablinger et al., Nucl.Phys. B921 (2017) 585

Publications: Mathematics

JB, S. Kurth, Phys. Rev. D **60** (1999) 014018
JB, Comput. Comput.Phys.Comm. 133 (2000) 76
JB, Comput. Phys. Commun. 159 (2004) 19
JB, Comput. Phys. Commun. 180 (2009) 2143; 0901.0837
JB, D. Broadhurst, J. Vermaseren, Comput. Phys. Commun. 181 (2010) 582
JB, M. Kauers, S. Klein, C. Schneider, Comput. Phys. Commun. 180 (2009) 2143
JB, S.Klein, C. Schneider, F. Stan. J. Symbolic Comput. 47 (2012) 1267
J. Ablinger, JB, C. Schneider, J. Math. Phys. 52 (2011) 102301, J. Math. Phys. 54 (2013) 082301
J. Ablinger, JB, 1304.7071 [Contr. to a Book: Springer, Wien]
J. Ablinger, JB, C. Raab, C. Schneider, J. Math. Phys. 55 (2014) 112301
J. Ablinger, A. Behring, JB, A. De Freitas, A. von Manteuffel, C. Schneider, Comp. Phys. Commun. 202 (2016) 33
JB, C. Schneider, Phys. Lett. B 771 (2017) 31
A. Ablinger et al., DESY 16-147, arXiv:1706.01299
JB, M. Round, C. Schneider arXiv:1706.03677 [cs.SC]
J. Ablinger, JB, P. Marquard, N. Rana, C. Schneider, Nucl.Phys. B939 (2019) 253
J. Ablinger, JB, M. Round, C. Schneider, arXiv:1809.07084.