

Search for Bound States in Doubly Heavy Tetraquarks with Lattice QCD

Martin Pflaumer

pflaumer@itp.uni-frankfurt.de

Goethe-Universität Frankfurt am Main
in collaboration with Luka Leskovec, Stefan Meinel, Marc Wagner

The 14th International Workshop on Heavy Quarkonium
March 15, 2021



Motivation (1)

Experimental background

- Experimentally observed states $Z_b(10610)^+$ and $Z_b(10650)^+$
- Mass suggests a bottomonium state $\bar{b}b$ but would be electrically neutral
⇒ Quantum numbers can be described with four-quark structure

Motivation (1)

Experimental background

- Experimentally observed states $Z_b(10610)^+$ and $Z_b(10650)^+$
- Mass suggests a bottomonium state $\bar{b}b$ but would be electrically neutral
 \Rightarrow Quantum numbers can be described with four-quark structure

Theoretical study

- We study similar but less challenging systems
- Quark content: $\bar{Q}\bar{Q}'qq'$, here: $\bar{b}\bar{b}ud$, $\bar{b}\bar{b}us$, $\bar{b}\bar{c}ud$
- In the limit $m_Q \rightarrow \infty$ stable tetraquark was shown

[J. Carlson, L. Heller and J. A. Tjon, Phys. Rev. D **37**, 744 (1988)]

[A. V. Manohar and M. B. Wise, Nucl. Phys. B **399**, 17 (1993)]

[E. J. Eichten and C. Quigg, Phys. Rev. Lett. **119**, no. 20, 202002 (2017)]

Motivation (2)

Born-Oppenheimer study of doubly-heavy tetraquarks:

- i.e. static heavy quarks (\bar{b} -quarks)
- Prediction of a **bound tetraquark** in $\bar{b}\bar{b}ud$ sector with $I(J^P) = 0(1^+)$ and $M_{\bar{b}\bar{b}ud} - (M_B + M_{B^*}) \approx -90 \text{ MeV}$

[Z. S. Brown and K. Orginos, Phys. Rev. D **86**, 114506 (2012)]

[P. Bicudo *et al.* [ETMC], Phys. Rev. D **87**, no. 11, 114511 (2013)]

[P. Bicudo, K. Cichy, A. Peters, B. Wagenbach, M. Wagner, Phys. Rev. D **92**, no. 1, 014507 (2015)]

[P. Bicudo, J. Scheunert and M. Wagner, Phys. Rev. D **95**, no. 3, 034502 (2017)]

Motivation (2)

Born-Oppenheimer study of doubly-heavy tetraquarks:

- i.e. static heavy quarks (\bar{b} -quarks)
- Prediction of a **bound tetraquark** in $\bar{b}\bar{b}ud$ sector with $I(J^P) = 0(1^+)$ and $M_{\bar{b}\bar{b}ud} - (M_B + M_{B^*}) \approx -90 \text{ MeV}$

[Z. S. Brown and K. Orginos, Phys. Rev. D **86**, 114506 (2012)]

[P. Bicudo *et al.* [ETMC], Phys. Rev. D **87**, no. 11, 114511 (2013)]

[P. Bicudo, K. Cichy, A. Peters, B. Wagenbach, M. Wagner, Phys. Rev. D **92**, no. 1, 014507 (2015)]

[P. Bicudo, J. Scheunert and M. Wagner, Phys. Rev. D **95**, no. 3, 034502 (2017)]

- Evidence for a $\bar{b}\bar{b}ud$ **resonance** in the $I(J^P) = 0(1^-)$ channel with $M_{\bar{b}\bar{b}ud} - (M_B + M_B) \approx +20 \text{ MeV}$, $\Gamma \approx 100 \text{ MeV}$

[P. Bicudo, M. Cardoso, A. Peters, M.P. and M. Wagner, Phys. Rev. D **96**, no. 5, 054510 (2017)]

Motivation (3)

Searching for doubly-heavy tetraquark **bound states** in full lattice QCD using **Non-Relativistic QCD**:

- i.e. \bar{b} -quarks are treated non-relativistically.
- Expecting bound state for $\bar{b}\bar{b}ud$ in the $I(J^P) = 0(1^+)$ channel
 - Improve computation by more elaborated operators
- Previous studies predict a bound state for $\bar{b}\bar{b}us$ with $I(J^P) = \frac{1}{2}(1^+)$
 - Expected to be weaker bound than $\bar{b}\bar{b}ud$
- Status for $\bar{b}\bar{c}ud$ with $I(J^P) = 0(1^+)$ and $0(0^+)$ more unclear
 - Might be weakly bound or no binding

[A. Francis, R. J. Hudspith, R. Lewis and K. Maltman, Phys. Rev. Lett. **118**, no. 14, 142001 (2017)]

[P. Junnarkar, N. Mathur and M. Padmanath, Phys. Rev. D **99**, no. 3, 034507 (2019)]

[A. Francis, R. J. Hudspith, R. Lewis and K. Maltman, Phys. Rev. D **99**, no. 5, 054505 (2019)]

[L. Leskovec, S. Meinel, M.P. and M. Wagner, Phys. Rev. D **100**, no.1, 014503 (2019)]

[R. J. Hudspith, B. Colquhoun, A. Francis, R. Lewis and K. Maltman, Phys. Rev. D **102**, 114506 (2020)]

[M.P., L. Leskovec, S. Meinel and M. Wagner, arXiv:2009.10538 [hep-lat]].

Interpolating Operators for $\bar{b}b u d$

- Relevant thresholds are BB^* and B^*B^* (≈ 45 MeV heavier)
- Two types of interpolating operators:

Interpolating Operators for $\bar{b}b u d$

- Relevant thresholds are BB^* and B^*B^* (≈ 45 MeV heavier)
- Two types of interpolating operators:
 - **Local operators:**
 - Four quarks at the same space-time position
 - 3 operators: BB^* , B^*B^* , diquark-antidiquark

Interpolating Operators for $\bar{b}b u d$

- Relevant thresholds are BB^* and B^*B^* (≈ 45 MeV heavier)
- Two types of interpolating operators:
 - **Local operators:**
 - Four quarks at the same space-time position
 - 3 operators: BB^* , B^*B^* , diquark-antidiquark
 - **Nonlocal operators:**
 - Two mesons separated in space-time position
 - 2 operators: BB^* and B^*B^* scattering operators

Interpolating Operators for $\bar{b}b u d$


- Relevant thresholds are BB^* and B^*B^* (≈ 45 MeV heavier)
 - Two types of interpolating operators:
 - **Local operators:**
 - Four quarks at the same space-time position
 - 3 operators: BB^* , B^*B^* , diquark-antidiquark
 - **Nonlocal operators:**
 - Two mesons separated in space-time position
 - 2 operators: BB^* and B^*B^* scattering operators
 - Expectation:
 - **Local operators:** good overlap to **ground state** (stable four-quark)
 - **Nonlocal operators:** sizable overlap to **first excited state** (2 meson state)
- \Rightarrow Isolate ground state from higher excitations, especially first excited state

Energy Spectrum for the $\bar{b}\bar{b}ud$ system

- Due to point-to-all propagators, only 5×3 correlation matrix available (no scattering operator at source)
- Apply **multi-exponential matrix fitting**: employable also for non-symmetric matrices


$$C_{jk}(t) \approx \sum_{n=0}^{N-1} Z_j^n Z_k^n e^{-E_n t},$$

E_n : n -th energy eigenvalue
 $Z_j^n = \langle \Omega | \mathcal{O}_j | n \rangle$: overlap factor



Schematic representation of Wick contractions for different correlation matrix elements

Fit Results for $\bar{b}b\bar{u}d$




Results for the lowest two $\bar{b}b\bar{u}d$ energy levels relative to the BB^* threshold. Black box: local operator included. Red box: scattering operator included.

Overlap Factors

For fixed j : Z_j^n indicates relative importance of energy eigenstates $|n\rangle$

$$\mathcal{O}_j^\dagger|\Omega\rangle = \sum_{n=0}^{\infty} |n\rangle\langle n|\mathcal{O}_j^\dagger|\Omega\rangle = \sum_{n=0}^{\infty} Z_j^n |n\rangle.$$



The normalized overlap factors $|\tilde{Z}_j^n|^2 = \frac{|Z_j^n|^2}{\max_m (|Z_j^m|^2)}$ as determined on ensemble C005.

Scattering Analysis


- Relate *finite volume* energy spectrum E_n to *infinite volume* scattering amplitude
- Use Lüscher's formula to determine phase shift and *infinite volume* binding energy
- Confirmation that ground state is stable tetraquark.

Scattering Analysis and Chiral Extrapolation

Scattering Analysis

- Relate *finite volume* energy spectrum E_n to *infinite volume* scattering amplitude
- Use Lüscher's formula to determine phase shift and *infinite volume* binding energy
- Confirmation that ground state is stable tetraquark.

Chiral Extrapolation



Fit of the pion-mass dependence of E_{binding} . The vertical dashed line indicates the physical pion mass.

$$E_{\text{binding}}(m_{\pi,\text{phys}}) = (-128 \pm 24 \pm 10) \text{ MeV}$$

$$m_{\text{tetraquark}}(m_{\pi,\text{phys}}) = (10476 \pm 24 \pm 10) \text{ MeV}$$

Interpolating Operators for $\bar{b}\bar{b}us$ and $\bar{b}\bar{c}ud$

Interpolating Operators for $\bar{b}\bar{b}us$

$$I(J^P) = \frac{1}{2}(1^+)$$

relevant thresholds	B^*B_s, BB_s^* (\approx equal), $B^*B_s^*$ ($\approx +45$ Mev)
---------------------	--

local operators	$B^*B_s, BB_s^*, B^*B_s^*$, diquark-antidiquark
-----------------	--

nonlocal operators	$B^*B_s, BB_s^*, B^*B_s^*$
--------------------	----------------------------

Interpolating Operators for $\bar{b}b_{us}$ and $\bar{b}c_{ud}$

Interpolating Operators for $\bar{b}b_{us}$

$$I(J^P) = \frac{1}{2}(1^+)$$

relevant thresholds	B^*B_s, BB_s^* (\approx equal), $B^*B_s^*$ ($\approx +45$ Mev)
---------------------	--

local operators	$B^*B_s, BB_s^*, B^*B_s^*$, diquark-antidiquark
-----------------	--

nonlocal operators	$B^*B_s, BB_s^*, B^*B_s^*$
--------------------	----------------------------

Interpolating Operators for $\bar{b}c_{ud}$

$$I(J^P) = 0(1^+)$$


$$I(J^P) = 0(0^+)$$

relevant thresholds	B^*D, BD^* ($\approx +95$ Mev), B^*D^* ($\approx +140$ Mev)	$BD,$ B^*D^* ($\approx +185$ Mev)
---------------------	--	---

local operators	B^*D, BD^* , diquark-antidiquark	$BD,$ diquark-antidiquark
-----------------	---------------------------------------	------------------------------



nonlocal operators	BD^*, B^*D	BD
--------------------	--------------	------

Preliminary Results for $\bar{b}\bar{b}us$



- Found evidence for bound state with $E_{\text{binding}} \approx -80$ MeV
- Not affected by finite volume effects

Preliminary Results for $\bar{b}\bar{c}ud$



left: $\bar{b}\bar{c}ud$, $J = 0$. right: $\bar{b}\bar{c}ud$, $J = 1$.

- No evidence for bound states in $\bar{b}\bar{c}ud$ systems
- Lowest energy level corresponds to threshold

Summary

- Study bound states in doubly heavy tetraquarks
- Consider *local* and *nonlocal* interpolating operators
- Apply a *finite volume Lüscher analysis*

Summary

- Study bound states in doubly heavy tetraquarks
- Consider *local* and *nonlocal* interpolating operators
- Apply a *finite volume Lüscher analysis*
- Predict a **bound state** in the $\bar{b}\bar{b}ud$ channel with in $I(J^P) = 0(1^+)$ with $E_{\text{binding}} = (-128 \pm 24 \pm 10) \text{ MeV}$
- Evidence for **bound state** in $\bar{b}\bar{b}us$, $I(J^P) = \frac{1}{2}(1^+)$ sector with $E_{\text{binding}} \approx -80 \text{ MeV}$
- No evidence for bound tetraquark in $\bar{b}\bar{c}ud$, both $0(1^+)$ and $0(0^+)$

Summary

- Study bound states in doubly heavy tetraquarks
- Consider *local* and *nonlocal* interpolating operators
- Apply a *finite volume Lüscher analysis*
- Predict a **bound state** in the $\bar{b}\bar{b}ud$ channel with in $I(J^P) = 0(1^+)$ with $E_{\text{binding}} = (-128 \pm 24 \pm 10) \text{ MeV}$
- Evidence for **bound state** in $\bar{b}\bar{b}us$, $I(J^P) = \frac{1}{2}(1^+)$ sector with $E_{\text{binding}} \approx -80 \text{ MeV}$
- No evidence for bound tetraquark in $\bar{b}\bar{c}ud$, both $0(1^+)$ and $0(0^+)$

Outlook

- Finalize evaluation of $\bar{b}\bar{b}us$ and $\bar{b}\bar{c}ud$ outcomes
- More detailed analysis of threshold states in $\bar{b}\bar{c}ud \rightarrow$ candidates for resonances?

Summary

- Study bound states in doubly heavy tetraquarks
- Consider *local* and *nonlocal* interpolating operators
- Apply a *finite volume Lüscher analysis*
- Predict a **bound state** in the $\bar{b}\bar{b}ud$ channel with in $I(J^P) = 0(1^+)$ with $E_{\text{binding}} = (-128 \pm 24 \pm 10) \text{ MeV}$
- Evidence for **bound state** in $\bar{b}\bar{b}us$, $I(J^P) = \frac{1}{2}(1^+)$ sector with $E_{\text{binding}} \approx -80 \text{ MeV}$
- No evidence for bound tetraquark in $\bar{b}\bar{c}ud$, both $0(1^+)$ and $0(0^+)$

Outlook

- Finalize evaluation of $\bar{b}\bar{b}us$ and $\bar{b}\bar{c}ud$ outcomes
- More detailed analysis of threshold states in $\bar{b}\bar{c}ud \rightarrow$ candidates for resonances?

Thank You for Your Attention!

Lattice Setup

- Use gauge link configuration generated by RBC and UKQCD collaboration

[Y. Aoki *et al.* [RBC and UKQCD Collaborations], Phys. Rev. D **83**, 074508 (2011)]

[T. Blum *et al.* [RBC and UKQCD Collaborations], Phys. Rev. D **93**, no. 7, 074505 (2016)]

- 2 + 1 flavours **domain-wall fermions** and Iwasaki gauge action
- Five different ensembles which differ in

lattice spacing $a \approx 0.083 \text{ fm} \dots 0.114 \text{ fm}$,

lattice size $L \approx 2.65 \text{ fm} \dots 5.48 \text{ fm}$,

pion mass $m_\pi \approx 139 \text{ MeV} \dots 431 \text{ MeV}$


\Rightarrow explore dependence on L, m_π

- Smear **point-to-all propagators** for the up and down quarks

Scattering Analysis

- Relate *finite volume* energy spectrum E_n to *infinite volume* scattering amplitude for 2 energy levels in T_1^+ irrep
- Use Lüscher's formula and scattering momenta k_n^2 to determine phase shift
- Apply effective-range-expansion (ERE)

$$k \cot \delta_0(k) = \frac{1}{a_0} + \frac{1}{2}r_0k^2 + \mathcal{O}(k^4).$$



Plot of the effective-range-expansion for C005.
Blue curve: $ak \cot(\delta(k)) + |ak|$.
Vertical green line: Inelastic B^*B^* threshold


Scattering Analysis

- Relate *finite volume* energy spectrum E_n to *infinite volume* scattering amplitude for 2 energy levels in T_1^+ irrep
- Use Lüscher's formula and scattering momenta k_n^2 to determine phase shift
- Apply effective-range-expansion (ERE)

$$k \cot \delta_0(k) = \frac{1}{a_0} + \frac{1}{2}r_0k^2 + \mathcal{O}(k^4).$$

- Search bound state pole of scattering amplitude below threshold at

$$\cot \delta_0(k_{\text{BS}}) = i, \quad \text{so:} \quad -|k_{\text{BS}}| = \frac{1}{a_0} - \frac{1}{2}r_0|k_{\text{BS}}|^2$$



Scattering Analysis


- Relate *finite volume* energy spectrum E_n to *infinite volume* scattering amplitude for 2 energy levels in T_1^+ irrep
- Use Lüscher's formula and scattering momenta k_n^2 to determine phase shift
- Apply effective-range-expansion (ERE)

$$k \cot \delta_0(k) = \frac{1}{a_0} + \frac{1}{2}r_0k^2 + \mathcal{O}(k^4).$$

- Search bound state pole of scattering amplitude below threshold at


$$\cot \delta_0(k_{\text{BS}}) = i, \quad \text{so:} \quad -|k_{\text{BS}}| = \frac{1}{a_0} - \frac{1}{2}r_0|k_{\text{BS}}|^2$$

- Results essentially identical to the finite-volume energy levels
- Confirmation that ground state is stable tetraquark.



Plot of the effective-range-expansion for C005.
Blue curve: $ak \cot(\delta(k)) + |ak|$.
Vertical green line: Inelastic B^*B^* threshold

Comparison of Different Results for $\bar{b}\bar{b}ud$



Comparison of $\bar{b}\bar{b}ud$ tetraquark binding energies with $I(J^P) = 0(1^+)$ (black: this work; blue: lattice NRQCD; red: lattice QCD computations of static $\bar{b}\bar{b}$ potentials and solving the Schrödinger equation; green: effective field theories and potential models).