Thermal photon measurements with the future MPD experiment at NICA

D. Ivanishchev, D. Kotov, E. Kryshen, M. Malaev, V. Riabov, Yu. Riabov
(NRC «Kurchatov Institute» - PNPI) for the MPD collaboration



This work was supported by RFBR according to the research project № 18-02-40045

Outline

- Heavy ion collisions at NICA
- Motivation for thermal photons study in heavy-ion collisions
- Photon conversion method
- Feasibility studies for thermal photons measurement in the MPD
- Conclusions and outlook

Heavy ion collisions at NICA





- A mega-science project NICA, Dubna, JINR
- Modernization of existing Nuclotron facility
- Parameters:
  - ✓ relativistic ions up to Au, $\sqrt{s_{NN}}$ = 4-11 GeV
  - ✓ polarized p and d, $\sqrt{s_{NN}}$ = 27 GeV (for p)
  - ✓ luminosity 10^{27} $cm^{-2}s^{-1}$
- Working experiment: BM@N (fixed target)
- Experiments under construction: MPD, SPD (collider)

- Study of the phase diagram in the region of high baryonic density and intermediate temperatures
- Extension of modern heavy-ion programs at RHIC and the LHC to lower energies

Thermal radiation in heavy ion collisions





- Photons leave the medium without interaction
- Black body radiation: inverse slope proportional to T_{eff}

Thermal photons from QGP



Naïve:



Hot medium
Large yield

Thermal photons from hadron gas





Warm medium
Moderate yield

Photon spectra at RHIC and LHC

PHENIX (AuAu @ 200 GeV)



T_{eff} = 239 ± 25 (stat) ± 7 (syst) MeV

PHENIX: Phys. Rev. Lett. 104 (2010) 132301

ALICE (PbPb @ 2760 GeV)



T_{eff} = 297 ± 12 (stat) ± 41 (syst) MeV

ALICE: Phys. Lett. B754 (2016) 235

Effective temperature vs energy



J. Phys.: Conf. Ser. 1070 (2018) 012012

Challenge: decay photons

Inclusive photon spectra are dominated by decay photons

$$R_\gamma = \gamma_{\text{inc}} / \gamma_{\text{decay}}$$



Relative contributions of different hadrons to the total decay photon spectrum as a function of the decay photon transverse momentum



ALICE: Phys. Lett. B754 (2016) 235

Photon reconstruction: two methods

- Electromagnetic calorimeters
  - Efficient at $p_T > 2$ GeV/c
  - Hardware trigger capabilities

- Photon conversion $\gamma \rightarrow e^+e^-$ in the material
  - $P = 1 - \exp(-7/9\, x/X_0)$
  - Efficient at $0.5 < p_T < 4$ GeV/c
  - Much better resolution at low p_T







MPD experiment at NICA



- CMS Energy: 4-11 GeV
- Design luminosity: 10^{27} cm^{-1} s^{-1}
- Stage 1: TPC, TOF, ECAL, FHCal, FFD
- Stage 2: + ITS + EndCap

MPD: NIM A 628 (2011) 99-102

Photon conversion centers





Main conversion structures in Stage 1:

- Beam pipe: 0.3% X_0
- Inner TPC barrel structures: 2.4% X_0

Future:

- Inner tracking system
- Dedicated photon converter (cylindrical metal pipe) under investigation

Conversion reconstruction efficiency



Typical cuts on electrons:

- $|\eta|<1$
- $p_T > 50$ MeV/c
- at least 20 hits in TPC
- +/-4σ electron PID in TPC/TOF

Typical cuts on ee pair:

- Small DCA ($\chi^2 < 10$)
- Vertex R > 10 cm
- Direction to vertex:
  $\theta < \exp(-2.777-2.798 * p_T) + 0.0175$
- $m_{ee} < 22.6 + 17.4 * p_T$
- ee plane orientation wrt B:
  $\Psi_{Pair} < 0.1$ rad

- Studied with MPDROOT Stage 1 setup
- Using MpdParticle to build secondary vertices
- Cuts optimized to maximize signal significance
- Contribution of (non-gamma) background < 10-20%
  - can be further improved with tighter cuts



UrQMD and PHSD predictions at NICA energies



- UrQMD and PHSD generators: good agreement in neutral meson cross sections

Neutral meson reconstruction





- Using 20M minimum bias URQMD events
- Pion signal is clearly visible in a wide p_T range
- Statistics not enough to study eta reconstruction







Neutral meson reconstruction efficiency



- Embedding technique used to study reconstruction efficiency vs p_T
- 700 000 min. bias UrQMD events @ 11 GeV
- 500 π^0 + 500 η embedded with flat p_T distribution
- Neutral meson reconstruction efficiency ~ 10^{-4}
- π^0 peak is significantly narrower with conversion method compared to ECAL





Corrected π^0 spectra





- Efficiency-corrected spectra are extrapolated down to 0 p_T:
  - Tsallis function
  - Two-component model (Bylinkin, Rostovtsev)
  - Hagedorn function
- The obtained fits cab be used to calculate photon spectra from π^0 decays
- Extrapolation uncertainties are significant only at low photon p_T < 0.3 GeV/c

Integrated direct photon yield

No reliable predictions for photons in UrQMD/PHSD -> using data driven method





Phys. Rev. Lett. 123, 022301

The integrated direct photon yield:

- scales as $(dN_{ch}/d\eta)^{1.25}$ in a wide range of multiplicities/collision energies
- the scaling is violated in small collision systems / small multiplicities ($dN_{ch}/d\eta < 20$)
- AA yield is a factor of ~10 larger than the N_{coll}-scaled yield in pp

Assuming this scaling still holds at lower energies, we can expect universal multiplicity scaling for

- (0-60)% centralities at 11 GeV
- (0-40)% centralities at 4 GeV

p_T-differential direct photon yields



PHENIX, PRL 123, 022301
Khachatryan, Praszałowicz EPJC 80 (2020) 670



thermal | power law

- Universal scaling of p_T-differential direct photon yields at moderate p_T is observed at RHIC/LHC
- It can be used to predict p_T spectra of direct photons at NICA energies for $p_T > 0.6$ GeV/c
- Switch to thermal spectrum at $p_T < 0.6$ GeV/c: $dN/dp_T \sim p_T \exp(-p_T/T)$
- Using conservative effective temperature T = 150 MeV (see e.g. *PRC* 93 (2016) 054901)

Reconstructed direct photons





- Reconstructed direct photon spectra can be obtained from predicted direct photon yields multiplied by the photon reconstruction efficiency

Inclusive photon spectra and R_γ ratio

$$R_\gamma = \frac{\gamma_{\rm inc}}{\gamma_{\rm decay}} = \frac{\gamma_{\rm inc}/\pi^0}{\gamma_{\rm decay}/\pi^0_{\rm param}}$$

Inclusive photon spectrum was simulated as a sum of direct and decay photon spectra

Excess over 1 shows the fraction of direct photons





- Systematic uncertainties on R_γ can be reduced to ~5%
- Conclusion: direct photon yields can be extracted with good accuracy down to low p_T

$$\gamma_{\rm direct} = \left(1 - \frac{1}{R_\gamma}\right) \cdot \gamma_{\rm inc}$$

Conclusions and outlook

- Photons are valuable probes of dense hadronic matter produced in heavy ion collisions
- Photon conversion method is a powerful tool to measure photon and neutral meson spectra
- Reconstruction of thermal photon yields looks promising at MPD
- Next:
  - Evaluate perspectives of thermal photon flow measurements
  - Feasibility studies on the dedicated converter and Stage 2 setup

BACKUP

Electron PID





TPC dE/dx: 0.2 - 0.5 GeV/*c*

TOF: p_T < 0.3 GeV/*c*

Selecting tracks in +/-4σ around the mean value:





Optimization of electron pair selection cuts





Fit functions for pt distributions



- Two-component model (Bylinkin, Rostovtsev):

$$E\frac{\mathrm{d}^3\sigma}{\mathrm{d}p^3} = A_\mathrm{e} \exp \frac{(M - \sqrt{p_\mathrm{T}^2 + M^2})}{T_\mathrm{e}} + A\left(1 + \frac{p_\mathrm{T}^2}{n_\mathrm{br} T^2}\right)^{-n_{br}}$$

- Hagedorn fit:

$$\frac{dN}{dy dp_\mathrm{T}} = p_\mathrm{T}\ A\ \left(\exp\left(-a * p_\mathrm{T} - |b| p_\mathrm{T}^2\right) + \frac{p_\mathrm{T}}{p_0}\right)^{-n}$$

- Tsallis fit:

$$E\frac{d^3\sigma}{dp^3} = \frac{1}{2\pi}\frac{d\sigma}{dy}\frac{(n-1)(n-2)}{(nT)^2}\left(1 + \frac{m_T}{nT}\right)^{-n}$$