LXX International conference "NUCLEUS –2020. Nuclear physics and elementary particle physics. Nuclear physics technologies"

Contribution ID: 86

Type: Oral report

MICROSCOPIC DESCRIPTION OF ISOSCALAR GIANT MONOPOLE RESONANCE IN $^{118-132}\mathrm{Sn}$

Monday, 12 October 2020 15:55 (25 minutes)

The study of nuclear giant resonances has long been a subject of extensive theoretical and experimental research. The multipole response of nuclei far from the β -stability line and the possible occurrence of exotic modes of excitation present a growing field of research. In particular, the study of the isoscalar giant monopole resonances (ISGMR) in neutron-rich nuclei is presently an important problem not only from the nuclear structure point of view [1] but also because of the special role they play in many astrophysical processes such as prompt supernova explosions [2] and the interiors of neutron stars [3]. One of the successful tools for describing the ISGMR is the quasiparticle random phase approximation (QRPA) with the self-consistent mean-field derived from Skyrme energy density functionals (EDF) [4]. Such an approach can describe the properties of the low-lying states reasonably well by using existing Skyrme interactions. Due to the anharmonicity of the vibrations there is a coupling between one-phonon and more complex states [5]. The main difficulty is that the complexity of calculations beyond standard QRPA increases rapidly with the size of the configuration space, and one has to work within limited spaces. Using a finite rank separable approximation for the residual particle-hole interaction derived from the Skyrme forces one can overcome this numerical problem [6-8]. In the present report, we analyze the effects of phonon-phonon coupling (PPC) on the E0 strength distributions of neutron-rich tin isotopes. Using the same set of the EDF parameters we describe available experimental data for ^{118,120,122,124}Sn [9] and give prediction for ^{130,132}Sn [10]. The effects of the PPC leads to a redistribution of the main monopole strength to lower energy states and also to higher energy tail.

- 1. J.P. Blaizot // Phys. Rep. 1980. V. 64. P. 171.
- 2. H.A. Bethe // Rev. Mod. Phys. 1990. V. 62. P. 801.
- 3. N.K. Glendenning // Phys. Rev. Lett. 1986. V. 57. P. 1120.
- 4. N. Paar, D. Vretenar, E. Khan, G. Colò // Rep. Prog. Phys. 2007. V. 70. P. 691.
- 5. V.G. Soloviev // Theory of Atomic Nuclei: Quasiparticles and Phonons. 1992. Bristol/Philadelphia.
- 6. N.V. Giai, Ch. Stoyanov, V.V. Voronov // Phys. Rev. C. 1998. V. 57. P. 1204.
- 7. A.P. Severyukhin, V.V. Voronov, N.V. Giai // Phys. Rev. C. 2008. V. 77. P. 024322.
- 8. A.P. Severyukhin, V.V. Voronov, N.V. Gia // Eur. Phys. J. A. 2004. V. 22. P. 397.
- 9. T. Li, U. Garg, Y. Liu et al. // Phys. Rev. C. 2010. V. 81. P. 034309.
- 10. N.N. Arsenyev, A.P. Severyukhin // in preparation.

Primary authors: Dr ARSENYEV, Nikolay (Joint Institute for Nuclear Research); Dr SEVERYUKHIN, Alexey (Joint Institute for Nuclear Research)

Presenter: Dr ARSENYEV, Nikolay (Joint Institute for Nuclear Research)

Session Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei

Track Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei.