

Study of ground states of $^{10,11}\text{B}$, $^{10,11}\text{C}$ nuclei by Feynman's continual integrals method

Tuesday, October 13, 2020 6:35 PM (20 minutes)

The wave functions of the ground states of few-body nuclei $^{10,11}\text{B}$, $^{10,11}\text{C}$ were calculated by Feynman's continual integrals method in Euclidean time [1–3]. The algorithm of parallel calculations was implemented in C++ programming language using NVIDIA CUDA technology [4]. Calculations were performed on the NVIDIA Tesla K40 accelerator installed within the heterogeneous cluster of the Laboratory of Information Technologies, Joint Institute for Nuclear Research, Dubna.

The studied isotopes are considered as cluster nuclei with the following configurations: ^{10}B ($2\alpha + n + p$), ^{11}B ($2\alpha + n + n + p$), ^{10}C ($2\alpha + p + p$) and ^{11}C ($2\alpha + n + p + p$). Results of the cluster model were compared with results of the shell model of deformed nuclei [5, 6].

References

1. R.P.Feynman and A.R.Hibbs. Quantum Mechanics and Path Integrals (McGraw-Hill, New York, 1965).
2. E.V.Shuryak and O.V.Zhirov // Nucl. Phys. B. 1984. V.242. P.393.
3. V.V.Samarin, M.A.Naumenko // Phys. Atom. Nucl. 2017. V.80. P.877.
4. M.A.Naumenko and V.V.Samarin // Supercomp. Front. Innov. 2016. V.3. P.80.
5. V.V.Samarin // Phys. Atom. Nucl. 2010. V.73. P. 1416.
6. V.V.Samarin // Phys. Atom. Nucl. 2015. V.78. P. 128.

Primary author: Prof. SAMARIN, Viacheslav (JINR)

Presenter: Prof. SAMARIN, Viacheslav (JINR)

Session Classification: Poster session 1 (part 1)

Track Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei.