

## ASTROPHYSICAL S-FACTOR OF THE DIRECT ${}^3\text{He}(\alpha,\gamma){}^7\text{Be}$ CAPTURE REACTION IN CLUSTER MODELS

Saturday 17 October 2020 11:45 (25 minutes)

The direct  ${}^3\text{He}(\alpha,\gamma){}^7\text{Be}$  radiative capture reaction is studied in the framework of two- and three-body potential cluster models [1,2]. E1 and E2 transitions are described at the long-wavelength approximation. The two-body model is based on a simple Gaussian form  ${}^3\text{He}$ -potential of Dubovichenko  $V_D^a$  from Ref.[1], with a modification in  $d$  waves. The new potential parameters are  $V_0=-180$  MeV,  $\alpha=0.4173$   $\text{fm}^{-2}$  and  $V_0=-190$  MeV,  $\alpha=0.4017$   $\text{fm}^{-2}$  in the  $d_{3/2}$  and  $d_{5/2}$  partial waves, respectively. The potential describes correctly the phase shifts in the  $s$ ,  $p$ ,  $d$  and  $f$  waves and binding energies of the ground  $p_{3/2}$  and the first excited  $p_{1/2}$  bound states. As can be seen in Fig.1, the modification of the potential in  $d$  waves allows to improve the description of the astrophysical S factor for the direct  ${}^3\text{He}(\alpha,\gamma){}^7\text{Be}$  radiative capture reaction at intermediate energies  $E>0.5$  MeV in comparison with the results of Ref.[1]. In the three-body model the  ${}^7\text{Be}$  nucleus is described as a bound state of  $\alpha+p+d$  in the Hyperspherical Lagrange mesh method. The initial state is factorized into the  $p+d$  bound state and the  $\alpha+{}^3\text{He}$  scattering state. The  $\alpha d$ -potential is from Ref.[2], while  $\alpha N$ -potential was taken from Ref.[3]. The  $pd$ -potential of the Gaussian form [4] with parameters  $V_0=-34.92$  MeV,  $\alpha=0.15$   $\text{fm}^{-2}$  and  $V_0=2.4$  MeV,  $\alpha=0.01$   $\text{fm}^{-2}$  are used in the even and odd partial waves, respectively. The  $\alpha{}^3\text{He}$ -potential is the same as in the two-body model. The three body bound state wave functions of  ${}^7\text{Be}$  was corrected at  $R=6$  fm with the help of the Whittaker asymptotics.

Fig. 1. (a) Astrophysical S factor within the two- and three-body models in comparison with the available experimental data. Panel (b) highlights the low energy region.

1. E.M. Tursunov et al. // Phys. Rev. C. 2018. V.97. id.035802.
2. E.M. Tursunov et al. // Phys. Rev. C. 2018. V.98. id.055803.
3. V.T. Voronchev et al. // Few-Body Syst. 1995. V. 18. p. 191.
4. S. Dubovichenko et al. // Phys. Elem. Part. At. Nucl. 1997. V.28. p.1529.

**Primary authors:** TURSUNOV, E.M. (Institute of Nuclear Physics, Academy of Sciences, 100214, Ulugbek, Tashkent, Uzbekistan); TURAKULOV, S.A. (Institute of Nuclear Physics, Academy of Sciences, 100214, Ulugbek, Tashkent, Uzbekistan; Tashkent Railway Engineering Institute, Uzbekistan); DUSNAZAROV, E.M. (Institute of Nuclear Physics, Academy of Sciences, 100214, Ulugbek, Tashkent, Uzbekistan)

**Presenter:** TURAKULOV, S.A. (Institute of Nuclear Physics, Academy of Sciences, 100214, Ulugbek, Tashkent, Uzbekistan; Tashkent Railway Engineering Institute, Uzbekistan)

**Session Classification:** Section 2. Experimental and theoretical studies of nuclear reactions

**Track Classification:** Section 2. Experimental and theoretical studies of nuclear reactions.