



Effect of detuning impedance on TMCI for constant wake and zero chroma

E. Métral and G. Rumolo

- ◆ Comparison between Burov-Danilov_1998 (extended to constant wake) and 2-particle model developed by GiovanniR (12/08/19)



Reminder on 2-particle model (GiovanniR, 12/08/19)







**Burov-Danilov_1998 extended to constant wake
=> EM's implementation... possible errors are
mine ;-)**



$$\kappa_x = D(z)/W_x(z)$$



+ 1 in x for CRW flat chamber
- 1/2 in y for CRW flat chamber



$\kappa = 0.0$ 



$\kappa = 0.1$ 



 $\kappa = 0.2$ 



$\kappa = 0.3$ 



$\kappa = 0.4$ 


$\kappa = 0.5$ 

$\kappa = 0.6$ 


$\kappa = 0.7$ 



$\kappa = 0.8$ 



$\kappa = 0.9$ 




$\kappa = 1.0$






$\kappa = 1.1$ 


$\kappa = 0.0$ 




$\kappa = -0.1$




 $\kappa = -0.2$ 





$\kappa = -0.3$







$\kappa = -0.4$








$\kappa = -0.5$







$\kappa = -0.6$




 $\kappa = -0.7$ 





$\kappa = -0.8$







$\kappa = -0.9$








$\kappa = -1.0$






$\kappa = -1.1$







Conclusion

- ◆ Good agreement between the 2 models and confirmation of the predictions from the 2-particle model (GiovanniR, 12/08/19)
- ◆ The air-bag model of these slides (Burov-Danilov_1998 extended to constant wake for zero chroma and considering only 2 modes, -1 and 0) is fully described by the following matrix to be diagonalized

$$\begin{pmatrix} -1 + \frac{\kappa}{2} I_{norm} & \frac{2 I_{norm}}{\pi^2} (1 - \kappa) \\ \frac{2 I_{norm}}{\pi^2} (-1 - \kappa) & \frac{I_{norm}}{2} (-1 + \kappa) \end{pmatrix}$$