

High p_T identified hadrons in large and small systems measured in PHENIX

Balázs Ujvári

University of Debrecen, on behalf of the PHENIX collaboration

March 4. 2020.



1 PHENIX

- Detector
- Collisions
- Hadron identification


2 High p_T hadrons in large system

- Mesons
- Baryons

3 High p_T hadrons in small system


- π^0 , Φ p+Au, d+Au, He+Au
- R_{dAu} of mesons + baryons

Detector




PHENIX, beam view


Collisions




Au+Au
 $N_{\text{part}}=148$, $N_{\text{coll}}=315$, $b=7.8$ fm



Cu+Au
 $N_{\text{part}}=147$, $N_{\text{coll}}=292$, $b=4.3$ fm



Au+Au
 $N_{\text{part}}=212$, $N_{\text{coll}}=523$, $b=6.2$ fm




Cu+Au
 $N_{\text{part}}=202$, $N_{\text{coll}}=431$, $b=1.5$ fm

Au+Au and Cu+Au collisions with comparable N_{part}


Beam-beam counters are used to determine the event centrality. The mean number of participating nucleons (N_{part}) and the binary collisions (N_{coll}) in various centrality intervals are estimated with a Glauber model Monte-Carlo simulation folded with the BBC response.

Hadron identification π^0 and η




Invariant mass distribution for gamma-gamma pair


Hadron identification K_S^0 and K^*



The K_S^0 mesons are reconstructed by combining the π^0 candidates in pairs within the same event.




The K^{*0} and \bar{K}^{*0} mesons are reconstructed via the $K^+\pi^-$ and $K^-\pi^+$




Invariant mass distributions obtained with the “two kaons PID” and “no PID” methods after subtraction of the combinatorial background estimated using the event-mixing technique. Plot on the top corresponds to integrated p_T range whereas plot on the bottom is for the range $2 < p_T$ (GeV/c) < 3 . The “no PID” spectrum is fit to the sum of a Breit-Wigner function convolved with a Gaussian function to account for the Φ signal, and a polynomial function to account for the residual background.

Mesons




π^0 invariant yield








η invariant yield

Meson R_{AA} Cu+Au




R_{AB} for K^* and Φ meson $>$ R_{AB} for π^0 and η at low p_T in central events
 Difference disappears at high p_T




R_{AA} for Φ , η and π^0 in Cu+Au in different centrality bins


Meson R_{AA} in U+U




Strange quark dependence also present in U+U collisions at low p_T



Nuclear modification factors of π^0 , η and K_S mesons as functions of p_T for different U+U centrality intervals.





Comparison of $\pi^0 R_{AB}$ measured in Cu+Au, Au+Au and Cu+Cu collisions. It suggests that π^0 suppression mostly depends on the energy density and size of the produced medium. Corona-effect?





Ratios of η and π^0 yields measured as a function of p_T in different centrality intervals.

The QGP does not affect the jet fragmentation into these light mesons.




Similar results in Au+Au.




R_{AA} of K_S in different collision systems.

Parameter	"Glauber 1"	"Glauber 2"
R (fm)	6.81	6.86
a (fm)	0.60	0.42
β_2	0.280	0.265
β_4	0.093	0



Enhancement of $(p + \bar{p})/2$ yield at moderate p_T

Baryons





R_{AA} of $(p + \bar{p})/2$ consistent across Cu+Au and Au+Au at similar N_{part}

Summary large systems



- 1 Strangeness dependence in R_{AA} at low p_T in central events
- 2 R_{AA} for all meson species converges at high p_T
- 3 Ratios of η and π^0 is similar to other large systems
- 4 Corona-effect is either non-existent or very small
- 5 Same baryon enhancement seen in Cu+Au and Au+Au

Small System









π^0 , Φ p+Au, d+Au



π^0 , Φ p+Au, d+Au, He+Au






Ordering:


$$R_{pAu} > R_{dAu} > R_{HeAu}$$

in 0-20% at intermediate p_T
range.


Hint of π^0 suppression in
central collisions.



$R_{pAu} \approx R_{dAu} \approx R_{HeAu}$
 Hint of π^0 enhancement
 in peripheral collisions
 at high p_T .




R_{dAu} of mesons



Light mesons R_{dAu} exhibit similar shape in the whole low p_T range, in contrast to large system at low p_T

R_{dAu} of mesons + baryons





Protons R_{dAu} shows enhancement at moderate p_T as in heavy-ion

Summary small systems


- 1 π^0, Φ $R_{pAu} > R_{dAu} > R_{HeAu}$ in 0-20% at intermediate p_T range
- 2 π^0, Φ $R_{pAu} \approx R_{dAu} \approx R_{HeAu}$ in peripheral collisions
- 3 Light mesons R_{dAu} exhibit similar shape in contrast to heavy-ion
- 4 Protons R_{dAu} shows enhancement at moderate p_T as in heavy-ion

The research was financed by the Thematic Excellence Programme of the Ministry for Innovation and Technology in Hungary (ED_18-1-2019-0028), within the framework of the Space Sciences thematic programme of the University of Debrecen

NKFIH, OTKA-131991



The η/π^0 and K_S/π^0 as a function of p_T for different U+U centrality intervals



The comparison of η/π^0 ratios obtained in U+U and Au+Au.