



Contribution ID: 140

Type: Talk

Exploring the deformation of nuclei with correlation between anisotropic flow and transverse momentum from STAR

Thursday 15 July 2021 15:10 (20 minutes)

In the relativistic heavy-ion collisions the mean transverse momentum ($\langle p_T \rangle$) and anisotropic flow (v_n , $n=2,3$) have been found to be tightly correlated with the size and initial geometry eccentricity of the produced fireball, respectively. It provides a novel tool to image the deformation of the atomic nuclei at extremely short time scale ($< 10^{-24}$ s).

In this talk, we present measurements of correlations between v_n and $\langle p_T \rangle$ by using the Pearson correlation coefficient ($\rho(v_n\{2\}^2, \langle p_T \rangle)$) as a function of multiplicity in Au+Au and U+U collisions at top RHIC energy. Unlike in Au+Au collisions, a sign-change behavior has been found for $\rho(v_2\{2\}^2, \langle p_T \rangle)$ in central U+U collisions due to nuclei deformations. While $\rho(v_3\{2\}^2, \langle p_T \rangle)$ has been found to be similar between two collision systems. Comparing with several model calculations in the ultra-central regions, such measurements will help us to constrain the quadrupole deformation parameter (β_2) of the atomic nuclei.

Preferred track

Collectivity & Multiple Scattering

Author: HUANG, Shengli (Stony Brook University)

Presenter: HUANG, Shengli (Stony Brook University)

Session Classification: Collectivity and multiple-scattering

Track Classification: Collectivity and multiple-scattering