

Contribution ID: 231

Type: Flash-talk and poster/pre-recorded talk

## Investigations of the longitudinal broadening of two-particle transverse momentum correlations from STAR

Thursday 15 July 2021 19:32 (2 minutes)

Two-particle transverse momentum correlator is a powerful technique for understanding the dynamics of relativistic heavy-ion collisions. Among these, the transverse momentum correlator  $G_2(\Delta\eta, \Delta\varphi)$  is of particular interest for its potential sensitivity to the shear viscosity per entropy density  $\eta/s$  of the quark-gluon plasma formed in heavy-ion collisions [1,2]. We used Au–Au at  $\sqrt{s_{NN}} = 200$  GeV data to investigate: (i) the self-correlations associated with the definition of the collision centrality [3], (ii) the longitudinal broadening of the  $G_2(\Delta\eta, \Delta\varphi)$  correlator, (iii) the long range azimuthal dependence of the  $G_2(\Delta\eta, \Delta\varphi)$ . We will present the centrality and event shape dependence of the longitudinal width  $\sigma(\Delta\eta)$  and the azimuthal harmonics  $a_n^{pT}$  of the  $G_2(\Delta\eta, \Delta\varphi)$  correlator. Our measurements are compared to similar LHC measurements as well as with calculations using the UrQMD, AMPT, and EPOS models [4]. These comparisons are expected to reflect the efficacy of the  $G_2(\Delta\eta, \Delta\varphi)$  correlator to differentiate among theoretical models as well as to constrain the  $\eta/s$ .

[1] S. Gavin and M. Abdel-Aziz, Phys. Rev. Lett. 97, 162302 (2006)

[2] V. Gonzalez, et al., Eur.Phys.J.C 81 (2021) 5, 465

[3] N. Magdy, et al., arXiv:2101.01555

[4] N. Magdy, et al., arXiv:2105.07912

### Preferred track

Collectivity & Multiple Scattering

**Primary author:** Dr ABDELRAHMAN, Niseem (University of Illinois at Chicago)

**Presenter:** Dr ABDELRAHMAN, Niseem (University of Illinois at Chicago)

**Session Classification:** Poster Session