Searching for Dark Photons at the LHeC & FCC-he

Zeren Simon Wang

1909.02312
work with M. D'Onofrio and O. Fischer

November 27, 2019
Sixth LHC LLP Community Workshop
Ghent University, Ghent, Belgium

asia pacific center for
theoretical physics

Motivation

- Dark Matter
- Hidden sector theories predict new particles to interact with the Standard Model (SM) field content via feebly coupled mediator particles
- Non-gravitational portals between the dark sector and the SM:
  - scalar: dark Higgs
  - pseudoscalar: axion-like particles
  - vector: dark photons
  - neutrino: heavy neutral leptons
- Studied at the HL-LHC, CEPC, FCC-ee/hh, beam-dump, and fixed-target experiments
- Sensitivity of electron-proton colliders may be unique: LHeC & FCC-he

The dark photon model

- Extend the SM gauge group by an additional (broken) gauge group: $\underline{U(1)_X}$

$$\mathcal{L} \supset -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} - \frac{\epsilon}{2}X_{\mu\nu}F^{\mu\nu}$$

- Applying a field re-definition to get rid of the kinetic mixing term $X_{\mu\nu}F^{\mu\nu}$

$$\mathcal{L} \supset -\sum_f \bar{f}\,\epsilon\, e\, q_f\, \not{A}'\, f$$

- Two parameters: mass $m_{\gamma'}$ and mixing ϵ
- <u>Production</u> and <u>decay</u> controlled by one parameter ϵ
- Focus on <u>MeV-GeV</u> mass range in this work

LHeC and FCC-he, and dark photon production

	E_e (GeV)	E_p (TeV)	$\sqrt{s}$ (TeV)	$\mathcal{L}_{\text{int}}$ (ab^{-1})
LHeC	60	7	1.3	1
FCC-he	60	50	3.5	3







DIS: $Q^2 \gg m_p^2 \simeq 1$ GeV^2

Signal estimation







$$N_{\rm dv}(\sqrt{s}, \mathcal{L}, m_X, \epsilon)$$
$$= \sigma(M, \epsilon)\, \mathcal{L} \times \int D(\vartheta, \gamma)\, P_{\rm dv}(x_{\rm min}(\vartheta), x_{\rm max}(\vartheta), \Delta x_{\rm lab}(\tau, \gamma))\, d\vartheta d\gamma$$

$$P_{\rm dv} = {\rm Exp}\left(\frac{-x_{\rm min}}{\Delta x_{\rm lab}}\right) - {\rm Exp}\left(\frac{-x_{\rm max}}{\Delta x_{\rm lab}}\right)$$

$$x_{\rm min} = 200\mu m, x_{\rm max} = \infty, \Delta x_{\rm lab} = \tau_{\rm lab}|\vec{v}| = \beta_{\gamma'}\gamma_{\gamma'}\,\tau c$$

	$\eta(e/\mu)$	η(jets)
LHeC	(-4.3,4.9)	(-5,5)
FCC-he	(-5.0,5.2)	(-5.5,5.5)

In practice, apply $|\eta| < 4.7$

Dark photon decays



[Raggi, Kozhuharov,'15]

$$\Gamma(\gamma' \to l^+ l^-) = \frac{1}{3}\,\alpha_{\rm QED}\, m_{\gamma'}\, \epsilon^2 \sqrt{1 - \frac{4m_l^2}{m_{\gamma'}^2}} \left(1 + \frac{2m_l^2}{m_{\gamma'}^2}\right)$$

$$\Gamma_{\rm total}(\gamma') = \frac{\Gamma(\gamma' \to e^- e^+)}{\mathrm{BR}(\gamma' \to e^- e^+)}$$

Background

- Real low-energy photons interacting with the detector material or the beam pipe $\rightarrow e^- e^+$ pairs
  - location of the secondary vertex coincides with the detector material or the beam pipe
  - easily rejected
- Long-lived hadrons such as K_S, K_L, and Λ
  - lifetime far away from IP (3cm, 15m, and 8cm)
  - hadronic activity is aligned with the proton beam and propagates mostly into the forward hemisphere of the detector
  - their primary decay channels are only marginally consistent with our signal signature
  - their masses are well known

Results I: 90% C.L. sensitivity reaches



Results II: 90% C.L. sensitivity reaches



Nbkg=0, signal efficiency = 100%
Nbkg=0, signal efficiency = 20%
Nbkg=100, signal efficiency = 100%
Nbkg=100, signal efficiency = 20%



Results III: Comparison



[Physics Briefing Book: Input for the European Strategy for Particle Physics Update 2020 [1910.11775]]

Summary

- Portals to the dark sector
- Vector portal and dark photon model
- Electron-proton colliders may explore a unique territory in the parameter space for MeV-GeV dark photons

Thank You!