

Searching for long-lived particles at the LHC: Sixth workshop of the LHC LLP Community



Contribution ID: 1

Type: **not specified**

Searching for Dark Photons at the LHeC and FCC-he

Wednesday, 27 November 2019 14:15 (15 minutes)

Extensions of the Standard Model (SM) gauge group with a new $U(1)_X$ predict an additional gauge boson. Through kinetic mixing with the SM photons featured by a coupling ϵ , the ensuing so-called dark photons γ' , which acquire mass as a result of the breaking of the gauge group $U(1)_X$, can interact with the SM field content. These massive dark photons can therefore decay to pairs of leptons, hadrons, or quarks, depending on their mass $m_{\gamma'}$. In this work, we discuss searches for dark photons in the mass range around and below one GeV at the LHeC and FCC-he colliders. The signal is given by the displaced decays of the long-lived dark photon into two charged fermions. We discuss the impact of conceivable irreducible (SM and machine-related) backgrounds and different signal efficiencies. Our estimates show that the LHeC and FCC-he can test a domain that is complementary to other present and planned experiments.

Primary authors: D'ONOFRIO, Monica (University of Liverpool (GB)); FISCHER, Oliver (Karlsruhe Institute of Technology); WANG, Zeren Simon (APCTP, Korea)

Presenter: WANG, Zeren Simon (APCTP, Korea)