

20th Conference of Czech and Slovak Physicists



Report of Contributions

Contribution ID: 1

Type: **not specified**

Registration

Monday 7 September 2020 08:00 (9 hours)

Contribution ID: 2

Type: **not specified**

Welcome

Monday 7 September 2020 10:00 (30 minutes)

Contribution ID: 3

Type: **not specified**

Welcome party

Monday 7 September 2020 19:00 (3 hours)

Contribution ID: 6

Type: **not specified**

**Conference dinner: 19:00 - 19:30 arrival | 19:30 first
toast | 21:30 end of catering | 22:00 restaurant closed**

Wednesday 9 September 2020 19:30 (2 hours)

Contribution ID: 7

Type: **not specified**

Conference closing

Thursday 10 September 2020 15:30 (5 minutes)

Surprise for the persistent individuals - goodbye drink
(foyer of building A - behind the lecture room T1)

Contribution ID: 11

Type: Oral

Magnonic Analog of Black/White-Hole Horizons in Superfluid $^3\text{He-B}$

Monday 7 September 2020 10:30 (45 minutes)

Súčasná technológia neumožňuje experimentálne študovať a overovať naše teoretické modely a predstavy o fundamentálnych fyzikálnych javoch prebiehajúcich v takých astro-fyzikálnych objektoch, akými sú čierne diery. Zatiaľ jediným spôsobom ako experimentálne verifikovať tieto modely (a predstavy) o fyzike čiernych dier je najšť laboratórny fyzikálny systém, ktorý s istými obmedzeniami umožňuje simuláciu vlastností čiernych dier. Cieľom prednášky je prezentovať ako teoretický model, tak aj experimentálne výsledky získané na novom fyzikálnom systéme umožňujúcom simuláciu vlastností horizontu udalostí v čiernych/bielych dierach. Modelový systém je založený na jave spinovej (magnónovej) supratekutosti v supratekutom $^3\text{He-B}$. Ako experimentálny nástroj, ktorý modeluje vlastnosti čiernych/bielych dier sme využili tzv. spinovo precesujúce vlny šíriace sa na pozadí spinového toku medzi dvoma Bose-Einsteinovými kondenzátmi magnónov vo forme homogénne precesujúcich domén. V súhlase s teoretickým modelom sme experimentálne ukázali formovanie a prítomnosť horizontu udalostí pre spinovo precesujúce vlny, vrátane efektu zosilnenia týchto vln. Vzhľadom k tomu, že experiment je realizovaný pri teplote blízko absolútnej nuly ($600\ \mu\text{K}$), prezentovaný modelový systém tak umožňuje aj štúdium spontánneho Hawkingovho žiarenia.

The current mankind technologies do not allow experimental study and thus verify our theoretical models and concepts on fundamental physical phenomena that take place in such astrophysical objects like the black-holes are. At present the only possibility how to verify experimentally these theoretical models on physics of the black-holes is to find a laboratory system which, obviously with some limitations, mimics some properties of the black holes. An aim of this lecture is to introduce and present as a theoretical model, so the experimental results of the experiment made in a limit of absolute zero temperature ($600\ \mu\text{K}$) studying the spin wave analogue of black/white hole horizon using a spin (magnonic) superfluidity in superfluid $^3\text{He-B}$. As an experimental tool simulating the properties of the black/white horizon we used the spin-precession waves propagating on the background of the spin supercurrents between two Bose-Einstein condensates of magnons in form of the homogeneously precessing domains. We provide experimental evidence of the white hole formation for spin precession waves in this system, together with observation of an amplification effect. Moreover, the estimated temperature of the spontaneous Hawking radiation in this system is about four orders of magnitude lower than the system's background temperature what makes it a promising tool to study the effect of spontaneous Hawking radiation.

Author: MARCEL ČLOVEČKO, EMIL GAŽO, MARTIN KUPKA, PETER SKYBA

Presenter: SKYBA P. (Centrum fyziky nízkých teplôt, Ústav experimentálnej fyziky, SAV, Košice)

Session Classification: Plenary Talks

Contribution ID: 12

Type: **Oral**

QUANTUM WALKS

Monday 7 September 2020 11:15 (45 minutes)

Quantum walks represent a generalization of the random walk to the coherent evolution of the quantum particle on a graph or a lattice. We provide an overview of the concept, focusing on the discrete-time quantum walk, and illustrate the applications in quantum simulations and quantum computation. Finally, we introduce a very successful experimental implementation of quantum walks which is based on photonic time-multiplexing, and review the recent experiments.

Author: ŠTEFANĀK M. (FJFI ČVUT v Praze)

Presenter: ŠTEFANĀK M. (FJFI ČVUT v Praze)

Session Classification: Plenary Talks

Contribution ID: 13

Type: **Oral**

EXPERIMENT KATRIN LIMITED THE NEUTRINO MASS TO LESS THAN 1 eV

Monday 7 September 2020 13:30 (45 minutes)

Neutrinos are together with photons the most abundant particles in the Universe. The knowledge of the neutrino mass is important not only for particle physics but also for cosmology. Oscillation experiments proved that at least two of the known three neutrino mass states have the mass different from zero, while at least one of these states has the mass greater than 0.05 eV. Precision spectroscopy of electrons emitted in the nuclear beta decay offers a principal possibility to determine the neutrino mass directly in a model independent way. However, only upper limits of the neutrino mass were determined until now. The most advanced Karlsruhe Tritium Neutrino Mass experiment (KATRIN) operates with a windowless gaseous source of molecular tritium. The energy of emitted electrons is analyzed with the electrostatic retarding spectrometer with magnetic adiabatic collimation. Recently, the first four-week science run of the KATRIN limited the neutrino mass to less than 1.1 eV. This is twice better than the previous limit achieved after several years of measurement. The KATRIN aims to the 0.2 neutrino mass sensitivity after 1000 days of measurements. Further on the hypothetical sterile neutrinos in the eV and keV mass ranges will be searched for.

Author: OTOKAR DRAGOUN FOR THE KATRIN COLLABORATION

Presenter: DRAGOUN O. (Nuclear Physics Institute of the Czech. Acad. Sci., Řež)

Session Classification: Plenary Talks

Contribution ID: 14

Type: **Oral**

CHARACTERIZATION AND BIOAPPLICATION OF MAGNETIC NANOPARTICLES

Monday 7 September 2020 14:15 (35 minutes)

Magnetic nanoparticles (NPs), especially SPIO nanoparticles, have long been used as MRI (Magnetic Resonance Imaging) contrast agents and as advantageous nanoplatforms for drug delivery, taking advantage of their unique magnetic properties and ability to function at the molecular and cellular level. In our centre smart design and development of SPIO nanoparticles in mild conditions with desired physico-chemical properties (size, composition, surface modification, magnetization, relaxivity, biocompatibility, etc.) have been performed to fulfil desired bioapplication. In this lecture the most promising results will be presented and consequently discussed advanced in vitro and in vivo MRI application of such prepared SPIO nanomaterials or nanocomposites. The particular attention will be devoted to labeling and MRI detection of stem cells, and potential clinical application of maghemite/bentonite nanocomposite as an per-oral contrast agent.

Author: POLÁKOVÁ K. (Regional Centre of Advanced Technologies and Materials, Palacký University in Olomouc)

Presenter: POLÁKOVÁ K. (Regional Centre of Advanced Technologies and Materials, Palacký University in Olomouc)

Session Classification: Plenary Talks

Contribution ID: 15

Type: **Oral**

NITRIDE SEMICONDUCTORS - PROPERTIES AND APPLICATIONS

Monday 7 September 2020 14:50 (35 minutes)

Why are the III-nitride semiconductors so unique that they became the second most important semiconductor material after silicon? Why it was so difficult to prepare high quality nitride layers? It was so difficult that scientist, which have partly solved this problem were five years ago awarded by Nobel Prize. What are the most perspective applications for nitrides? What problems have to be solved in nitride technology? Answers to these four questions concerning nitride properties, applications, technology and open problems will be presented in our contribution. We will show how the piezoelectric field in nitride heterostructures can be utilized in transistors with high electron mobility, but on the other how it can complicate the carrier confinement in quantum wells and design of optoelectronic applications. Application in lasers and light emitting diodes, in high electron mobility transistors (HEMTs) and in scintillators will be discussed with their advantages and limitations. We will touch also the most important technological problems and challenges, such as high dislocation density and possible ways to suppress it, the p-type doping of AlGa_N layers, problems with limited indium atom incorporation. Results of scintillator structure development in our group with the world record InGa_N QW number will be presented.

Author: ALICE HOSPODKOVÁ, TOMÁŠ HUBÁČEK, JIŘÍ OSWALD, FRANTIŠEK HÁJEK, KARLA KULDOVÁ, AND EDUARD HULICIUS

Presenter: HOSPODKOVÁ A. (Institute of Physics CAS, Prague)

Session Classification: Plenary Talks

Contribution ID: 16

Type: **Oral**

Role of photovoltaic solar cells in the ongoing energy systems transformation

Monday 7 September 2020 16:00 (35 minutes)

Photovoltaic solar energy conversion years has become a decisive part of the electricity generation. The global installed photovoltaic (PV) capacity surpassed 600 GW in 2020 and the terawatt installed capacity is going to be achieved within the following decade. The dramatic fall of costs made PV the cheapest source of new electricity generation (without subsidies) in large part of the world. PV exhibited ~25 % annual growth rate during the last 5 years. PV produces about 3 % of electricity globally, about 5% in the EU and it is expected to reach 15 % share of EU electricity generation by 2030.

PV is dominated by silicon wafer-based cells for which previously unforeseen low system prices have been reached. The advantages of photovoltaics based on silicon thin film (lower consumption of energy-costly silicon and thus shorter energy payback time) have not been sufficient to overcome the disadvantage of lower efficiencies. Thus the development and production of silicon thin film photovoltaics nearly stopped. The other types of thin film based PV (CdTe, CIS) as well as concentrated PV approaches struggle as well. The record efficiency of silicon based cell reached 26.7 % by using the silicon heterojunction (SHJ) with interdigitated back contacts composed of intrinsic and either n or p-type hydrogenated amorphous silicon layers with thicknesses ~ 10 nm. The record SHJ cell has been fabricated by photolithography, unsuitable for mass production. A simpler stencil mask based technology has been developed in the NextBase H2020 project. The tunnel-IBC approach further simplifies the process by eliminating the patterning and aligning the hole collector contacts. These advances offer a chance for a restart of photovoltaic production in Europe.

Innovation goes on to further enhance light-to-power conversion efficiency but this is not possible for single junction silicon based cells, as they are close to the fundamental Shockley Queisser limit (~29.6 %). Much higher efficiencies are possible with multijunction designs, but so far they have been demonstrated using III-V semiconductors, which are not suitable to large scale deployment. The next step in PV evolution will be the dual junction (tandem) cells of silicon with a suitable partner. The most perspective is the combination of perovskite –silicon tandem cells, with potential efficiency above 45% and current research results reaching 29%.

Author: FEJFAR A. (Institute of Physics, Czech Academy of Sciences, Prague)

Presenter: FEJFAR A. (Institute of Physics, Czech Academy of Sciences, Prague)

Session Classification: Plenary Talks

Contribution ID: 17

Type: **Oral**

ORGANIC SEMICONDUCTORS

Monday 7 September 2020 16:35 (35 minutes)

Organic semiconductors are a broad family of organic molecular materials that exhibit certain properties similar to inorganic semiconductors. Interestingly, molecular materials show only weak forces between the molecules which gives an opportunity of new deposition methods. The low-temperature evaporation in a vacuum is applicable for various small molecules, whereas “wet technologies” using organic material solubility in solvents are famous of large molecules and polymers. The thin-film fabrication technology is not only unique property of these materials; organic semiconductors exhibit semiconducting properties even without any doping. Hence, the organic semiconductors do not represent only alternative semiconducting materials, but it is also an exciting challenge for electronics and device physics. It has been found that the semiconducting properties have different microscopic origin than the inorganic materials even though the macroscopic behaviour is sometimes almost identical. Interestingly, in contrast with very first ideas researcher commented that “organic semiconductors” can be misleading term and suggests to call these materials “organic semi-insulators” but it is too late for such a correction. Actually, it is fascinating to state “Yes, it works, but we are still not sure why”.

Author: WEIS M.**Presenter:** WEIS M.**Session Classification:** Plenary Talks

Contribution ID: 18

Type: Oral

TIME-RESOLVED OPTICAL AND TERAHERTZ STUDIES OF THIN-FILM FERROMAGNETS AND ANTIFERROMAGNETS

Monday 7 September 2020 17:10 (35 minutes)

Recent breakthroughs in electrical detection and manipulation of antiferromagnets have opened a new avenue in the research of non-volatile spintronic devices. Antiparallel spins in antiferromagnets lead to the insensitivity to magnetic field perturbations, multi-level stability and ultra-fast spin dynamics, which is also a major experimental challenge [1]. In this contribution we show how the know-how that we achieved in the pump-probe study of spin transport in non-magnetic heterostructures AlGaAs/GaAs [2], diluted ferromagnetic semiconductor (Ga,Mn)As [3] or terahertz magneto-resistance [4] can be transferred to the research of antiferromagnetic metal CuMnAs. In particular, we show the latest progress in studies of magnetic anisotropy [5], dynamics of heat dissipation [6] and even switching of memory devices by single femtosecond laser pulse [7] or terahertz excitation [8].

[1] P. Němec et al., Nature Physics 14, 229 (2018)

[2] L. Nádvořník et al., Scientific Reports 6, 22901 (2016).

[3] P. Němec et al., Nature Physics 8, 411 (2012)

[4] L. Nádvořník et al., submitted to Nature Communications.

[5] V. Saidl, et al., Nature Photonics 11, 91 (2017)

[6] M. Surýnek et al., J. Appl. Phys. 127, 233904 (2020)

[7] Z. Kašpar et al., arXiv 1909.09071

[8] J. Heitz et al., submitted to Nano Letters.

Presenter: NÁDVORNÍK L. (Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Plenary Talks

Contribution ID: 19

Type: **Oral**

ELECTROMAGNETIC STRUCTURE OF MESONS AND BARYONS

Tuesday 8 September 2020 08:30 (45 minutes)

Present state of elementary particles to be specified by a set of quantum numbers is summarized. An existence of four types of interactions between elementary particles is reminded with an emphasis that not all particles are able of all four interactions. Those to be able of strong interactions are named "hadrons" - compound of "quarks". Compound from quark-antiquark are "mesons" and compound from three quarks are baryons. Recently four, even five quarks hadrons are confirmed experimentally, not to be in contradiction with QCD. Nevertheless X,Y,Z particles are out of any classification into multiplets to be known for standard mesons and baryons. Hadrons in interaction with charged leptons manifest space structure to be called electromagnetic (EM) structure, first time revealed for protons before the quark model of hadrons was established. It is described by "EM form factors (FFs)" the number of which depends on spin of hadron. The Unitary and Analytic model of FFs is presented to be based on analyticity and existence of vector-meson-resonances with quantum numbers of photon. Model depends on some unknown coupling constant ratios. Therefore it is applicable only to hadrons for which some experimental data exist. As a result only the EM structure of the nonet of pseudoscalar mesons and the $\frac{1}{2}^+$ octet baryons is completely described in the framework of this model up to now.

Author: C.ADAMUŠČÍN, E.BARTOŠ, S.DUBNIČKA, A.LIPTAJ, A.Z.DUBNIČKOVÁ

Presenter: DUBNIČKA S (Institute of Physics, Slovak Academy of Sciences, Bratislava)

Session Classification: Plenary Talks

Contribution ID: 20

Type: **Oral**

SELECTED RESULTS OF THE ATLAS EXPERIMENT WITH SIGNIFICANT CONTRIBUTION OF CZECH AND SLOVAK PHYSICISTS

Tuesday 8 September 2020 09:15 (35 minutes)

The ATLAS experiment is the largest particle physics detector installed at the Large Hadron Collider at CERN, which measures the proton-proton (or lead-lead) collisions at the world-highest energies. The precision measurements of the physics processes and particles' properties probe our current theory of elementary particles, the so-called Standard Model (SM). The discovery of the Higgs boson in 2012 opened a new territory to be explored - its detailed properties and interactions. Another exciting goal for now and the future is the search for new physics phenomena behind the SM.

The ATLAS Collaboration produces plenty of physics results every year. This talk concentrates on recent analyses with significant contribution of Czech and Slovak physicist participating in this experiment, for instance the measurement of selected Higgs boson decays, top-quark physics, rare decays of B-mesons and heavy-ion physics.

Author: DAVIDEK, Tomas (Charles University (CZ))

Presenter: DAVIDEK, Tomas (Charles University (CZ))

Session Classification: Plenary Talks

Contribution ID: 21

Type: **Oral**

Explosions in the Universe and the Origin of Chemical Elements

Tuesday 8 September 2020 10:20 (45 minutes)

Recent detection of gravitational waves from merging black holes and neutron stars opened a new window for studying the Universe and further accelerated the tremendous progress in multiwavelength time-domain astronomy. This talk will review exciting new developments in the quest for uncovering the origin of chemical elements in the Universe and understanding the birth of neutron stars and black holes. These major unsolved astrophysical problems are associated with some of the most luminous explosions and involve a wide range of physics such as nuclear matter at high densities, neutrinos, strong magnetic fields, and formation of dust and molecules.

Author: PEJCHA O. (Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles university)

Presenter: PEJCHA O. (Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles university)

Session Classification: Plenary Talks

Contribution ID: 22

Type: **Oral**

TEN YEARS OF COMPASS TOKAMAK OPERATION

Tuesday 8 September 2020 11:05 (35 minutes)

The COMPASS tokamak, originally constructed and operated at UKAEA in UK, has been reinstalled at the Institute of Plasma Physics of the CAS in 2009 and experienced a fruitful decade of scientific exploration. In this contribution, we will summarize the main achievements, which have contributed to the physics understanding of tokamak plasmas and development of nuclear fusion related technologies. These include: (i) investigation of the power decay lengths in limited plasmas, which directly impacted the design of the ITER inner wall plasma-facing components, (ii) error field (EF) experiments using the unique set of the tokamak high-field-side error field coils, (iii) power exhaust studies, (iv) runaway electron characterization and mitigation programme, and others.

COMPASS tokamak will be shut down before the end of 2020 to make space for a completely new machine: COMPASS Upgrade. Characteristics of this challenging device will be presented.

Author: M. KOMM ON BEHALF OF THE COMPASS TEAM

Presenter: KOMM M. (Institute of Plasma Physics of the CAS, Prague)

Session Classification: Plenary Talks

Contribution ID: 23

Type: **Oral**

Czech participation in new European Large Research Infrastructure FAIR

Tuesday 8 September 2020 11:40 (20 minutes)

Czech Republic is aspirant member of the Facility for Antiproton and Ion Research (FAIR) since March 2019. Unique new European Large Research Infrastructure FAIR is currently built up at the campus of German national laboratory for heavy ion research GSI Darmstadt. It will cover research in nuclear and hadron physics as well as in associated fields. It is included in the European Strategy Forum on Research Infrastructures (ESFRI). The current status of FAIR construction will be discussed as well as current status and future enlargement of Czech participation in all four pillars of FAIR, i.e. in CBM, NuSTAR, PANDA a APPA. Now days Czech scientist from NPI CAS, Silesian University at Opava and Czech Technical University are involved in studies of compressed baryonic matter (HADES, CBM); team from Silesian University is involved in studies related to nuclear astrophysics (NuSTAR); teams from Charles University and Czech Technical University participate in RD and construction of detectors for PANDA pillar devoted to particle physics. APPA pillar is devoted to application of nuclear and hadronic physics in further fields, like plasma physics, radiobiology, testing of detectors for European Space Agency (ESA) etc. Small group from NPI CAS interested in radiobiology participates now in APPA, further Czech groups involved in other fields covered by APPA as well as by other pillars, are welcomed. Investments of Czech groups in experiments planned to be carried out in FAIR are currently covered mainly by Large Research Infrastructure FAIR-CZ.

Author: KUGLER A. (NPI CAS, Řež)

Presenter: KUGLER A. (NPI CAS, Řež)

Session Classification: Plenary Talks

Contribution ID: 24

Type: **Oral**

TRENDY VO FYZIKÁLNO M VZDELÁVANÍ, OČAKÁVANIA A REALIZÁCIA

Tuesday 8 September 2020 13:30 (35 minutes)

Abstrakt: Vzdelávanie je procesom sprístupňovania nadobudnutých poznatkov a overených postupov z vybranej oblasti ľudskej činnosti, ponúkajúcim využívanie a nadväznosť na existujúcu úroveň vedy, techniky či spoločenských zvyklostí. Kvalitným vzdelávaním nie iba informujeme o minulosti, ale najmä iniciujeme smerom k budúcnosti. Fyzika, ako veda, má obrovský vzdelávací potenciál z pohľadu rozvoja vedeckej gramotnosti, štúdia novodobých vedeckých tém, prepájania základného a aplikovaného výskumu, interdisciplinarita a silnej väzby na informatiku. Fyzika, ako vyučovací predmet, je unikátnym priestorom pre aktívne poznávanie, bádateľskú výučbu, formatívne hodnotenie, kolaboratívne učenie, rozvoj zručností a spôsobilostí, vyšších poznávacích procesov. Náročnosť sprístupňovania vzdelávacieho obsahu vo fyzike si vyžaduje spoluprácu vedcov, didaktikov a učiteľov v oblastiach tvorby kurikula, aplikácie vyučovacích metód, popularizácie a neformálneho vzdelávania. Vysoké očakávania z pohľadu profilu absolventa, kladú požiadavky na vzdelávanie, ktoré položí silné základy pre neustálu adaptáciu, inováciu, kreativitu, či riešenie globálnych výziev, o ktorých dnes iba uvažujeme. Vďaka národným a medzinárodným projektom sa už dnes realizujú vzdelávacie koncepty, ktoré veríme, že naplnia naše očakávania. Koncept gymnaziálnych tried so zameraním na informatiku, prírodné vedy a matematiku, ako jeden z produktov národného projektu IT Akadémia, je v príspevku predstavený s dôrazom na inováciu obsahu a metód fyzikálneho vzdelávania, profiláciu absolventa, jeho orientáciu a pripravenosť na vysokoškolské štúdium STEM odborov.

Author: KIREŠ M. (Ústav fyzikálnych vied Prírodovedecká fakulta UPJŠ v Košiciach)

Presenter: KIREŠ M. (Ústav fyzikálnych vied Prírodovedecká fakulta UPJŠ v Košiciach)

Session Classification: Parallel sessions

Contribution ID: 25

Type: **Oral**

ŠKOLSKÝ MERACÍ SYSTÉM S VYUŽITÍM MIKROKONTROLÉRA

Tuesday 8 September 2020 14:05 (20 minutes)

Zvyšovanie výpočtového výkonu počítačových komponentov, ktoré je sprevádzané znižovaním ich cien je predpokladom vzniku nových progresívnych oblastí, ktoré postupne získavajú na popularite, vďaka čomu získavajú masový rozmer. Tieto nové technológie rýchlo prenikajú do rôznych oblastí nášho života a školstvo nie je výnimkou. V čase, keď väčšina žiakov vlastní digitálne zariadenie sa využitie týchto zariadení aj na vzdelávacie účely javí ako dobré riešenie. Predpokladom takéhoto riešenia je existencia systému, ktoré umožňuje okrem plnenia edukačných úloh aj zabezpečovať komunikáciu so zariadeniami učiteľov a študentov. Školský elektronický merací systém predstavuje technický prostriedok pre realizáciu merania, ktorý následne poskytuje namerané údaje užívateľovi pre ich ďalšie spracovanie. Elektronické meracie systémy predstavujú zaujímavý nástroj pre zefektívnenie výučby fyziky na základných a stredných školách. V príspevku autori predstavujú vlastný vyvinutý školský merací systém základom ktorého je mikrokontrolér ESP32. Veľkou výhodou systému je jeho univerzálnosť a škálovateľnosť, vďaka čomu môže systém plniť aj špecifické požiadavky užívateľa. Modulárnosť systému zase zjednodušuje prácu s jednotlivými komponentmi a sprehladňuje jeho štruktúru aj pri jeho pomerne vysokej komplexnosti. Vďaka bezdrôtovej technológii a použitiu kompatibilných komunikačných protokolov je možné sa k systému pripájať pomocou zariadení bez ohľadu na ich operačný systém. Súčasťou návrhu systému je aj návrh plošného spoja elektroniky a zapuzdrenia komponentov.

Author: S. ILKOVIČ, M. CHOMA**Presenter:** CHOMA M. (Faculty of Humanities and Natural Sciences, Prešov University, Prešov)**Session Classification:** Parallel sessions

Contribution ID: 26

Type: **Oral**

MODELOVANIE ZMENY TIAŽOVÉHO ZRÝCHLENIA A JEHO VPLYVU NA TLAK V KVAPALINE

Tuesday 8 September 2020 14:25 (20 minutes)

Cieľom príspevku je ponúknuť učiteľom fyziky spôsob ako rozvinúť žiacke predstavy o tlaku v kvapalinách, o jeho príčine a dôsledkoch. Zaoberáme sa preto chápaním pojmov tiaž, tiažová sila, gravitačná sila. Venujeme sa tiež pokusu s kvapalinou v beztiažovom stave, kedy na telesá ponorené v kvapaline nepôsobí vztlaková sila. V príspevku navrhujeme experiment zameraný na overenie lineárnej závislosti tlaku v kvapaline od tiažového zrýchlenia. Zmenu tiažového zrýchlenia modelujeme využitím princípu Atwoodovho pádostroja. Experiment odporúčame realizovať so žiakmi vyšších ročníkov stredných škôl a gymnázií.

Author: KOVÁČ M. (Fakulta didaktiky matematiky, fyziky a informatiky, Univerzita Komenského v Bratislave)

Presenter: KOVÁČ M. (Fakulta didaktiky matematiky, fyziky a informatiky, Univerzita Komenského v Bratislave)

Session Classification: Parallel sessions

Contribution ID: 27

Type: **Oral**

Faktory ovlivňující vnímání demonstračních experimentů studenty: Videostudie

Tuesday 8 September 2020 14:45 (20 minutes)

Experiment je základním prvkem nejen ve fyzice jako vědě, ale také ve výuce fyziky. Ačkoliv je v poslední době více výzkumné pozornosti věnováno samostatnému experimentování žáků v rámci badatelské výuky, významnou roli stále ve školách hraje demonstrační experiment. Na katedře didaktiky fyziky MFF UK má projekt předvádění demonstračních experimentů pro studenty středních škol tradici více než 30 let a v současné době nabízí celkem sedm tematických celků.

Výzkum provedený ve školním roce 2017/18 na vzorku cca 5100 studentů však ukázal, že tyto tematické celky jsou studenty vnímány poměrně odlišně zejména z hlediska jejich zájmu, vnitřní motivace a subjektivně vnímané užitečnosti a hodnoty. Nabízí se tedy otázka, zda existují parametry, které předurčují pozitivní vnímání demonstračních pokusů studenty – ať už by mělo jít o požadavky na performeru, o způsob práce s publikem či o výběr jednotlivých experimentů.

Cílem tohoto příspěvku je představit výsledky videostudie, které byly podrobeny videozáznamy všech sedmi tematicky odlišných vystoupení, a interpretovat tyto výsledky ve spojení s tím, jak pozitivně či negativně studenti daná vystoupení vnímali.

Author: ALEXANDR NIKITIN, MARIE SNĚTINOVÁ, PETR KÁCOVSKÝ

Presenter: KÁCOVSKÝ P. (Katedra didaktiky fyziky MFF UK, Praha, Česká republika)

Session Classification: Parallel sessions

Contribution ID: 28

Type: **Oral**

THE PROTON AND DEUTERON ACTIVATION AT NPI CAS AND SPIRAL2/NFS FRANCE

Tuesday 8 September 2020 13:50 (20 minutes)

Knowledge of experimental activation cross sections is important for creation and evaluation of evaluated libraries (TENDL) used by various programs for design accelerators, their components, reactors, but also for nuclear structure study. The proton and deuteron activation cross sections have been studied at NPI CAS (Nuclear Physics Institute) for many years at variable energy cyclotron U120M. Deuteron energy is limited to 20 MeV. For this reason we took advantage of the opportunity to extend this research to the new facility SPIRAL2/NFS (Neutrons For Science) constructed at GANIL/SPIRAL2 in France. Charge-particle irradiation chamber was developed and constructed at NPI CAS. The chamber is equipped with a pneumatic transfer system that allows measurement of activated isotopes with half-lives in minute region. The first test with protons was made at the end of 2019.

The proton induced reaction cross sections on iron were investigated by stacked-foil activation technique with initial proton beam of energy 20 MeV from the cyclotron U-120M of the NPI CAS, Řež and the measurements confirmed the data of previous authors. We determined production cross sections for ^{58m}Co and ^{58g}Co for the first time. We also determined the production cross sections for ^{51}Mn which were incorrectly ascribed to ^{51}Cr . The investigation will continue for higher proton energy interval up to 33 MeV at SPIRAL2/ NFS with the equipment allowing measurement of products with half-lives in minute region (e.g. ^{54m}Co , ^{53m}Fe) and using isotopically enriched targets.

Author: E. ŠIMEČKOVÁ, R. BĚHAL, V. GLAGOLEV, J. MRÁZEK, J. NOVÁK, M. ŠTEFÁNIK

Presenter: ŠIMEČKOVÁ E. (NPI CAS, Řež)

Session Classification: Parallel sessions

Contribution ID: 29

Type: **Oral**

Measurements of open charm hadrons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV by the STAR experiment

Wednesday 9 September 2020 10:30 (20 minutes)

At RHIC energies, charm quarks are primarily produced in hard partonic scatterings at early stages of ultra-relativistic heavy-ion collisions. This makes them an ideal probe of the Quark-Gluon Plasma (QGP) produced in these collisions, since they experience the whole evolution of the hot and dense medium. STAR is able to study the interactions of charm quarks with the QGP through direct reconstruction of hadronic decays of D^\pm , D^0 , D_s , and Λ_c^\pm hadrons. This is

possible thanks to an excellent pointing resolution provided by the Heavy Flavor Tracker.

In this talk, we will present the most recent results on open charm hadron production from the STAR experiment. In particular, we will discuss the nuclear modification factors of D^\pm and D^0 mesons which give access to the charm quark energy loss in the QGP. We will also discuss D_s / D^0 and Λ_c^\pm / D^0 yield ratios as functions of transverse momentum and collision centrality which help us understand better the charm quark hadronization process in heavy-ion collisions. In addition, we will present the rapidity-odd directed flow of D^0 mesons, which can be used to probe the initial tilt of the QGP bulk and effects of early-time magnetic field.

Author: VANĚK J. (Nuclear Physics Institute, Czech Academy of Sciences, Rez)

Presenter: VANĚK J. (Nuclear Physics Institute, Czech Academy of Sciences, Rez)

Session Classification: Parallel sessions

Contribution ID: 30

Type: **Oral**

DETECTION FOR (N,G*) REACTIONS BY FAST NEUTRONS

Tuesday 8 September 2020 14:10 (20 minutes)

Precise nuclear data are of great importance in the design of fusion and ADS devices. The studies of prompt and delayed gamma rays from (n,g*) reactions bring important knowledge as it is one of the main sources of nuclear data such as cross-section, nuclear energy levels, etc.

A central part of the capability is the Cyclotron U-120 and neutron converter coupled to the collimator. The collimated neutron beam of fast neutrons (with energies up to 33 MeV) was recently acquired at the Laboratory of Fast Neutrons of the NPI CAS. The detection system for the prompt gammas irradiated in the reaction of studied material with fast neutrons is being constructed. The setup consists of an array of HpGe detectors. The first experiments performed have shown that the setup was able to detect delayed gammas with decay times of a few ms. Future tests will focus on detecting prompt gammas.

Author: DANIIL KOLIADKO, MARTIN ANSORGE, MITJA MAJERLE, JAN NOVÁK

Presenter: KOLIADKO D. (Nuclear Physics Institute of the Czech Academy of Sciences, Řež)

Session Classification: Parallel sessions

Contribution ID: 31

Type: **Oral**

NUCLEAR DATA MEASUREMENTS WITH COLLIMATED FAST NEUTRON BEAMS PRODUCED BY CYCLOTRON-DRIVEN NEUTRON GENERATORS

Tuesday 8 September 2020 14:30 (20 minutes)

At the Nuclear Physics Institute at the laboratory of Fast Neutron Generators, a new neutron collimator was recently constructed. Simultaneously a new neutron converter coupled to the collimator was constructed and successfully tested last year. Collimated beams of fast neutrons bringing new experimental possibilities especially for on-beam measurements with semiconductor detectors. Motivated by the development of future fission and fusion energy projects the nuclear data measurements of cross-sections for nuclear reactions induced by fast neutrons are under preparation (or were already conducted) at the Department of Nuclear Reactions at the NPI CAS. A new vacuum chamber with silicon detector composed telescopes will allow us to perform precise measurements of double-differential cross-sections for interaction (n, cp) induced by fast neutrons with kinetic energies from 5 to 33 MeV. The current status of development and preparations for experiments with the new vacuum chamber will be given.

Moreover already measured experimental total cross-section data for reaction nat O(n,tot) will be shown and a particular experimental approach to transmission experiments will be described.

Author: MARTIN ANSORGE, MITJA MAJERLE, JAN NOVÁK, DANIIL KOLIADKO

Presenter: ANSORGE M. (Nuclear Physics Institute of the Czech Academy of, Řež)

Session Classification: Parallel sessions

Contribution ID: 32

Type: **Oral**

Recent results of Upsilon production measured with the STAR experiment

Wednesday 9 September 2020 11:50 (20 minutes)

Heavy-ion collisions allow to recreate conditions present in the universe shortly after the Big Bang when the quark-gluon plasma can be formed. The properties of such plasma can be studied using Upsilon mesons. Each of Upsilon states is expected to dissociate at a different temperature reached in the plasma, through Debye-like screening of color charges. Thus, their production yields are suppressed to different levels with respect to the yield in p+p collisions. Additional cold nuclear matter effects can be studied in p+A or d+A collisions. Furthermore, the production mechanism of these heavy mesons is not fully understood and this can be studied in p+p collisions as well.

In this talk, we will present an overview of recent measurements on the production of Upsilon states by the STAR experiment. These include a comprehensive study performed in Au+Au, p+p, and p+Au collisions at $\sqrt{s_{NN}} = 200$ GeV as well as p+p collisions at $\sqrt{s} = 500$ GeV.

Author: LESZEK KOSARZEWSKI FOR THE STAR COLLABORATION

Presenter: KOSARZEWSKI L. (Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague)

Session Classification: Parallel sessions

Contribution ID: 33

Type: **Oral**

FYZIKA PRO NEJMENŠÍ A JEJICH UČITELE

Tuesday 8 September 2020 15:40 (20 minutes)

Co zajímá nejmenší "fyziky" z mateřských škol a prvního stupně základních škol, co se jim líbí a co všechno dokáží, a co na to jejich učitelé? Co vše lze prostřednictvím jednoduchých fyzikálních pokusů u dětí rozvíjet? Jak fyzika pomáhá spolupráci učitelů z různých typů škol? Při odpovídání na předchozí otázky vychází autorka ze své mnohaleté zkušenosti se zprostředkováváním fyziky předškolním a mladším školním dětem a jejich učitelům v rámci vlastního autorského programu Pohádková fyzika i v rámci Elixíru do škol.

Author: HOUFKOVÁ J. (Katedra didaktiky fyziky MFF UK, Praha)

Presenter: HOUFKOVÁ J. (Katedra didaktiky fyziky MFF UK, Praha)

Session Classification: Parallel sessions

Contribution ID: 34

Type: **Oral**

SMÍCHOVSKÁ PRŮMYSLOVKA TŘETÍHO VĚKU

Tuesday 8 September 2020 16:00 (20 minutes)

Smíchovská střední průmyslová škola se dlouhodobě snaží o zavádění nových forem výuky, se kterými souvisí také snaha o status otevřené instituce přístupné všem generacím. Centrem všeho dění jsou samozřejmě studenti školy, kteří mají možnost se všeobecně rozvíjet, a to nejen klasicky v hodinách, ale i jako asistenti či vedoucí workshopů pro mateřské a základní školy a od roku 2015 také pro seniory. Smíchovská průmyslovka třetího věku je dlouhodobý projekt podpořen grantem magistrátu hl. města Praha, který formou pravidelných přednášek a workshopů pomáhá k rozvíjení technických schopností seniorů ale i sociálních vazeb mezi mladou generací a seniory.

Author: KRAJČOVÁ V. (Smíchovská střední průmyslová škola, Praha)

Presenter: KRAJČOVÁ V. (Smíchovská střední průmyslová škola, Praha)

Session Classification: Parallel sessions

Contribution ID: 35

Type: **Oral**

CANCELED!!! STRASTI A SLASTI UČITEĽA FYZIKY V ROKU 2020

Tuesday 8 September 2020 16:20 (20 minutes)

Vyučovací proces, nielen predmetu fyzika, má svoje viac či menej zaužívané pravidlá. Zvyčajne sa ich adeпти na učiteľstvo dozvedajú na pri štúdiu na vysokej škole. V tomto príspevku chceme odpovedať na niekoľko otázok a podať informácie, ktoré sa možno pri štúdiu nedozvedia. Napríklad:

Je vyučovanie fyziky iné oproti iným predmetom? Ako sa mení úloha učiteľa fyziky v posledných rokoch? Sú vytvorené podmienky na kvalitné vyučovanie fyziky na základných a stredných školách? Ako môže vyzerat „dobrá“ vyučovacia hodina fyziky? Čo sa zmenilo na metodickéj pomoci organizácii nazvanej Metodicko-pedagogické centrum v posledných desaťročiach? Kde sú IT technológie rozumné využiteľné a kde ich pomoc je skôr prekážkou? Je 3D tlač pre obyčajného učiteľa fyziky prínosom? Boli sme pripravení na online vyučovanie fyziky a zvládli sme ho?

CANCELED

Author: BEŇUŠKA J.

Presenter: BEŇUŠKA J.

Session Classification: Parallel sessions

Contribution ID: 36

Type: **not specified**

Kulatý stůl k otázkám české terminologie ve fyzice (má ještě smysl psát o fyzice česky?)

Tuesday 8 September 2020 16:45 (1 hour)

Presenter: VALENTA J.

Session Classification: Parallel sessions

Contribution ID: 37

Type: **Oral**

STRUCTURAL AND MAGNETIC PHASE DIAGRAMS OF $\text{RMn}_{1-x}\text{Fe}_x\text{O}_3$ SYSTEM WITH PEROVSKITE CRYSTAL STRUCTURE

Tuesday 8 September 2020 15:40 (35 minutes)

The transition metal oxides, with perovskite structure, have been largely studied because these materials exhibit strong coupling between orbital, electronic, spin and lattice degrees of freedom. The atomic substitution at the dodecahedral sites in stoichiometric compounds has been extensively studied and recently, interest has turned on to the effect of the substitution at the octahedral sites, where the experimental studies have been mainly focused on the magnetic behaviour, and few on the multiferroic properties. In our paper we focus on construction of structural and magnetic phase diagrams of $\text{RMn}_{1-x}\text{Fe}_x\text{O}_3$ system with perovskite crystal structure ($R = \text{Nd, Pr, Tb and Dy}$). Our study was performed on single crystals prepared by optical floating zone method. We studied creation of the substitutional solid solutions in whole concentration range, evolution of Jahn-Teller (JT) effect and orbital ordering with substitution at the octahedral sites with non-active JT ion Fe^{3+} . Construction of magnetic phase diagrams was undertaken with particular emphasis to determination of magnetic structure by means of magnetization, heat capacity and neutron diffraction measurements. We paid special attention to tuning of magneto-electric (ME) coupling in multiferroic compounds (RMnO_3 , $R = \text{Tb, Dy}$) with magnetically induced ferroelectricity by low concentration doping with Fe.

Author: MARIAN MIHALIK, MATÚŠ MIHALIK, MÁRIA ZENTKOVÁ

Presenter: MIHALIK M. (Institute of Experimental Physics, Slovak Academy of Sciences, Košice)

Session Classification: Parallel sessions

Contribution ID: 38

Type: Oral

EXPERIMENTAL STUDY OF EMERGENT GROUND STATE BEHAVIOUR IN RE_iNi_5 (RE = Ce, Yb, Gd) SYSTEM

Tuesday 8 September 2020 16:15 (20 minutes)

The study of intermetallic compounds has a long history. Rare earth intermetallic compounds are systematically investigated more than 3 decades already. Polycrystalline samples $(\text{Ce,Gd,Yb})\text{Ni}_5$ were prepared by different ways, where interesting physical problems were studied.

In this work we focused on competition between the effect of spin fluctuations and the possible existence of quantum criticality, a dimension response of physical properties and the response of chemical pressure on the physical properties.

First of all, structural analysis confirms required crystal structure and scanning electron microscopy shows the chemical compositions. After that, magnetic and thermal properties were performed by using several experimental techniques. The results will be displayed.

Collected experimental data show, that substitution of rare earth atoms in a compound has very important role. An interesting physical property can be observed due to acquired information.

Author: A. DZUBINSKA¹, M. REIFFERS², M. GIOVANNINI³, J. R. FERNANDEZ⁴, R. VARGA¹, J. I. ESPESO³ AND J. C. GOMEZ SAL

Presenter: DŽUBINSKÁ (CPM-TIP, UPJS, 040 11 Košice)

Session Classification: Parallel sessions

Contribution ID: 39

Type: **Oral**

THE SEARCH FOR MULTIFERROELECTRICITY: THE CASE OF NICKEL TELLURIDES

Tuesday 8 September 2020 16:35 (20 minutes)

Multiferroic materials exhibiting coupled magnetic and electric orderings are intensely studied due to an interest in both their fundamental properties and potential applications based on the magnetoelectric coupling. In general, magnons are collective spin excitations of various types. In multiferroics, the electric component of the electromagnetic radiation can excite magnons via the dynamic magnetoelectric coupling. These excitations are then called electromagnons; they can be difficult to distinguish from classical magnons. One possible method consists in comparing terahertz or infrared spectra with Raman spectra. In fact, the electromagnons, unlike magnons, must follow the same selection rules as polar phonons—they must be active in both kinds of spectra.

In this presentation, we will illustrate our approach by the study of nickel tellurides [1,2,3]. Indeed, a colossal magnetoelectric coupling has been found in Ni_3TeO_6 . For the first time, we managed to synthesise its isostructural compounds by substitution of Ni with Mn or Co. Low-temperature spectra revealed electromagnons sensitive to magnetic field, which proves the multiferroic character of the compounds.

Author: C. KADLEC,¹ S. SKIADOPOULOU,¹ M. RETUERTO,² F. KADLEC,¹ F. BORODAVKA,¹ M. MÍŠEK,¹ M. GREENBLATT² AND S. KAMBA¹

Presenter: KADLEC CH. (1 Institute of Physics, Czech Academy of Sciences, Na Slovance 2, Prague)

Session Classification: Parallel sessions

Contribution ID: 40

Type: **Oral**

FRUSTRATED SHASTRY-SUTHERLAND METALLIC SYSTEMS TmB₄ AND ErB₄

Tuesday 8 September 2020 16:55 (20 minutes)

Geometrically frustrated lattices play an important role in emergent quantum mechanical phases, which have been rather well investigated in electronic insulators. A system that exhibits both lattice frustration and metallic behaviour are the rare-earth (RE) tetraborides REB₄. They have a tetragonal structure with magnetic RE ions embedded in boron network and forming a lattice topologically equivalent to the frustrated Shastry-Sutherland lattice with the strong Ising single-ion anisotropy.

In our contribution we compare the behaviour of TmB₄ and ErB₄ by investigating the angular dependencies of their thermal and magnetic properties in a wide range of temperatures (T) and magnetic fields (H) as well as transport properties under pressure up to 3 GPa. Compared are also the rotating magneto-caloric effects (R-MCE) of both compounds. Spin-electron model was suggested to explain the complex behaviour of R-MCE, which is based on the idea of two interacting systems: the localized spins of RE ions and the itinerant electrons in conduction band. The received results from Monte Carlo approach successfully reproduce the observed heating and cooling regions in H - T phase diagrams of TmB₄. Thus, our study shows that measurements of R-MCE can be an effective tool for investigating the microscopic properties of magnetization processes.

Author: S. GABÁNI¹, MAT. ORENDÁČ¹, G. PRISTÁŠ¹, J. BAČKAI², E. GAŽO¹, K. SIEMENSMEYER³, N. SHITSEVALOVA⁴, K. FLACHBART

Presenter: GABÁNI S. (Institute of Experimental Physics SAS, Košice)

Session Classification: Parallel sessions

Contribution ID: 41

Type: **Oral**

SKÚMANIE MIKROŠTRUKTÚRY OCELÍ POMOCOU MÖSSBAUEROVEJ SPEKTROMETRIE

Tuesday 8 September 2020 17:15 (20 minutes)

Mikroštruktúra materiálov determinuje aj ich makroskopické fyzikálne vlastnosti. Tie je nevyhnutné poznať hlavne v prípade ocelí, ktoré sa používajú ako základný materiál pre konštrukčné prvky jadrových zariadení. Ukazuje sa, že chróm v rôznych zliatinách má antikoročné vlastnosti kvôli povrchovej segregácii a tvorbe pasivačnej vrstvy. Práve preto sú ocele s prímiesou chrómu vhodným kandidátom pre tlakové nádoby jadrových reaktorov. Pre výskum mikroštruktúry ocelí je mimoriadne vhodná Mössbauerova spektroskopia, ktorá pomocou hyperjemných interakcií medzi jadrami ^{57}Fe a príslušným elektrónovým obalom citlivo popisuje ich lokálne usporiadanie. Poskytuje informáciu o štruktúrnom usporiadaní (typ kryštalických fáz) a aj magnetickom stave. V prezentácii budú diskutované rôzne spôsoby vyhodnocovania Mössbauerových spektier modelových zliatin typu Fe-Cr s obsahom Cr v rozsahu 1-50 at. %. Pre nízke koncentrácie Cr (do 15 at. %) je vhodná metóda binomického rozdelenia, u vyšších zas distribúcie hyperjemných magnetických polí. Zloženie modelových vzoriek bolo overené metódou röntgenovskej fluorescencie, ich povrch bol skúmaný pomocou rastrovacej elektrónovej mikroskopie. Získané výsledky budú použité pri skúmaní mikroštruktúry reálnych typov konštrukčných ocelí. Práca bola podporená projektom VEGA 1/0130/20.

Author: DÁVID KOŠOVSKÝ, TOMÁŠ KMJEČ, MARCEL MIGLIERINI

Presenter: KOŠOVSKÝ D. (Ústav jadrového a fyzikálneho inžinierstva, Slovenská technická univerzita v Bratislave)

Session Classification: Parallel sessions

Contribution ID: 42

Type: **not specified**

kulatý stůl "Jak učíme fyziku - postřehy a nápady"

Tuesday 8 September 2020 18:30 (1 hour)

Session Classification: Programme open to general public

Contribution ID: 43

Type: **Oral**

TAJOMSTVÁ FUJARY A PÍŠŤALIEK

Tuesday 8 September 2020 19:30 (1 hour)

Píšťalky a najmä fujara sú neodmysliteľnou súčasťou slovenskej kultúry a svetového kultúrneho dedičstva. Najviac osobitými slovenskými píšťalami sú fujara (najdlhší nástroj tohto typu na svete) a koncovka (píšťala bez dierok). Fujara si získala celosvetovú pozornosť najmä špecifickou farbou zvuku, koncovka zasa svojou jednoduchou konštrukciou a spôsobom hry na ňu. Oba tieto nástroje však môžu slúžiť aj ako učebné pomôcky pri výučbe viacerých oblastí fyziky, napríklad šírenie zvuku a stojace vlny, hydrodynamika tekutín, alebo základy harmónie z pohľadu frekvencie zvuku.

Na koncovke a fujare možno pekne ilustrovať (počúť) rôzne typy stojacich vln (vyššie harmonické kmitočty) za okrajovej podmienky ako otvoreného, tak aj uzavretého konca. Vybudením vyšších harmonických kmitočtov sa dajú vysvetliť tradičné hudobné ozdoby hráčov na fujaru, ako sú rozfuk a prefuk. Prúdenie plynu za vzniku vírov je kľúčovým faktorom pri vzniku kladnej spätnej väzby a trvalého tónu. Pytagorejská teória harmónie spolu so štruktúrou vyšších harmonických tónov generovaných v píšťale umožňuje vysvetliť spôsob hry na koncovku a tiež to, prečo minimálny počet dierok na píšťale sú tri (ako má aj fujara) a prečo majú píšťaly najčastejšie 6 dierok.

Ukážky rôznych (aj primitívnych) nástrojov a hry na ne môže spestriť výučbu fyziky a tiež zvýrazniť väzbu fyziky na iné odbory.

Presenter: KUNDRACIK F. (Katedra experimentálnej fyziky, Fakulta matematiky, fyziky a informatiky, Univerzita Komenského v Bratislave)

Session Classification: Programme open to general public

Contribution ID: 44

Type: **Oral**

VODÍK –PALIVO PRO UDRŽITELNOU ENERGETIKU

Tuesday 8 September 2020 20:30 (1 hour)

Odklon od využívání fosilních paliv vyžaduje hledání nových nosičů energie, které budou nezbytné pro dosažení ambiciózního cíle uhlíkově neutrální energetiky. Přestože podíl výroby elektrické energie z obnovitelných zdrojů, především solární a větrné, neustále roste, významnější závislost na těchto zdrojích bude spojena s řešením různých problémů, spojených především s vlivy počasí a geografického umístění na jejich efektivitu. Obtížná regulace produkce elektrické energie z obnovitelných zdrojů vyžaduje hledání způsobu reverzibilního ukládání energie a nových chemických nosičů energie. Zatím se jako nejperspektivnější jeví využití vodíku, a je zřejmé, že vodík bude důležitou součástí energetického mixu budoucnosti. Pokles ceny solární a větrné energie vede k rozvoji výroby „čistého“ vodíku elektrolýzou vody a umožňuje tak ukládání přebytků energie. Energii z vodíku lze zpětně přeměnit ve vodíkovém cyklu na energii elektrickou v palivových článcích, které mohou být jak stacionární, tak mobilní. Mobilní prostředky jsou v podstatě elektrické hybridní systémy, kde je baterie doplněna vodíkovou nabíječkou. Vodík se uplatňuje v manipulační technice, bude využit v elektrickém letectví, kamionové a lodní dopravě. Vodík jako nosič energie má výhodu rychlého čerpání a rovněž delšího dojezdu. Vodíková budoucnost je neoddelitelně spojena s využitím baterií a hybridní systémy budou tím víc nezbytné, čím větší spotřebu bude mít dané zařízení.

Presenter: MATOLÍN V. (Matematicko-fyzikální fakulta, Univerzita Karlova, Praha)

Session Classification: Programme open to general public

Contribution ID: 45

Type: **not specified**

Valná hromada Slovenské fyzikální společnosti CANCELED!!

Wednesday 9 September 2020 08:30 (1h 30m)

Canceled

Contribution ID: 46

Type: **not specified**

Slavnostní předání cen Milana Odehnala, zvaná přednáška vítěze

Wednesday 9 September 2020 08:30 (1h 30m)

Contribution ID: 47

Type: **Oral**

STUDENTS' SOLVING OF MULTIPLE-CHOICE TASKS FOCUSED ON GRAPH SLOPE INTERPRETATION OBSERVED BY THE EYE-TRACKING METHOD

Wednesday 9 September 2020 10:30 (20 minutes)

We employed the eye-tracking method (Eyetracker by Tobii, 300 Hz) in order to observe students' strategies of choosing an option when they solved multiple-choice tasks focused on graph slope interpretation. Each student solved 7 multiple-choice tasks focused on graphs in kinematics. Most tasks were adopted from Beichner's TUG-K test (Beichner, 1994). Particularly, they were focused on position-time graphs and determining velocity from the graphs as it is stated in the Objective 1 of the Beichner's original test. Altogether 23 high school students participated in the study (8 women, 15 men). Each stem-text, stem-graph and each option was marked as a separate Area Of Interest for every task. We provide comparison of attention allocation between two students group: those, who solved a test task correctly and those who not. Based on the data analysis, we observed and described different students understanding of the graph slope concept. Moreover, we applied the cluster analysis in order to divide students into groups according to their attention spent on each option. We obtained very similar results as in the previous case.

Beichner, R. J. (1994). Testing student interpretation of kinematics graphs. *American Journal of Physics*, 62, 750–762.

Presenter: KEKULE M. (Katedra didaktiky fyziky, MFF UK, Praha)

Session Classification: Parallel sessions

Contribution ID: 48

Type: **Oral**

DOPLŇKOVÉ KURZY (BRIDGING COURSES) PRO STUDENTY 1. ROČNÍKŮ NA VYSOKÝCH ŠKOLÁCH

Wednesday 9 September 2020 10:50 (20 minutes)

Cílem příspěvku je prezentovat výsledky mezinárodního šetření o existenci a obsahu překlenovacích kurzů pro absolventy středních škol, kteří nastupují do 1. ročníku studia přírodovědných předmětů na vysokých školách. Byli dotazováni jak studenti, tak i vysokoškolští pedagogové, zda jsou takovéto kurzy na vysokých školách nabízeny, zda studenti tyto kurzy navštěvují, popř. zda by byl o takovéto průpravné kurzy zájem a jaký by měl být jejich obsah (kritická místa kurikula středních škol). Je známo, že v ČR je diskutována úroveň znalostí matematiky absolventů středních škol, zda mají potřebné kompetence pro studium technických oborů, popř. fyziky na technických vysokých školách a přírodovědeckých fakultách (aktuálně také zrušení povinné maturity z matematiky). Je zajímavé porovnat tuto situaci s tím, zda podobné problémy řeší zahraniční vysoké školy a popř. jaké kurzy jsou na vysokých školách nabízeny, na jaké učivo je kladen důraz. Uvedeme příklady nabídky doplňkových kurzů na Přírodovědecké fakultě v Olomouci a příklady z Rakouska, Itálie a Litvy.

Presenter: HOLUBOVÁ R. (Přírodovědecká fakulta Univerzity Palackého Olomouc)

Session Classification: Parallel sessions

Contribution ID: 49

Type: **Oral**

STRATÉGIE ZVOLENÉ ŠTUDENTMI PRI TVORBE KOMENTÁROV K TICHÝM VIDEÁM

Wednesday 9 September 2020 11:10 (20 minutes)

Konstruktivistické prístupy, ktoré sa v súčasnosti uplatňujú vo vzdelávaní v podobe bádateľsky orientovanej výučby, zdôrazňujú okrem rozvíjania porozumenia aj rozvoj zručností a spôsobilostí súvisiacich s činnosťami, ktoré vedú ku konštrukcii poznatkov. Počas prípravy študentov učiteľstva fyziky sa však často stretávame so situáciou, kedy študent uprednostňuje vyučovacie stratégie, ktorými si počas svojho vzdelávania prešiel on sám, a má tendenciu ich napodobňovať. Preto považujeme za dôležité poskytovať študentom učiteľstva neustálu spätnú väzbu a vzor v konštruktivistickom sprístupňovaní poznatkov. Jednou z viacerých príležitostí, kedy môžeme takúto spätnú väzbu našim študentom poskytnúť, je aj realizácia aktivít s tichým videom. Tiché video je typ úlohy, v ktorej študenti nahrávajú komentár k videu trvajúcemu spravidla jednu až dve minúty, zobrazujúcemu nejaký fyzikálny jav, pokus, experiment. Príspevok obsahuje javovú analýzu, na základe ktorej sme určili stratégie zvolené študentmi učiteľstva fyziky pri tvorbe komentárov k trom tichým videám.

Presenter: GORČÁKOVÁ S. (Katedra didaktiky matematiky, fyziky a informatiky, FMFI UK, Bratislava)

Session Classification: Parallel sessions

Contribution ID: 50

Type: **not specified**

Využití konceptuálních úloh a grafického znázorňování v úvodním kurzu kvantové fyziky

Wednesday 9 September 2020 11:30 (20 minutes)

V poslední době se i ve vysokoškolské výuce, zejména v úvodních kurzech během prvních let studia, objevuje posun od tradičně pojaté výuky k aktivizujícím výukovým formám. Jejich prostřednictvím se prohlubuje konceptuální pochopení základních myšlenek, které se tak vzájemně doplňuje s řemeslným zvládnutím konkrétních technik řešení úloh typických pro dané fyzikální téma. Výzkumy prováděné v zahraničí ukazují, že tento přístup má pozitivní efekt jak na pochopení probírané látky, tak na schopnost řešit tyto úlohy.

Konceptuální a aktivizující přístup využíváme ve výuce úvodního kurzu kvantové fyziky pro budoucí učitele již několik let. Velkou výzvou zůstává shánění vhodných materiálů – konceptuálních úloh, otázek do diskuze, námětů na skupinovou práci či vhodných nástrojů pro vizualizaci jednotlivých problémů. V našem případě je navíc velmi vhodné používat materiály v češtině, aby si budoucí učitelé dostatečně zažili terminologii i způsoby diskutování, protože ve své praxi se budou muset obejít bez matematického aparátu, který tvoří přirozenou a jedinou dostatečně výstižnou „řeč“ kvantové fyziky.

V příspěvku představíme ucelenou kolekci konceptuálních úloh i vhodných nástrojů a vizualizací vytvořených jak na základě zahraničních zdrojů, tak vyvinutých přímo ve zmíněném kurzu. Tyto materiály jsou volně dostupné všem zájemcům a mohou tak obohatit výuku úvodních kurzů kvantové fyziky i na dalších školách a oborech.

Author: ZDEŇKA KOUPILOVÁ, PETR KÁCOVSKÝ

Presenter: KOUPILOVÁ Z. (Katedra didaktiky fyziky MFF UK, Praha)

Session Classification: Parallel sessions

Contribution ID: 51

Type: **Oral**

Smart biometric recording system for real-time evaluation of motility on organism with knee pathology

Wednesday 9 September 2020 11:50 (20 minutes)

Lab rats and rabbits are irreplaceable models for study of the orthopedic pathologies. Traditional analysis of these pathologies is based on postmortem histological analysis or X-ray or invasive sampling, however modern monitoring of orthopedic pathology and regeneration needs the smart and noninvasive methods for quantification of the motility of the limbs and organism. Advances in imaging technologies have enhanced our possibilities to quantify the typical movement of the organism.

Lab animals were held in typical lab cage. We developed an autonomous optical setup which recorded animal position and displayed the trajectory and typical characteristic of the movement (speed, acceleration, shading of the walls of cage, recognition of biorhythmic period).

The system was based on the HD 10 fps camera. Export of the frames was committed to OpenCV and mathematical conversion of motion to x-y coordinates was curbed by set of Python utilities.

Comparison of animals with defects and healthy animals gives basic criterion for automatic recognition of pathological state and basic criterion for evaluation of regeneration process. Upgrade of these biometric system should be used also for clinical and home rehabilitation in the future.

Supported by the Ministry of Education, Youth and Sports project "FIT" (Pharmacology, Immunotherapy, nanoToxicology) CZ.02.1.01/0.0/0.0/15 003/0000495

Author: AMIR ENIKEEV A , JOSEF SKOPALÍK A,B , TOMÁŠ PARÁK B , KAROL ŠVÍK C , JANA KOLÁŘOVÁ A , VRATISLAV AMIR ENIKEEV A , JOSEF SKOPALÍK A,B , TOMÁŠ PARÁK B , KAROL ŠVÍK C , JANA KOLÁŘOVÁ A , VRATISLAV ČMIEL A , JOSEF MAŠEK

Presenter: SKOPALÍK J. (Dept. of Biomedical Engineering, Faculty of Electrical Engineering and Communication, University of Technology, Brno)

Session Classification: Parallel sessions

Contribution ID: 52

Type: **Oral**

Nuclear Methods for Astrophysical Purposes

Tuesday 8 September 2020 13:30 (20 minutes)

It is widely known that nucleosynthesis takes place into stars. Element lighter than Iron and heavier than Lithium (partially produced in primordial nucleosynthesis) are in fact produced during stellar evolution by means of the interaction between charged and neutral particles. About charged particles its interactions is strongly hampered by the presence of the Coulomb barrier –usually around some MeV –that is much higher than the typical energies of thermal agitation of particles inside a star (between tenth and thousands of keV). This fact generates the strong, exponential fall of the cross-section that can be observed in many experiments that have been conducted through the years. This often makes really hard –when not virtually impossible –the study of this reactions in our laboratories around the world. Regarding the reactions involving neutrons in the entrance channel, experiments are complicated by the many difficulties involved in produce a suitable neutron beam. To overcome these problems many indirect methods have been developed through the last decades, and in particular the Asymptotic Normalization Coefficient (ANC) method and the Trojan Horse Method (THM) have proven to be quite useful to study reactions involving charged particles and neutrons in the entrance channel.

Author: GIUSEPPE FERDINANDO D'AGATA, JAROMIR MRAZEK ¹, VACLAV BURJAN ¹, ANASTASIA CASSISA

Presenter: D'AGATA G.F. (Nuclear Physics Institute of the Czech Academy of Sciences, Řež)

Session Classification: Parallel sessions

Contribution ID: 53

Type: **Oral**

RECENT RESULTS OF INCLUSIVE JET PRODUCTION IN AU+AU COLLISIONS AT \sqrt{s} NN = 200 GEV BY THE STAR EXPERIMENT

Wednesday 9 September 2020 10:50 (20 minutes)

It has been established that the Quark-Gluon Plasma (QGP), an exotic state of deconfined matter, is created in high-energy heavy-ion collisions. Jets are a very important probe of this hot and dense nuclear matter, since they emerge from the fragmentation of hard-scattered partons (quarks and gluons) that are created during the early stages of the collisions. Therefore, measurements of modifications to jet properties in heavy-ion collisions relative to those in proton-proton collisions can provide insight into understanding interactions between hard-scattered partons and the QGP.

This talk aims to present recent results of jet production in Au+Au collisions at \sqrt{s} NN = 200 GeV by the STAR Collaboration at the Relativistic Heavy Ion Collider. We will focus on the measurement of inclusive jet production as a function of transverse momentum (p_T) of jets and collision centrality. Jets are reconstructed using charged tracks from the Time-Projection Chamber and neutral energy from the Barrel Electromagnetic Calorimeter towers, using the anti- k_T algorithm with jet resolution parameter $R = 0.2, 0.3, \text{ and } 0.4$. The large combinatorial background is suppressed by requiring a high- p_T leading hadron in accepted jet candidates. Jet yield suppression is observed for central relative to peripheral Au+Au collisions, which is attributed to medium-induced parton energy loss. The measured distributions are compared to theoretical calculations incorporating jet quenching, which will improve our understanding of medium-induced energy loss of jets at RHIC energies.

Author: ROBERT LICENIK (FOR THE STAR COLLABORATION)

Presenter: LÍČENÍK R. (Nuclear Physics Institute, Czech Academy of Sciences, Řež)

Session Classification: Parallel sessions

Contribution ID: 54

Type: **Oral**

Search for jet quenching effects in high-multiplicity proton-proton collisions at $\sqrt{s} = 13$ TeV

Wednesday 9 September 2020 11:10 (20 minutes)

QCD jets are modified ("quenched") by their interactions with the quark-gluon plasma (QGP) formed in high-energy nuclear collisions. The measurement of jet quenching in small collision systems can tell us about the limits of QGP formation and the nature of equilibration in QCD, but to date no significant jet quenching has been observed in small systems. In this talk, the ALICE Collaboration reports results of a novel approach to jet-quenching measurements in high-multiplicity pp collisions at $\sqrt{s} = 13$ TeV, searching for modification of di-jet azimuthal acoplanarity measured by the semi-inclusive distribution of jets recoiling from a high-p T hadron. Charged-jet reconstruction is carried out using the anti-k T algorithm with $R = 0.4$ and a data-driven statistical method is used to correct the measured jet yield for uncorrelated background, which includes multi-partonic interactions. High-multiplicity (HM) pp events are selected based on charged-particle multiplicity registered in forward scintillator detectors and their acoplanarity distributions are compared to that for Minimum Bias (MB) events. Significant azimuthal broadening is observed in HM collisions, consistent with jet quenching. However, qualitatively similar features are also seen in pp collisions generated by the PYTHIA Monte Carlo event generator which does not include the simulation of jet quenching or any other QGP effects. We will discuss the current status of this analysis and prospects to understand the origin of this striking phenomenon.

Author: FILIP KRÍŽEK FOR THE ALICE COLLABORATION**Presenter:** KRÍŽEK F. (ÚJF AV ČR, Řež)**Session Classification:** Parallel sessions

Contribution ID: 55

Type: **Oral**

IDENTIFYING HEAVY-FLAVOR JETS USING VECTORS OF LOCALLY AGGREGATED DESCRIPTORS

Wednesday 9 September 2020 11:30 (20 minutes)

Jets of collimated particles arising from hard scattered partons have been studied extensively in hadron collisions. Jets serve a multitude of purposes as they are utilized in fundamental studies of the Standard Model (SM) and in searches for new particles. Recently, studies of jet interaction with the quark-gluon plasma (QGP) created in high energy heavy ion collisions are of growing interest, particularly towards understanding partonic energy loss in the QGP medium and its related modifications of the jet shower and fragmentation. Since the QGP is a colored medium, the extent of jet quenching and consequently, the transport properties of the medium are expected to be sensitive to fundamental properties of the jets such as the flavor of the parton that initiates the jet. Identifying the jet flavor enables an extraction of the mass dependence in jet-QGP interactions. We present a novel approach to tagging heavy-flavor jets at collider experiments utilizing the information contained within jet constituents via the JetVLAD model architecture. We show the performance of this model as characterized by common metrics and showcase its ability to extract high purity heavy-flavor jet sample at various realistic jet momenta and production cross-sections.

Author: GEORGY PONIMAKIN A,B,D , RAGHAV KUNNAWALKAM ELAYAVALLI C , JOSEF ŠIVIC D,E , JANA BIELČÍKOVÁ, JÖRN H. PUTSCHKE

Presenter: PONIMAKIN G. (Nuclear Physics Institute of the CAS, Rez)

Session Classification: Parallel sessions

Contribution ID: 56

Type: **Oral**

THE MOMENTUM SPECTRUM VERSUS THE PROBABILITY FLUX DIRECTION IN QUANTUM MECHANICS

Thursday 10 September 2020 11:25 (20 minutes)

Simple examples that the direction of the momentum, defined by its spectral support in a given state, and the direction of the probability flux may be locally uncorrelated in 1+1 dimensional space-time are given for the Schrödinger and the Dirac particle with non-zero mass. For the zero-mass Dirac particle, they are always correlated.

Jednoduché příklady ukazují, že směr hybnosti, definovaný jejím spektrálním nosičem v daném stavu, a směr toku pravděpodobnosti nemusí být lokálně korelovány pro Schrödingerovy a Diracovy částice s nenulovou hmotností v 1+1 rozměrném prostoročase. Pro Diracovy částice s nulovou hmotností jsou vždy korelovány.

Author: DITTRICH J. (Nuclear Physics Institute CAS, Řež.)

Presenter: DITTRICH J. (Nuclear Physics Institute CAS, Řež.)

Session Classification: Parallel sessions

Contribution ID: 57

Type: **Oral**

PROJEKT IT AKADÉMIA VO SVETLE ZMIEN VO FYZIKÁLNO M VZDELÁVANÍ

Wednesday 9 September 2020 13:30 (20 minutes)

V súčasnosti častokrát skloňované problémy s prírodovedným vzdelávaním sú v centre pozornosti národného projektu IT Akadémia, ktorého hlavným cieľom je vytvorenie modelu vzdelávania pre aktuálne a perspektívne potreby vedomostnej spoločnosti a trhu práce so zameraním na informatiku a digitálne technológie. V rámci príspevku predstavíme dva kľúčové ciele projektu: inovácia vzdelávania matematiky, informatiky a prírodných vied na základných a stredných školách a príprava učiteľov, v rámci ktorých sa sústreďme na fyzikálne vzdelávanie. Prvý z cieľov je zameraný na aktualizáciu obsahu, metód a foriem výučby. Rozsiahla zbierka metodických materiálov pre výučbu je vytvorená na princípoch aktívneho žiackeho poznávania, zmysluplného využívania digitálnych technológií a využívania stratégií formatívneho hodnotenia žiakov. Vytvorené metodické materiály sú v širokom meradle implementované a overované vo výučbe. Spätná väzba získaná z overovania a vzájomná spolupráca učiteľov z praxe a pracovníkov Univerzít vychovávajúcich budúcich učiteľov má napomôcť k reálnemu uplatneniu vytvoreného modelu vzdelávania v praxi. Nato, aby boli tieto zmeny úspešné, je však potrebné motivovať a pripraviť predovšetkým učiteľa, ktorý vo vzdelávaní zohráva kľúčovú úlohu. Z uvedeného dôvodu je vytvorený vzdelávací program, v rámci ktorého sa učitelia fyziky vzdelávajú v oblasti metód bádateľsky orientovanej výučby, digitálnych technológií a hodnotenia výsledkov vzdelávacieho procesu. V príspevku predstavíme základné myšlienky a ukážky vytvorených metodických materiálov a koncepciu vzdelávania učiteľov fyziky.

Author: ZUZANA JEŠKOVÁ, MARIÁN KIREŠ

Presenter: JEŠKOVÁ Z. (Prírodovedecká fakulta UPJŠ, Košice)

Session Classification: Parallel sessions

Contribution ID: 58

Type: **Oral**

VÝSKUM ÚROVNE POZNÁVACIEHO ZÁUJMU ŽIAKOV O FYZIKU

Wednesday 9 September 2020 13:50 (20 minutes)

Predkladaný príspevok je venovaný pedagogicko-psychologickej téme záujmu, ako jedného z významných činiteľov zohrávajúcich dôležitú úlohu v procese samostatného žiackeho bádania a učenia sa. Autori príspevku sústreďujú svoju pozornosť na záujem žiakov o predmet školskej fyziky. Dôraz pritom kladú na poznávací záujem. Hlavným cieľom príspevku je zistiť prítomnosť poznávacieho záujmu o fyziku na základných školách a následne určiť aktuálny stupeň jeho rozvoja. Na základe skutočnosti, že problematika poznávacieho záujmu v súčasnosti nie je v slovenskej literatúre metodicky dostatočne spracovaná, opierajú sa autori vo svojom výskume o metodiku I.J. Laninovej, ktorá vymedzuje tri základné úrovne poznávacieho záujmu. Vzhľadom na všeobecne známy fakt, že fyzika patrí medzi menej obľúbené predmety, doplnili autori príspevku Laninovej klasifikáciu ešte o ďalšiu, štvrtú, úroveň záujmu –tzv. nulový poznávací záujem, odpovedajúci absolútnemu nezájmu žiaka o tento predmet. Za účelom naplnenia vytyčeného výskumného cieľa pripravili autori vlastný neštandardizovaný dotazník určený predovšetkým pre žiakov siedmeho ročníka základných škôl. Analýza a vyhodnotenie výsledkov dotazníkového šetrenia sú súčasťou tohto príspevku. Autori v ňom prezentujú svoje zistenia týkajúce sa jednotlivých úrovní poznávacieho záujmu. Okrem toho, zaoberajú sa aj porovnávaním a skúmaním možného vplyvu niektorých faktorov ako pohlavie či lokalita školy na rozvoj poznávacieho záujmu.

Author: ANDREA KLUČAROVÁ, VLADIMÍR ŠEBEŇ

Presenter: KLUČAROVÁ A. (Katedra fyziky, matematiky a techniky, Fakulta humanitných a prírodných vied, Prešovská univerzita v Prešove)

Session Classification: Parallel sessions

Contribution ID: 59

Type: **Oral**

MOŽNOSTI HODNOTENIA ARGUMENTOV ŽIAKOV V RÁMCI VYUČOVANIA FYZIKY NA ZÁKLADNEJ ŠKOLE

Wednesday 9 September 2020 14:10 (20 minutes)

Človek je spoločenská bytosť a integrálnou súčasťou spoločenského života je aj komunikácia. Pri komunikácii musíme často zdôvodňovať svoje tvrdenia, presvedčať iných o ich pravdivosti –argumentovať. Schopnosť argumentovať však nie je schopnosť, ktorá je človeku daná pri narodení. Je však možné ju rozvíjať a zlepšovať napríklad tým, že žiaci počas hodín vyjadrujú svoje názory pri riešení rôznych príkladov, alebo na základe niektorých faktov predložia svoje tvrdenia. Na vyučovaní fyziky žiaci často majú uviesť svoje predpoklady, vysloviť hypotézy, alebo vysvetliť rôzne javy a deje. Podľa cieľov predmetu fyziky v štátnom vzdelávacom programe by mali byť spôsobilosti žiakov argumentovať rozvíjané počas vyučovania fyziky. V príspevku sa pozrieme na to, ako vyučovanie fyziky reaguje na požiadavku rozvíjania tejto kompetencie v našom vzdelávacom systéme na Slovensku. Na zistenie úrovne argumentácie sme najprv stanovili kritériá, na základe ktorých možno hodnotiť argumenty žiakov. Uvedieme ukážku konkrétnych odpovedí žiakov na fyzikálne úlohy, v rámci ktorých žiaci potrebovali uplatniť argumentáciu, a ich hodnotenie.

Author: TÜNDE KISS, KLÁRA VELMOVSKÁ

Presenter: KISS T. (Oddelenie didaktiky fyziky, Fakulta matematiky, fyziky a informatiky, Univerzita Komenského, Bratislava)

Session Classification: Parallel sessions

Contribution ID: 60

Type: **Oral**

PROBLEMATIKA KLASIFIKÁCIE VÝKONOV ŽIAKA V NEŠTANDARDIZOVANOM DIDAKTICKOM TESTE

Wednesday 9 September 2020 14:30 (20 minutes)

Hodnotenie, ako súčasť výchovno-vzdelávacieho procesu, je v poslednej dobe značne diskutovanou témou medzi odborníkmi v didaktike, učiteľmi, žiakmi, rodičmi ale i v celej spoločnosti. Diskutovanou témou je najmä úloha hodnotenia, jeho funkcie a postavenie vo výchovno-vzdelávacom procese. Taktiež sa vedú úvahy, či je hodnotenie v podobe, akej sa realizuje, objektívne a naozaj posudzuje a klasifikuje žiakov podľa vhodných kritérií tak, aby boli rovnaké výkony žiakov hodnotené totožným klasifikačným stupňom. Proces hodnotenia a klasifikácie žiackych výkonov považujú učitelia za jednu z najnáročnejších úloh učiteľského povolania. Tento fakt je potvrdzovaný viacerými výskumami. Príspevok sa zameria na problematiku náročnosti klasifikácie výkonov žiakov v neštandardizovanom didaktickom teste. V realizovanom prieskume sme analyzovali klasifikáciu výkonov žiakov v neštandardizovaných testových úlohách študentami učiteľstva na FMFI UK. Úlohou prieskumu bolo porovnať jednotlivé hodnotenia navrhnutého neštandardizovaného didaktického testu študentami učiteľstva medzi nimi navzájom a s hodnotením podľa navrhnutej schémy hodnotenia, ktorá vychádza z revidovanej Bloomovej taxonómie. Študenti učiteľstva hodnotili výkony žiakov v didaktickom teste bez kľúča hodnotenia a správnych odpovedí. Do prieskumu sa zapojilo aj niekoľko učiteľov fyziky z praxe a získané dáta sme taktiež podrobili analýze. V príspevku uvidíme priebeh prieskumu, získané výsledky a odporúčania pre pedagogickú prax, ktoré z prieskumu vyplývajú.

Author: JAKUB ČEVAJKA, KLÁRA VELMOVSKÁ

Presenter: ČEVAJKA J. (Fakulta Matematiky, Fyziky a Informatiky, Univerzita Komenského v Bratislave)

Session Classification: Parallel sessions

Contribution ID: 61

Type: Oral

CO NÁM FYZIKOVÉ CHTĚJÍ ŘÍCT K FYZIKÁLNÍMU VZDĚLÁVÁNÍ NA STŘEDNÍCH A ZÁKLADNÍCH ŠKOLÁCH

Wednesday 9 September 2020 14:50 (20 minutes)

Cílem výzkumu, o jehož části pojednává tento příspěvek, je zjistit představy zainteresovaných aktérů o fyzikálním kurikulu. Mezi tyto aktéry patří mimo jiné fyzikové–vědci, další přírodovědci, didaktikové fyziky a učitelé fyziky. Na základě sběru a analýzy jejich představ budou formulovány podněty k proměně současného kurikula fyziky pro základní a střední školy.

Cílem příspěvku je zprostředkovat zkušenosti se sběrem a analýzou dat od fyziků–vědců. Podněty vzešlé od této skupiny aktérů mohou přispět k hlubší reflexi fyzikálního vzdělávání, protože fyzikové stojí do značné míry mimo oblast vzdělávání, zato jsou výrazně ukotveni v oboru. Výzkumný problém byl formulován následující otázkou: Co podle českých fyziků by měla zahrnovat výuka fyziky na základních a středních školách? Data byla získána prostřednictvím rozhovorů v rámci metodologie zakotvené teorie. Výsledky této části výzkumu budou využity k výzkumu mezi dalšími skupinami aktérů a měly by vyústit ve formulování doporučení k tvorbě učebnic fyziky.

Jako předběžné závěry rozhovorů, které byly provedeny s 29 z 60 oslovených fyziků, můžeme uvést následující podněty pro výuku: výrazné propojení výuky fyziky s běžným životem; provádění žákovských experimentů; výuka menšího počtu témat, zato v širších souvislostech; společné vzdělávání různých skupin žáků; zajímavost pro žáky; zařazování současné fyziky, včetně dosud nevyřešených problémů; podpora mimoškolních aktivit.

Author: PETR KOLÁŘ, VOJTĚCH ŽÁK

Presenter: KOLÁŘ P. (Matematicko-fyzikální fakulta, Univerzita Karlova, Praha)

Session Classification: Parallel sessions

Contribution ID: 62

Type: Oral

SOFT MAGNETIC COMPOSITES: PREPARATION, PROPERTIES AND APPLICATIONS

Wednesday 9 September 2020 13:30 (35 minutes)

Nepretržitý pokrok v oblasti magneticky mäkkých materiálov prináša nové možnosti ich využitia v modernej elektrotechnike a elektronike. Cieľom hľadania progresívnych materiálov je snaha o zníženie ekonomických nákladov na ich výrobu a prevádzku z nich zhotovených výrobkov v priemyselných aplikáciách. Magneticky mäkké kompozity (MMK) vyrobené práškovou metalurgiou otvárajú nové možnosti pri navrhovaní a výrobe dielov aj pre nové typy elektromotorov. Vďaka vysokému elektrickému odporu majú MMK v aplikáciách s vysokofrekvenčným premagnetovaním nižšie celkové magnetické straty v porovnaní s klasickými materiálmi. Za magnetické vlastnosti feromagnetík je zodpovedná ich štruktúra, fyzikálne vlastnosti, doménová štruktúra a jej reakcie na magnetické pole. Tieto vlastnosti je možné zlepšiť alebo optimalizovať správnu voľbou a vhodným spracovaním magneticky mäkkého materiálu. MMK sa skladajú z jednotlivito obalených feromagnetických častíc prášku, ktoré sú pokryté elektricky izolačnou vrstvou. Súčasný výrobný postup zahŕňa lisovanie prášku do konečného tvaru a tepelné spracovanie, ktorého cieľom je uvoľnenie napätí indukovaných v priebehu lisovania. Základné metodiky skúmania magnetických vlastností sú meranie hysteréznych slučiek v širokom pásme frekvencií a použitie impedančnej spektroskopie. Koncept MMK založený na využití izolačných vlastností povrchovej vrstvy častíc môže priniesť výrazné výsledky v oblasti prípravy feromagnetík rôznorodých tvarov s optimalizovanými vlastnosťami pre konkrétne aplikácie.

Author: JÁN FÜZER, PETER KOLLÁR, SAMUEL DOBÁK, MAGDALÉNA STREČKOVÁ, ZUZANA BIRČÁKOVÁ, RADOVAN BUREŠ, MÁRIA FÁBEROVÁ

Presenter: FÜZER J. (Ústav fyzikálnych vied, Prírodovedecká fakulta, Univerzita P. J. Šafárika, Košice)

Session Classification: Parallel sessions

Contribution ID: 63

Type: **Oral**

NOVEL DIRECTIONS IN STUDIES OF ELECTRON-HOLE LIQUID IN SEMICONDUCTORS: WIDE GAP AND 2D MATERIALS

Wednesday 9 September 2020 14:05 (40 minutes)

The low-temperature electron-hole plasma may undergo phase transition to the liquid state: the electron-hole liquid (EHL). The transition is allowed in dense plasma below critical temperature T_C and it exists only in the thermodynamic limit, i.e. when the lifetime of the electron-hole system is long enough to reach the thermal equilibrium. For that reason, the macroscopic and relatively long-lived EHL is exclusively observed in the indirect semiconductors. This liquid phase is the two component degenerate Fermi liquid. EHL has a form of spatially separated “droplets” within a cloud of EHP or excitons. In close analogy with conventional liquids EHL has constant density at given temperature –it is related to the quasiparticle distance for which their energy is minimized. Decay of EHL droplets via electron-hole recombination is faster than decay of free excitons - fortunately, this decay includes radiative recombination processes giving rise to photoluminescence (PL) which is the most important messenger bringing information on EHL. The investigation of the intriguing new “matter” was conducted mostly in 1970ies in Si and Ge. EHL was later observed also in low-dimensional semiconductor structures. Here we present recent results on wide-gap semiconductors (especially 4H-SiC) and the graphene-like materials like MoS₂.

Author: VALENTA J. (katedra chemické fyziky a optiky, Matematicko-fyzikální fakulta, Univerzita Karlova, Praha)

Presenter: VALENTA J. (katedra chemické fyziky a optiky, Matematicko-fyzikální fakulta, Univerzita Karlova, Praha)

Session Classification: Parallel sessions

Contribution ID: 64

Type: **Oral**

NITRIDE MULTIPLE QUANTUM WELL CHALLENGE

Wednesday 9 September 2020 14:45 (20 minutes)

Nitride semiconductor heterostructures are widely used for light-emitting, high-power and high-frequency applications. But InGaN/GaN multiple quantum well (QW) structures are also potential candidates for scintillation detectors. This application needs completely different design of the heterostructure in comparison with LED one. It opens new problems that have not been solved yet. Main technological challenge for scintillator structure design is the demand for thick active regions with a higher number of QWs compared to that for LED structures due to the high penetration depth like high energy electrons or X-ray radiation. Another challenge is usually extremely low excitation intensity of ionizing radiation. Under such conditions, the excitonic QW luminescence can have even lower intensity than different kinds of defect bands originating either in GaN or in InGaN QWs. Our ability to realize scintillators on an InGaN/GaN base will be presented and influence of number of QWs in the structure on luminescence properties will be described and discussed.

Author: JIŘÍ OSWALD ¹, FRANTIŠEK HÁJEK ^{1,2}, ALICE HOSPODKOVÁ ¹, TOMÁŠ HUBÁČEK ¹, VÍTĚZSLAV JARÝ ¹, KARLA KULDOVÁ ¹, JIŘÍ PANGRÁC ¹

Presenter: OSWALD J. (Institute of Physics CAS, v.v.i., Prague)

Session Classification: Parallel sessions

Contribution ID: 65

Type: **Oral**

NITRIDE SEMICONDUCTORS – PROPERTIES AND APPLICATIONS

Why are the III-nitride semiconductors so unique that they became the second most important semiconductor material after silicon? Why it was so difficult to prepare high quality nitride layers? It was so difficult that scientist, which have partly solved this problem were five years ago awarded by Nobel Prize. What are the most perspective applications for nitrides? What problems have to be solved in nitride technology? Answers to these four questions concerning nitride properties, applications, technology and open problems will be presented in our contribution. We will show how the piezoelectric field in nitride heterostructures can be utilized in transistors with high electron mobility, but on the other how it can complicate the carrier confinement in quantum wells and design of optoelectronic applications. Application in lasers and light emitting diodes, in high electron mobility transistors (HEMTs) and in scintillators will be discussed with their advantages and limitations. We will touch also the most important technological problems and challenges, such as high dislocation density and possible ways to suppress it, the p-type doping of AlGa_N layers, problems with limited indium atom incorporation. Results of scintillator structure development in our group with the world record InGa_N QW number will be presented.

Author: ALICE HOSPODKOVÁ, TOMÁŠ HUBÁČEK, JIŘÍ OSWALD, FRANTIŠEK HÁJEK, KARLA KULDOVÁ, AND EDUARD HULICIUS

Presenter: HOSPODKOVÁ A. (Institute of Physics CAS, Prague)

Session Classification: Parallel sessions

Contribution ID: 66

Type: **Poster**

VIDEOANALÝZA V STEM VZDELÁVANÍ

Wednesday 9 September 2020 15:50 (30 minutes)

STEM vzdelávanie zamerané na Science, Technology, Engineering a Mathematics zahŕňa v sebe prípravu vysokoškolsky vzdelaného absolventa technickej univerzity, ktorý v priebehu prípravy získal zručnosti zo všetkých spomínaných oblastí. Podľa Bloomovej taxonómie poznávacích cieľov takéto vzdelávanie je realizované na vyšších stupňoch, ako je analýza, aplikácia a hodnotiace posúdenie. Videoanalýza, ako jedna z metód výuky či štúdia, umožňuje rozvíjať matematické zručnosti vo fyzikálnom vzdelávaní a následne pripraviť študentov technického zamerania pre budúcu inžiniersku prax.

Predkladaný príspevok informuje o možnostiach využitia videoanalýzy vo výuke fyziky v inžinierskom vzdelávaní. Zároveň poukazuje na to, že využitím tejto metódy vo výuke je možné zlepšiť u študentov chápanie pojmu sily, aplikácie Newtonových pohybových zákonov, čo potvrdzujú aj výsledky z FCI (Force Concept Inventory) testu.

Presenter: HOCKICKO P. (Katedra fyziky, Fakulta elektrotechniky a informačných technológií, Žilinská univerzita v Žiline)

Session Classification: Poster session

Contribution ID: 67

Type: **Poster**

Elektronická Sběrka řešených úloh z fyziky a Sběrka fyzikálních pokusů

Wednesday 9 September 2020 15:50 (30 minutes)

Sběrka řešených úloh z fyziky a Sběrka fyzikálních pokusů vznikají na Katedře didaktiky fyziky MFF UK. Jedná se o podpůrné materiály pro studenty a učitele základních, středních i vysokých škol.

Sběrka řešených úloh z fyziky je vyvíjena již 14 let a obsahuje přes 900 úloh v češtině. Je primárně určena vysokoškolským studentům a žákům středních škol, lze v ní ale nalézt i úlohy na úrovni základní školy. Sběrka slouží k prohlubování a opakování učiva či k přípravě na zápočty a zkoušky, hojně ji využívají rovněž pedagogové. Protože jedním z našich cílů je seznamovat uživatele s technikami řešení fyzikálních úloh, obsahuje Sběrka nejenom jejich podrobně komentovaná řešení, ale i strukturované nápovědy, ilustrativní obrázky či interaktivní prvky.

Od roku 2015 vzniká také Sběrka fyzikálních pokusů, jejímž cílem je poskytovat učitelům inspiraci pro experimenty na úrovni základní a střední školy. Experimenty, kterých je aktuálně zveřejněno 165, jsou zpracovány do jednotné struktury s důrazem na videozáznam vzorového provedení a technické a metodické poznámky. Tam, kde je to možné, obě Sběrky propojujeme.

Dále vzniká paralelně Sběrka řešených úloh z matematiky s úlohami na vysokoškolské úrovni a také anglické verze vybraných částí všech uvedených Sběrek.

Presenter: SNĚTINOVÁ M. (Katedra didaktiky fyziky MFF UK, Praha)

Session Classification: Poster session

Contribution ID: 68

Type: **Poster**

ÚČAST ČESKOSLOVENSKA NA ZALOŽENÍ SPOJENÉHO ÚSTAVU JADERNÝCH VÝZKUMŮ

Wednesday 9 September 2020 15:50 (30 minutes)

V lednu 1956 bylo Československo přizváno k účasti na založení ústavu pro jaderný, který měl být obdobou Laboratoře Evropské organizace pro jaderný výzkum (CERN) v Ženevě pro země tzv. lidové demokracie. Československé stranické a vládní orgány nabídku přijaly. Na konferenci o ústavu, která se konala ve dnech 20.–26. března 1956 v Moskvě, byla československou vládou vyslána sedmičlenná delegace vedená tehdejším ministrem energetiky Františkem Vlasákem. Zde byla dne 26. března 1956 podepsána Dohoda o založení Spojeného ústavu jaderných výzkumů. Československo se stalo jednou z jedenácti zakládajících zemí ústavu.

S využitím archivních dokumentů budou v příspěvku doloženy bližší okolnosti československé účasti na založení Spojeného ústavu jaderných výzkumů (obsah iniciačního dokumentu, složení a návrhy československé delegace na moskevské konferenci a volba Václava Votruby jedním z prvních dvou náměstků ředitele ústavu).

Presenter: TĚŠÍNSKÁ E. (Ústav pro soudobé dějiny AV ČR Praha)

Session Classification: Poster session

Contribution ID: 69

Type: **Poster**

CALDEIRA-LEGGET MODEL FOR PARTICLE-BATH SYSTEMS IN THE PRESENCE OF A MAGNETIC FIELD

Wednesday 9 September 2020 15:50 (30 minutes)

The Brownian motion of a particle immersed in a bath of charged particles is considered when the system is placed in a magnetic field. The widely accepted Caldeira-Legget particle-bath model is modified so that not only the charged Brownian particle (BP) but also the bath particles respond to the external field. For stationary systems, two equations for the BP motion across the field are derived. They are of the type of generalized Langevin equations with two memory functions. The time correlation function of the thermal force is connected with one of these functions through the fluctuation-dissipation theorem but, unlike all previous theories, it is found to depend on the external field. In the absence of the magnetic force, the other memory function disappears. Analytical expressions are obtained for the velocity correlation functions and other relevant quantities such as the mean square displacement and the diffusion coefficient of the BP for different distributions of the eigenfrequencies of the bath oscillators. Assuming the Drude distribution of the frequencies, it is found that at long times the motion of the particle is sub-diffusive, with the exponent $1/2$. The case of the fractional thermal noise is also analyzed.

Presenter: LISÝ V. (Department of Physics, Faculty of Electrical Engineering and Informatics, Technical University of Košice)

Session Classification: Poster session

Contribution ID: 70

Type: **Poster**

BROWNIAN MOTION IN A BATH AFFECTED BY AN EXTERNAL HARMONIC POTENTIAL

Wednesday 9 September 2020 15:50 (30 minutes)

The Brownian motion of a particle in a bath of other particles is effectively described by the generalized Langevin equation (GLE). Following Kubo, it is usually assumed that if external forces act on the system, they do not affect the thermal force and the memory function that enter the GLE. The action of such forces is restricted to the Brownian particle (BP), leaving the bath particles unaffected by the external field. However, there are many physical situations, when not only the BP but also the bath particles are subjected to the external field. We show that for stationary systems in a harmonic potential the corresponding generalization of the Zwanzig-Caldeira-Legget theory leads to the GLE for which Kubo's fluctuation-dissipation theorem remains valid but both the memory function and the thermal force depend on the elastic constant of the confinement potential. As a result, the correlation functions describing the random motion of the BPs change in comparison with those in the original model as well. We discuss possibilities to calculate these functions and show several specific solutions for them depending on the frequency distribution of the bath oscillators and the coupling between the bath and the BP.

Presenter: TÓTHOVÁ J. (Department of Physics, Faculty of Electrical Engineering and Informatics, Technical University of Košice)

Session Classification: Poster session

Contribution ID: 71

Type: **Poster**

BIPARTITE ENTANGLEMENT IN THE SPIN-1/2 ISING-HEISENBERG PLANAR LATTICE CONSISTING OF INTER-CONNECTED TRIGONAL BIPYRAMIDS

Wednesday 9 September 2020 15:50 (30 minutes)

The present work deals with the exactly solvable spin-1/2 Ising-Heisenberg model on an infinite but regular two-dimensional lattice composed of identical inter-connected bipyramidal plaquettes with the aim to clarify a bipartite entanglement between the Heisenberg spins at zero as well as finite temperatures. The quantity called concurrence is used as an indicator for determining a strength of this quantum-mechanical correlation.

It is demonstrated that the Heisenberg spins of each bipyramidal plaquette can be mutually entangled at zero temperature only if the two-fold degenerate spontaneously ordered quantum phase characterized by a symmetric quantum superposition of three possible up-up-down (or down-down-up) states of these spins is stable. Otherwise, the bipartite quantum entanglement of the Heisenberg spins is totally absent. Interestingly, the entanglement between the Heisenberg spin pairs persists also at finite temperatures, even far above the critical temperature of the model if the exchange anisotropy between these spins is sufficiently strong. The entangled Heisenberg spin states can also be thermally activated above non-entangled ground state if the values of the exchange anisotropy parameter are taken sufficiently close to the boundary with the quantum ground-state phase.

Presenter: GÁLISOVÁ L.GALISOVA.LUCIA@GMAIL.COM (Institute of Manufacturing Management, Faculty of Manufacturing Technologies with the seat in Prešov, Technical University of Košice)

Session Classification: Poster session

Contribution ID: 72

Type: **Poster**

PHASE BIREFRINGENCE DISPERSION FUNCTION AS A COMPLEMENTARY TOOL FOR INVESTIGATION OF NEMATIC LIQUID CRYSTAL STRUCTURE

Wednesday 9 September 2020 15:50 (30 minutes)

If we put a several tens of microns thin sample of nematic liquid crystal with planar alignment of molecules with respect to the LC cell surface in between two crossed plane polarizers, illuminate it by white light and rotate appropriately, then using a fiber optic spectrometer we can observe a quasi-periodic spectral distribution of intensity of light behind the output polarizer. The spectral distribution of intensity of light emerges due to liquid crystal birefringence. Based on the polarizability models from which imply models describing the extraordinary and ordinary refractive indices n_e and n_o , respectively, the phase birefringence dispersion function of LC in the spectral region of wavelengths without resonant frequencies can be expressed. Comparison of the measured spectrum with that obtained by calculation according to the model one can find parameters characterizing mean resonance (absorption) wavelengths as well as coefficients expressing the strength of the influence of these absorption bands to resulting birefringence. The approach could complement absorption or ellipsometric measurements the aim of which was to characterize optical properties of a liquid crystal sample.

Presenter: TARJÁNYI N. (Department of Physics, Faculty of Electrical Engineering and Information Technology, University of Žilina)

Session Classification: Poster session

Contribution ID: 73

Type: **Poster**

Role of water molecules in the phase transitions in lawsonite

Wednesday 9 September 2020 15:50 (30 minutes)

Lawsonite [$\text{CaAl}_2\text{Si}_2\text{O}_7(\text{OH})_2 \cdot \text{H}_2\text{O}$] is an uncommon mineral found in metamorphic rocks, occurring at subduction zones, supposedly at depths of up to 250 km. It contains 11.5 wt% water in its crystal structure. At room temperature, it is orthorhombic, crystallizing in the Cmc₂m space group. Its structure contains a silicate tetrahedra framework with four formulas in the unit cell. Each chemical formula comprises one water molecule embedded in a structural cavity; these cavities form channels parallel with the c axis. The protons form O-H bonds, and they may diffuse along the channels. The static and dynamic orientations of the water molecules plays a key role in two low-temperature phase transitions—a structural one at 270 K, and a ferroelectric one at 124 K. We studied the lattice dynamics in a single crystal of lawsonite using infrared, Raman and THz time-domain spectroscopies. We have found a strong soft phonon mode linked to the ferroelectric phase transition. The observed unusual hardening of other phonons with heating is linked to anomalous temperature dependence of a unit cell parameter. In view of a dielectric anomaly identified earlier, our results show that the ferroelectric phase transition is of mixed displacive / order-disorder type.

Presenter: KADLEC F. (Institute of Physics, Academy of Sciences of the Czech Republic, Prague)

Session Classification: Poster session

Contribution ID: 74

Type: **Poster**

COMPLEX VOLUME CHANGES INDUCED BY HIGH PRESSURE AND EXTERNAL MAGNETIC FIELD IN Ni₂MnSn-BASED HEUSLER ALLOY

Wednesday 9 September 2020 15:50 (30 minutes)

The Heusler Ni₂Mn_{1+x}Sn_{1-x} alloys exhibit structural martensitic transformation from cubic (austenite) into orthorhombic (martensite) phase in a narrow composition range, $0,4 \leq x \leq 0,6$, only. The transformation is accompanied by a significant changes of volume, magnetization and by large anomalies in transport properties. We have studied forced volume magnetostriction, Joulian magnetostriction, thermal expansion and effect of high pressure on magnetization of the Ni_{1.92}Mn_{1.56}Sn_{0.52} alloy in wide range of temperature, magnetic field and pressure. The pronounced decrease of magnetization under pressure, $= -11.810 \cdot 10^{-3} \text{ GPa}^{-1}$, points to an itinerant character of magnetism of the alloy. Using the Maxwell relation, $d\omega/dH = -\rho dM/dP$, the received value of $d\ln M/dP$ can be perfectly compare with value of forced volume magnetostriction, $d\omega/dH = 3.1110 \cdot 10^{-6} \text{ T}^{-1}$, received by dilatometric measurement at field above 0.3 T. The Joulian magnetostriction at low temperature and low magnetic field (with $\Delta L_{\parallel} < 0$ and $\Delta L_{\perp} > 0$) confirms a competition between strengths of magnetocrystalline anisotropy and elastic energy in martensite of Ni_{1.92}Mn_{1.56}Sn_{0.52} that is characterized by a zig-zag twins structure. The relevant model proposed by O'Handley will be discussed. The observed change of volume during martensitic transformation of the alloy was verified by X-ray diffraction measurements.

Presenter: KAMARÁD J. (Institute of Physics ASCR, Prague)

Session Classification: Poster session

Contribution ID: 75

Type: **Poster**

NEW VIEW ON THE ORIGIN OF HIGH CONDUCTIVITY OF POLYANILINE FILMS PROTONATED BY HYDROCHLORIC ACID

Wednesday 9 September 2020 15:50 (30 minutes)

Polyaniline (PANI) is a material known for its high conductivity and also huge range of obtainable conductivities. In general, three types of charge carriers are considered in PANI salt, namely holes, polarons and protons. Many authors attribute the high conductivity of acid doped PANI to polarons [1-3]. We show that conductivity of polarons might not be necessarily the prevailing mechanism of charge transport. In our research in order to separate contributions from different charge carriers PANI/Si heterojunction was studied. Impedance spectra were measured to distinguish between polaron and hole mobilities. From V-A characteristics on PANI/Si structure contribution of polaron and hole conductivity was calculated. Dielectric spectra of heterojunction were measured to estimate conductivity of ions. Both measurements resulted in values in order of magnitude $\sim 10^{-7}$ S.cm⁻¹. On the contrary conductivity of several S.cm⁻¹ on films of PANI on glass substrates were obtained using ohmic gold electrodes. To explain discrepancy between obtained values of conductivity a model utilising redox reaction of hydrogen and chlorine was proposed and diffusion of hydrogen and chlorine molecules in PANI films were studied.

The work was supported by the project SVV 260 444/2018 and grant GAUK No. 190119 provided by Charles

- ☒ [1] Y. Harima, R. Patil, K. Yamashita, N. Yamamoto, S. Ito, A. Kitani, Chem. Phys. Lett. 345, 239-244 (2001).
- ☒ [2] M. Chipara, G. Aldica, D. Hui, M. Dimonie, K. T. Lau, L. Georgescu, I. Munteanu, H. Maroscoiu, J. Optoelect. Adv. Mater. 6, 297-305 (2004).
- ☒ [3] J. Kim, S. Park, and N. F. Scherer, J. Phys. Chem. B 112, 15576-15587 (2008).

Presenter: RUTSCHR. (Department of Macromolecular Physics, Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Poster session

Contribution ID: 76

Type: **Poster**

MÖSSBAUER SPECTROMETRY IN THE STUDY OF METALLIC GLASSES

Wednesday 9 September 2020 15:50 (30 minutes)

Metallic glasses are still attracting the interest of researchers namely for their very good soft magnetic properties. With the aim to enhance their practical applications, new compositions are continuously scrutinized. Here, we present Mössbauer spectrometry study of a novel Fe 51 Co 12 Si 16 B 8 Mo 5 P 8 metallic glass prepared in a form of ribbons by conventional method of planar flow casting. Samples in as-quenched state as well as after annealing at selected temperatures were investigated.

Mössbauer spectrometry is one of few analytical tools that can describe disordered amorphous systems. This can be done via hyperfine interactions between nuclei and electron shells which sensitively probe local short-range order arrangement. Experiments performed in a broad range of temperatures provide information on the evolution of microstructure. This is reflected by continuous modification of the hyperfine interactions from magnetic dipole towards electric quadrupole ones. Eventually, the Curie temperature of the investigated metallic glass can be established.

This work was supported by the grants VEGA 1/0130/20, APVV-16-0079 and by the European Regional Development Fund-Project "Center for Advanced Applied Sciences" No. CZ.02.1.01/0.0/0.0/16_019/0000778.

Presenter: MIGLIERINI M. (Institute of Nuclear and Physical Engineering, Slovak University of Technology in Bratislava)

Session Classification: Poster session

Contribution ID: 77

Type: **Poster**

SOLID-STATE ¹H NMR STUDY OF CORNSTARCH PLASTICIZED WITH UREA AND GLYCEROL DURING AGEING

Wednesday 9 September 2020 15:50 (30 minutes)

Thermoplastic starch, a group of starch-based materials, has often been under review because it can be used as low-cost bioplastics. Native starch is usually processed under the action of high temperature and shear stress with addition of plasticizers such as low molecular weight polyols and/or amide-containing molecules to obtain a moldable thermoplastic material. Such materials are sensitive to ageing and time dependent changes in their structure are of great importance for the use of materials in practice. In this study, solid-state magic-angle spinning ¹H NMR was used to characterize structural changes in cornstarch plasticized with glycerol and urea during one-year ageing. It was found that widths of the signals related to urea and glycerol CH/CH₂ groups did not change significantly while widths of the signals coming from water and glycerol OH groups decreased by about half after one year of storage. This could mean that urea formed stable hydrogen bonds with starch which were not affected too much during storage whereas hydrogen bonds between glycerol OH groups and urea were disrupted which resulted in increased mobility of glycerol and perhaps also phase separation of glycerol and water during ageing.

Presenter: ŠMÍDOVÁ N. (Department of Physics, Faculty of Electrical Engineering and Informatics, Technical University of Košice)

Session Classification: Poster session

Contribution ID: 78

Type: **Poster**

IMAGING HIGH-ASPECT RATIO NANOSTRUCTURES ON SOFT AND BRITTLE MATERIALS USING ATOMIC FORCE MICROSCOPY

Wednesday 9 September 2020 16:20 (30 minutes)

Atomic Force Microscopy (AFM) is a powerful imaging technique which provides very high-resolution images and topological data in scales of nanometers. The quality of the obtained data is determined by quality of used AFM tip and fine adjustment of measuring parameters and settings of the machine. The measurement becomes more challenging if the AFM probe is about to approach a soft and elastic surface such as silica gel polymers (polydimethylsiloxane) or biological membrane surfaces (insect wings) which due to their properties tend to bend and vibrate under the probe during scanning. The polymers tend to stick to the bottom of the tip and confuse the software controlling piezo motors during approach causing either false approach or putting a force on the sample, so it bends to concave meniscus or even puncture the sample surface. These malfunctions become even more problematic if a dense high-aspect ratio nanopillar network occurs on such surfaces. Yet, the ability to measure raw materials without any need of surface modification (as often needed for non-conductive materials for electron microscopy) is a decisive advantage for attempting to use AFM even on such materials. Here we present AFM topography characterization of dragonfly wings nanostructure and their polymer replicas manufactured from various soft polymers.

Presenter: ČERNÝ M. (Department of chemical physics and optics, Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Poster session

Contribution ID: 79

Type: **Poster**

THE STUDY OF LATTICE DYNAMICS IN LOW-DIMENSIONAL QUANTUM MAGNETS WITH CHAIN-LIKE CRYSTAL STRUCTURE

Wednesday 9 September 2020 16:20 (30 minutes)

Current work is focused on the lattice subsystem of Cu(en)Cl_2 , Cu(tn)Cl_2 and $\text{Cu(en)(H}_2\text{O)}_2\text{SO}_4$ ($\text{en} = \text{C}_2\text{H}_8\text{N}_2$, $\text{tn} = \text{C}_3\text{H}_{10}\text{N}_2$), quasi-two-dimensional quantum magnets with

one-dimensional polymeric structure. Magnetic layers were described within spin = 1/2 Heisenberg models on the rectangular and zig-zag-square lattice with nearest-neighbor couplings $J_1 > J_2$. The largest spatial anisotropy $R = -J_2/J_1$ and the strongest effect of interlayer coupling J' was observed in Cu(en)Cl_2 , while somewhat lower R and much weaker J' in Cu(tn)Cl_2 and $\text{Cu(en)(H}_2\text{O)}_2\text{SO}_4$.

Specific heat in zero magnetic field was measured from 2 to 300 K and Raman spectra at 300 K. In Cu(en)Cl_2 and Cu(tn)Cl_2 , specific heat anomalies observed at 138 K and 160 K, respectively, are associated with a structural phase transition. No structural phase transition was observed in $\text{Cu(en)(H}_2\text{O)}_2\text{SO}_4$. The contribution of acoustic modes is described within Debye approximation with Debye temperatures $\Theta_D = 147$ K, 109 K and 93 K for $\text{Cu(en)(H}_2\text{O)}_2\text{SO}_4$, Cu(en)Cl_2 and Cu(tn)Cl_2 , respectively. The larger Θ_D value of Cu(en)Cl_2

and the Raman shift towards higher energies correspond well with the Cu(en)Cl_2 specific heat values which are lower than those of Cu(tn)Cl_2 in the whole temperature region. The influence of acoustic modes on the magnetic correlations is discussed.

Presenter: VINNIK O. (Institute of Physics, P. J. Šafárik University, Košice)

Session Classification: Poster session

Contribution ID: **80**Type: **Poster**

PROPERTIES OF YELLOW BAND IN GAN LAYERS

Wednesday 9 September 2020 16:20 (30 minutes)

Nitride semiconductors became the second most important semiconductor material after silicon in the last two decades. Many daily used devices, such as blue light emitting diodes or high electron mobility transistors in electronic devices, are based on nitride semiconductors. Despite the nitride devices mass production, there are many open questions and not well understood phenomena which have to be solved.

Epitaxially grown GaN layers contain different bands in their luminescence spectra. There is very fast excitonic band and also different kinds of defect bands, which have mostly slow decay time. Fast decay time without any slow components is necessary for many scintillation applications. Especially defect band with the maximum emission around 2.2 eV (called yellow band) has very slow decay time (microsecond range) and needs to be suppressed in fast scintillators.

In our previous work [1], we have shown that different technological parameters during Metal Organic Vapour Phase Epitaxy (temperature, carrier gas, etc.) influence properties of yellow band quite significantly. In this work, we continue to study properties of yellow band of GaN layers grown with different parameters. Photoluminescence, time-resolved photoluminescence and other measuring techniques are used for investigation of yellow band properties. Origin of yellow band will be discussed.

[1] T. Hubáček et al., J. Cryst. Growth 531 (2020) 125383.

Presenter: HUBÁČEK T. (Institute of Physics CAS, v.v.i., Prague)

Session Classification: Poster session

Contribution ID: 81

Type: **Poster**

EVOLUTION OF MAGNETISM IN $\text{UCo}_{1-x}\text{Ir}_x\text{Ge}$ SYSTEM

Wednesday 9 September 2020 16:20 (30 minutes)

The uranium based intermetallics are intensively studied because of dual character of uranium 5f electrons. Unique coexistence of ferromagnetism and superconductivity has been revealed in URhGe and UCoGe crystalizing in the orthorhombic TiNiSi-type structure. Since UCoGe ($T_C = 2.5$ K) is very close to magnetic instability, it is a candidate to observe vanishing of ferromagnetic order and ferromagnetic quantum critical point. The chemical substitution was found as an effective tool to destabilize the ferromagnetic order in UCoGe by substitution on Co site by transition metal. $\text{UCo}_{1-x}\text{Ru}_x\text{Ge}$ or $\text{UCo}_{1-x}\text{Fe}_x\text{Ge}$ systems have showed vanishing of

ferromagnetism in quantum critical point at critical concentration ≈ 30 % of substituent element. However, in $\text{UCo}_{1-x}\text{Rh}_x\text{Ge}$ or $\text{UCo}_{1-x}\text{Pd}_x\text{Ge}$ where both parent compounds order magnetically, finite ordering temperature is conserved, thus quantum critical phenomena have not been observed.

We have decided to investigate $\text{UCo}_{1-x}\text{Ir}_x\text{Ge}$ system. Since UCoGe is ferromagnet and UIrGe antiferromagnet, its alloy system is promising candidate to observe interesting evolution of magnetism. We will show that $\text{UCo}_{1-x}\text{Ir}_x\text{Ge}$ is the first case of disappearance ordering temperature between two magnetic parent compounds. Therefore, two quantum critical point of different nature of ferromagnetic and antiferromagnetic-type are expected.

In order to investigate evolution of the magnetism we have prepared polycrystalline samples throughout whole concentration range. Subsequently, we have performed measurements of magnetization, electrical resistivity and heat capacity.

Presenter: HOVANČÍK D. (Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Poster session

Contribution ID: 82

Type: **Poster**

STRUCTURE AND GROWTH OF OMEGA-Ti NANOPARTICLES IN BETA-Ti SINGLE CRYSTALS STUDIED BY ANOMALOUS X-RAY DIFFRACTION

Wednesday 9 September 2020 16:20 (30 minutes)

Nanoparticles of hexagonal ω phase in bcc-Ti(Mo) single crystals (β phase) occur due to a diffusionless athermal β to ω transformation and they grow during follow-up ageing at elevated temperatures, while the alloying atoms (Mo in our case) are expelled from the nanoparticle volumes. We used anomalous x-ray diffraction for the study of local chemical composition of growing ω particles and we found that a Mo-rich shell at the particle/matrix interface is created during the particle growth. Moreover, the expelled Mo atoms create a cloud with higher Mo density around each particle and its Mo concentration profile was studied by anomalous x-ray diffraction and small-angle scattering.

Presenter: HOLÝ V. (Department of Condensed Matter Physics, Charles University, Prague)

Session Classification: Poster session

Contribution ID: 83

Type: **Poster**

ANALÝZA TOMOGRAFICKÝCH REKONSTRUKCÍ Z DAT TOKAMAKU COMPASS

Wednesday 9 September 2020 16:20 (30 minutes)

Tokamak COMPASS je vybaven dvěma sadami diagnostik, které jsou rutině využívány k počítání tomografických inverzí. Jedná se o polovodičové diody, které slouží buď jako bolometry (měří celkový vyzářený výkon), nebo jsou vybaveny beryliovým filtrem a měří měkké rentgenové záření.

V důsledku omezení pozorovacích úhlů je tomografická inverze na tokamacích nedostatečně určená, a proto se k inverzi dat často využívá algoritmus Minimalizace Fisherovy informace využívající Tichonovovu regularizaci. Tento příspěvek se zabývá analýzou takto spočítaných inverzí pomocí metod umožňujících zkoumat časový a prostorový vývoj emisivity plazmatu. Princip spočívá ve sledování oblastí zájmu, které mohou mít jak obdélníkový tvar, tak tvary odvozené od tvaru magnetických povrchů. Tyto metody lze využít ke zpracování rekonstrukcí z více diagnostik najednou.

V příspěvku budou prezentovány analýzy rekonstrukcí vybraných výbojů tokamaku COMPASS. Analýza rekonstrukcí měkkého rentgenového záření se zaměří na pilovou nestabilitu a vnitřní smyčkovou nestabilitu. Analýza dat z bolometrů bude zaměřena na jev tzv. oddělení plazmatu od divertoru a na studium okrajové nestability.

Presenter: SVOBODA J. (Ústav fyziky plazmatu AV ČR, Praha)

Session Classification: Poster session

Contribution ID: 84

Type: **Oral**

ENHANCING GOLD BY GOLD: PLASMON ENHANCED LUMINESCENCE OF GOLD NANOCCLUSERS ON GOLD NANORODS

Thursday 10 September 2020 08:30 (20 minutes)

One of the fundamental features of nanotechnology is that properties of a certain material can be changed just by varying its shape and size. This exactly is the case for the two types of gold nanostructures in our study, where luminescent gold nanoclusters (AuNCs) benefit from the presence of plasmonic gold nanorods (AuNRs). Despite the limited number of atoms, AuNRs with characteristic dimensions ranging from ~ 10 to 100 nm still retain their metallic properties. Consequently, collective oscillations of the nearly free conductive electrons (so called localized surface plasmon resonance; LSPR) can be excited in AuNRs upon interaction with an incoming electromagnetic radiation. In recent years, LSPR has been widely used to enhance the optical performance of other nearby objects (e.g. surface-enhanced Raman scattering or luminescence).

The electronic structure of AuNCs with diameter of ~ 1 nm is, on the other hand, rather molecular-like, which can result in near-infrared photoluminescence from the nanoclusters. In the present study, we attempt to increase the radiative rate in AuNCs by coupling their emission to plasmons of AuNRs in a core@shell AuNRs@AuNCs structure. By measuring the changes in luminescence of AuNCs upon attachment to AuNRs we demonstrate that it is indeed possible to enhance gold by gold.

Author: ONDŘEJ PAVELKA A, KLAUDIA KVAKOVÁ B, PETR CÍGLER B, JAN VALENTA A

Presenter: PAVELKA O. (Department of Chemical Physics and Optics, Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Parallel sessions

Contribution ID: 85

Type: **Oral**

DIELEKTRICKÁ SPEKTROSKOPIA FUNKCIONALIZOVANÝCH KARBÓNOVÝCH NANORÚROK

Thursday 10 September 2020 08:50 (20 minutes)

Štúdium dielektrických parametrov vedie k lepšiemu porozumeniu polarizačných a relaxačných javov v nových materiáloch. Táto štúdia je zameraná na mnohostenné karbónové nanorúrky funkcionalizované magnetickými nanočasticami. Pomocou dielektrickej spektroskopie sa študoval vplyv elektrického poľa na zmeny dielektrických parametrov i štruktúrneho usporiadania MWCNT/Fe₃O₄. Frekvenčná závislosť disipačného faktora sa merala v rámci frekvenčného rozsahu od 10 mHz do 2 MHz v rozmedzí teplôt od 15 °C do 65 °C pomocou kapacitnej metódy. Analýza nameraných údajov sa uskutočnila pomocou relaxačného Cole-Cole modelu. Pozorované nízkofrekvenčné relaxačné maximum disipačného faktora je vysvetlené Schwarzovou teóriou polarizácie elektrickej dvojvrstvy pričom s teplotou sa posúvalo do oblasti vyšších frekvencií. Elektrické pole vplýva na zmenu elektrického dipólového momentu nanorúrok, nanočastíc a orientáciu v smere elektrického poľa spojenú aj s tvorbou reťazcov.

Author: KÚDELČÍK J. (Katedra fyziky, Fakulta elektrotechniky a informačných technológií, Žilinská univerzita V Žiline)

Presenter: KÚDELČÍK J. (Katedra fyziky, Fakulta elektrotechniky a informačných technológií, Žilinská univerzita V Žiline)

Session Classification: Parallel sessions

Contribution ID: 86

Type: **Oral**

QUANTUM DOTS GROWN BY METALORGANIC VAPOR PHASE EPITAXY

Thursday 10 September 2020 09:10 (20 minutes)

This paper focuses on quantum dots (QDs) embedded inside semiconductor heterostructures prepared by Metalorganic Vapor Phase Epitaxy (MOVPE) technology and is based on our contribution in [1].

Light emission from atoms/molecules in a glass matrix or in gas is strongly monochromatic. Semiconductor direct-bandgap materials have much higher energy conversion efficiency than the other light sources, but they have a broad band or multimode light emission spectra. QDs inside semiconductor heterostructures can fundamentally improve the quality of spectrum, temperature dependencies and also light efficiency emission.

The main technological procedure used for MOVPE preparation of QDs embedded inside the heterostructure is self-assembled Stranski–Krastanov growth mode. Growth procedures and parameters will be briefly described.

Mostly used material combination for embedded QDs is InAs QDs in a GaAs matrix.

Embedded MOVPE-prepared QDs are currently used for semiconductor lasers, optical amplifiers, LEDs and photodetectors. High extinction coefficient of QDs is promising for possible optical applications. QDs can operate like a single-electron transistor and show the Coulomb blockade effect. QDs have also been suggested for quantum information processing . QD technology is relevant to solid-state quantum computation .

[1] Metalorganic Vapor Phase Epitaxy (MOVPE): Growth, Materials Properties and Applications”, edited by S.J.C. Irvine and P. Capper, published by Wiley (John Wiley Sons) 2019, autumn.

Author: EDUARD HULICIUS, ALICE HOSPODKOVÁ, AND MARKÉTA ZÍKOVÁ

Presenter: HULICIUS E. (Institute of Physics, Czech Academy of Sciences, Prague)

Session Classification: Parallel sessions

Contribution ID: 87

Type: **Oral**

Determining the alignment of thin MoS₂ layers from Raman and GIWAXS measurements

Thursday 10 September 2020 09:30 (20 minutes)

Thin MoS₂ layers grown by sulfurisation of Mo films can be aligned horizontally or vertically depending on the initial Mo thickness. So far, the vertical alignment of MoS₂ layers has been detected by TEM measurements. However, TEM provides information only from a very limited area of the sample and requires time consuming and laborious sample preparation. Alternatively, grazing incidence wide-angle X-ray scattering (GIWAXS) can be used for detecting the layer alignment in MoS₂ thin films. Still, this technique is not widely used. We present here that Raman spectroscopy (in combination with GIWAXS measurements) can provide information equivalent to that from GIWAXS or TEM on the layer alignment. Raman spectroscopy is a fast method, easy to use and needs no special sample preparation. We present measurements of the intensity of A_{1g} and E_{2g} lines as a function of the angle between the polarisation of the incident and scattered light in backscattering geometry. From the measurements, a depolarisation ratio for both Raman modes was obtained, showing its distinctive values for the two layer alignments indeed. We demonstrate the method by specifying the alignment of MoS₂ grown on a sapphire substrate partially coated by reduced graphene oxide flakes.

Author: M. HULMAN, ¹ M. SOJKOVÁ ¹, K. VÉGSÖ, ² P. ŠIFFALOVICĎ, ² N. MRKÝVKOVÁ

Presenter: HULMAN M. (Institute of Electrical Engineering, SAS, Bratislava)

Session Classification: Parallel sessions

Contribution ID: 88

Type: **Oral**

WHAT CAN WE LEARN FROM COMPARATIVE CHARACTERIZATION OF MATERIALS ON THE MACRO- AND NANO-SCALE

Thursday 10 September 2020 09:50 (20 minutes)

The physical and chemical properties of materials tested on the macro- or nano-scale may differ dramatically. For example, when measuring the Young's modulus of elasticity, we assume an "infinite" depth of material, which is not fulfilled especially when the Young's modulus is measured at the nano-scale. Another example, the interaction of nanomaterials with biological environment and their toxicity is dependent on properties of individual nanoparticles, such as shape, surface passivation, zeta-potential (surface charge in colloidal state), chemical reactivity etc. and conflicting results about the same nanomaterial toxicity are published quite often. We decided to study those materials with our home-designed micro-spectroscopy setup combined with atomic force microscopy device equipped with several additional modules. In this contribution we would like to present our knowledge about studying various systems at nanoscale and their comparison to bulk measurements. The new field of nano-reconnaissance can help us to understand for example the chemical reactivity of nanoparticles or reasons of secondary inflammatory reactions caused by deformation of proteins in the protein corona formed on the nanoparticle in a living organism.

Author: ANNA FUČÍKOVÁ 1 , MATĚJ ČERNÝ 1 , HELENA VALENTOVÁ 2 , JAN VALENTA

Presenter: FUČÍKOVÁ A. (Department of Chemical Physics and Optics, Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Parallel sessions

Contribution ID: 89

Type: **Oral**

LASER –PRACOVNÍ NÁSTROJ 21.STOLETÍ

Thursday 10 September 2020 08:30 (35 minutes)

Laserové systémy prošly od svého objevení v roce 1960 dlouhou cestu a v posledních letech se stávají stále častěji běžně využívaným pracovním nástrojem. Centrum HiLASE Fyzikálního ústavu se od svého vzniku zaměřuje na vývoj špičkových, zejména pulsních, diodově buzených pevnolátkových laserů pro moderní průmyslové aplikace. Takové lasery s vysokou účinností, středním výkonem až 1 kW a dobou trvání pulsu v řádu nanosekund až femtosekund jsou schopny zajistit v průmyslových podmínkách výrobu rychlých polovodičových čipů, zvýšit životnost a spolehlivost leteckých součástek, vrtat vysoce přesné otvory pro moderní motory, vytvořit funkční a antibakteriální povrchy na řadě materiálů a mnoho dalších.

Centrum Hilase v současné době disponuje laserovou technologií na bázi tzv. tenkých disků, kdy laser využívá výhody účinného chlazení velkoplošných krystalů s tloušťkou pouze stovek mikrometrů, popř. technologií kryogenně chlazených keramických desek pro pulsy s vyšší energií. Základní emisní oblast v okolí 1030 nm může být technikami nelineární optiky rozšířena do viditelné i střední infračervené oblasti. Přednáška shrne stav vývoje v laserovém centru HiLASE.

Author: SMRŽ M. (Institute of Physics, Czech Academy of Sciences, Prague,)

Presenter: SMRŽ M. (Institute of Physics, Czech Academy of Sciences, Prague,)

Session Classification: Parallel sessions

Contribution ID: 90

Type: **Oral**

VÝZKUM FÚZNÍHO PLAZMATU VE VÝBOJÍCH PLAZMOVÉHO FOKUSU

Thursday 10 September 2020 09:05 (20 minutes)

V plazmovém fokusu je husté a horké plazma vytvářeno kompresí proudu procházejícího plazmatem vlastním magnetickým polem. Prezentované výsledky vedoucí v deuteriové náplni k fúzní DD reakci byly prováděny v IPPLM ve Varšavě na aparatuře PF-1000, která dosahuje neutronový zisk 10 11 A při proudu 1–2 MA. Na aparatuře je využívána komplexní rentgenová, laserová, částicová a neutronová diagnostika. Interpretace publikovaných výsledků prokázala existenci organizovaných toroidálních a sférických struktur v komprimovaném sloupci a jejich vývoj. K produkci fúzních neutronů dochází při vzniku a zániku těchto struktur o hustotě $10^{24} - 10^{25}$ v m^{-3} a teplotách 50–200 eV. Tyto struktury a jejich vývoj jsou vysvětleny pomocí uzavřených proudů s poloidní a toroidální komponentou a jejich vzájemnou přeměnou. Magnetická pole jsou generována principem magnetického dynama, podobně jako v planetách nebo ve hvězdách. Fúzní neutrony vznikají srážkami rychlých deuteronů o energiích stovek keV s klidovými deuterony. Mechanismus urychlení těchto částic je netepečný a v prezentaci je vysvětlen na principu rekonekce magnetických silokřivek filamentárních proudů. Prezentované výsledky jsou porovnávány s pozorováním energetických částic v tokamakovém, laserovém a astrofyzikálním plazmatu.

This work was supported in part by the Research Program under Grants MSMT No. LLT17015, 8JPL19014, CZ.02.1.01/0.0/0.0/16019/0000778 and LTAUSA17084, GACR 19-02545S, IAEA CRP RC-19253 and SGS 16/223/OHK3/3T/13.

Author: PAVEL KUBES 1, MARIAN PADUCH 2, MAREK JAN SADOWSKI 2-3, JAKUB CIKHARDT 1, DANIEL KLIR 1, JOZEF KRAVARIK 1, KAREL REZAC 1, KRZYSZTOF TOMASZEWSKI 3, ROCH KWIATKOWSKI 3, EWA ZIELINSKA

Presenter: KUBEŠ P.

Session Classification: Parallel sessions

Contribution ID: 91

Type: **Oral**

PLASMALAB@CTU - NEW FACILITIES IN SUPPORT OF FUSION EDUCATION

Thursday 10 September 2020 09:25 (20 minutes)

The contribution focuses on a new plasma laboratory, currently in commissioning phase. Its aim is to provide doctoral and undergraduate students as well as non-fusion experts with hands-on experience and support for their experimental research relevant to modern fusion technology. The laboratory consists of four main sections: Electric and magnetic fields, Plasma, Optics, and the GOLEM tokamak [1]. All the experiments support remote control operation as much as feasible. The contribution will present the technical solutions of the particular experiments and first measurements.

Author: JANA BROTKÁOVÁ 1 , JAN MLYNÁŘ 1,2 , MIROSLAV PFEIFER 2,3 , VOJTĚCH SVOBODA

Presenter: BROTKÁOVÁ J. (FJFI ČVUT v Praze)

Session Classification: Parallel sessions

Contribution ID: 92

Type: **Oral**

ANALYZING MYSTERIOUS TURBULENCE

Thursday 10 September 2020 09:45 (20 minutes)

Turbulence is one of the most fascinating phenomena in nature existing from Ångström length-scales (quantized vortices in superfluid helium [La Mantia et al. 2016]) up to galactic scales [Vatistas 2010]. One of the main features of turbulence is the presence of interacting vortices of various strengths sizes and orientations. But the present description of turbulence is based on statistical analysis of velocities or pressures. The vortex filament method is used in numerical simulations of superfluid [Varga et al. 2017] or classical viscosity-damped [Marchevsky 2020] turbulent flows. We introduce the approach of individual vortices into experiments. We use the spatially resolved 2D velocity data obtained by using the experimental method Particle Image Velocimetry (PIV) to find the individual vortices in the turbulent field and then we analyze their properties statistically.

Author: DANIEL DUDA, VITALII YANOVYCH AND VÁCLAV URUBA

Presenter: DUDA D. (Faculty of Mechanical Engineering, University of West Bohemia in Pilsen)

Session Classification: Parallel sessions

Contribution ID: 93

Type: **Oral**

VZŤAHY MEDZI ELEKTRÓNOVOU ŠTRUKTÚROU A MIKROŠTRUKTÚROU TENKÝCH VRSTIEV POLYFLUORÉNU A KOPOLYMÉRU ZALOŽENÉHO NA FLUORÉNE

Thursday 10 September 2020 10:30 (20 minutes)

V posledných rokoch vzrástol význam syntézy konjugovaných kopolymérov spájajúcich v jednej molekule rôzne monoméry. Takéto materiály nachádzajú široké uplatnenie v mikroelektronike aj vo fotovoltike, keďže ich cieľaná syntéza umožňuje nielen vhodné nastavenie šírky zakázaného pásu, ale aj zvýšenie účinnosti konverzie slnečného žiarenia na elektrickú energiu spojením monomérov s dierovou a elektrónovou vodivosťou do jednej molekuly. Aj keď v organických polovodičoch dominujú molekulové vlastnosti nad štruktúrnymi, vzťahy medzi mikroštruktúrou tenkých vrstiev kopolymérov a ich elektrickými vlastnosťami sú silné a stále nie plne pochopené. V tomto príspevku porovnáme elektrónovú štruktúru a mikroštruktúru tenkých vrstiev modelového homopolyméru a kopolyméru syntetizovaného na jeho základe. Metódami ER-EIS a GIWAXS budeme analyzovať vplyv žihania na vlastnosti vrstiev, menovite na ich elektrónovú štruktúru a kryštalinitu.

Author: KATARÍNA GMUCOVÁ 1 , KAROL VÉGSÖ 1 , VOJTECH NÁDAŽDY 1 , TOMÁŠ VÁRY

Presenter: GMUCOVÁ K. (Fyzikálny ústav SAV, Bratislava)

Session Classification: Parallel sessions

Contribution ID: 94

Type: **Oral**

MAGNETIC, MAGNETOCALORIC, THERMAL AND TRANSPORT PROPERTIES OF $Gd_3Ni_2In_4$

Thursday 10 September 2020 10:50 (20 minutes)

The new polycrystalline intermetallic compound $Gd_3Ni_2In_4$ has been prepared. We have investigated the structural, magnetic, magnetocaloric, thermodynamic and transport properties. X-ray powder diffraction pattern displays that $Gd_3Ni_2In_4$ crystallizes in hexagonal $Lu_3Co_2In_4$ -type of crystal structure. The presence of two magnetic transitions, antiferromagnetic $T_N = 21$ K and ferromagnetic T_C at 55.5 K was observed in magnetization studies. The maximal value of magnetic entropy change, $-\Delta S_M$, determined from isothermal magnetization data in a magnetic field of 9 T is 4.57 J/kg K, which is spread over a wide temperature range ($\Delta T = 61.5$ K) and hence it yields to a relative cooling power (RCP) of 281 J/kg. In addition, the compound shows a significant positive magnetoresistance, $MR(T = 2\text{ K}) = 44\%$ in magnetic field $B = 9$ T. Taking into account these results one could conclude that $Gd_3Ni_2In_4$ compound is exhibiting successive reversible magnetic transitions. Thus, it may comprise a distinct class of magnetocaloric materials as they work in a wider temperature range than conventional refrigerant materials.

Author: K. ARUN ¹, T.P. RASHID ¹, MARIAN REIFFERS ², ANDREA DZUBINSKA ³, R. NAGALAKSHMI ¹

Presenter: REIFFERS M. (Faculty of Humanities and Natural Sciences, Prešov University, Prešov,)

Session Classification: Parallel sessions

Contribution ID: 95

Type: **Oral**

MICROSCOPIC ORIGIN OF LINEAR MAGNETO-OPTICAL EFFECT IN BCC FE

Thursday 10 September 2020 11:10 (20 minutes)

We present analysis of microscopic origin of magneto-optical spectra from the electronic structure of bcc Fe. The magneto-optical (MO) permittivity spectra are obtained by the WIEN2k code and compared to the corresponding experimental MO Kerr effect spectra. The ab-initio spectra are given by the Kubo formula, i.e., as a result of summation over all pairs of electronic bands and integration over the Brillouin zone that all sum up to the single MO spectrum. We investigate what features of electronic structure contribute to the outgoing total MO spectrum. We have also developed several novel ways of visualizing MO-related phenomena in the Brillouin zone.

It turns out that strong MO signal comes from several isolated k-points in the Brillouin zone, where the band degeneracy is avoided by the spin-orbit interaction. There are two major types of MO contributions determined by the nature of the avoided degeneracy (given by topology of the approaching bands), that contribute differently to the total MO spectrum. Both types are visualized and their contributions to the total spectrum are demonstrated in the form of local MO spectra.

Author: ONDŘEJ STEJSKAL, MARTIN VEIS, JAROSLAV HAMRLE

Presenter: STEJSKAL O. (Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Parallel sessions

Contribution ID: 96

Type: **Oral**

ELECTRICAL TRANSPORT PROPERTIES OF Ni_{1.92}Mn_{1.56}Sn_{0.52} MAGNETIC SHAPE MEMORY MATERIAL

Thursday 10 September 2020 11:30 (20 minutes)

The Ni_{1.92}Mn_{1.56}Sn_{0.52} belongs to a family of Heusler alloys with peculiar magnetic, structural and transport properties that can be potentially useful for applications. They show structural transition from high temperature cubic austenite (A) to low temperature orthorhombic martensite (M) with significant changes of volume, magnetization, resistivity and Hall effect. We have performed study of temperature dependence of electrical resistivity, Hall coefficient and heat conductivity of Ni_{1.92}Mn_{1.56}Sn_{0.52} in temperature region between 4 K and 400 K. The

ferromagnetic ordering accompanied by sharp change of resistivity T-derivative was established below $T_C = 320$ K. Structural A-M transition at cooling, $T_{A-M} = 282$ K, induces a large increase of resistivity and vice versa, M-A transition at heating, $T_{M-A} = 294$ K, induces a decrease of resistivity. It is worth to note that A-M transition is associated with transition from ferromagnetic austenite to martensite with almost zero magnetization. The re-entrance of magnetization at $T_c = 235$ K is accompanied with any resistivity anomaly. The maximum of resistivity around 150 K is followed by a shallow T^2 resistivity dependence. An unusual character of temperature dependence of the Hall effect points to dramatic changes in electronic structure during the structural transition.

Author: JIŘÍ KAŠTIL¹, JIŘÍ KAMARÁD¹, OLIVIER ISNARD²

Presenter: KAŠTIL J. (Institute of Physics ASCR, Prague)

Session Classification: Parallel sessions

Contribution ID: 97

Type: **Oral**

WIDE TEMPERATURE RANGE MAGNETOCALORIC EFFECT IN DY 6.5 CO 2 SI 2

Thursday 10 September 2020 11:50 (20 minutes)

We report that this alloy has been formed in three phases Dy 5 Si 3 , Dy 3 Co 2.2 Si 1.8 , and Dy 3 Co with four successive magnetic transitions. The important role in the magnetic behavior plays competing ferromagnetic and antiferromagnetic interactions in the alloy. The number of successive magnetic transitions together with competing magnetic interactions exhibit broadened magnetocaloric effect (MCE) peak with temperature width of $\Delta T = 83.8$ and 93.2 K and consequently a large RC value of 474 and 739 J/kg, for a magnetic field change of 0–5 and 0–7 T, respectively.

Author: K. ARUN 1 T. P. RASHID 1 IVAN CURLIK 2 SERGEJ ILKOVIC 2 MARIAN REIFFERS 2 ANDREA DZUBINSKA 3 R. NAGALAKSHMI

Presenter: ILKOVIČ S. (Faculty of Humanities and Natural Sciences, Prešov)

Session Classification: Parallel sessions

Contribution ID: 98

Type: **Oral**

Advanced scanning probe microscopy of low-dimensional molecular systems revealing exotic quantum states

Thursday 10 September 2020 10:30 (35 minutes)

Low dimensional materials offer very interesting material and physical properties due to reduced dimensionality. At present, 2D materials are the focus of attention. However, 1D systems often show far more exotic features, such as Tomonaga-Luttinger liquid, Peierls distortion, etc.. In this talk, we will present π -conjugated molecular chains of distinct topological classes formed on metallic surfaces by on-surface synthesis, whose physical properties were investigated by low temperature UHV scanning probe microscopy supported by theoretical analysis. We will reveal fundamental connection between topological phase transition and the level crossing driving a variation of the π -conjugation of polymers [1]. The proximity of the quantum phase transition, in principle, allows us to design intrinsically conductive polymers despite of the Peierls distortion.

[1] B. Cierra et al. Nature Nanotechnology. (2020). DOI: 10.1038/s41565-020-0668-7

Author: JELÍNEK P. (Institute of Physics of Czech Academy of Sciences, Prague)

Presenter: JELÍNEK P. (Institute of Physics of Czech Academy of Sciences, Prague)

Session Classification: Parallel sessions

Contribution ID: 99

Type: **Oral**

Unusual properties of piezo-resonators at millikelvin temperature range and high magnetic fields

Thursday 10 September 2020 11:05 (20 minutes)

We present unusual properties of the commercial quartz tuning forks (QTFs) operating at resonance frequencies of 32 kHz, 77 kHz and 100 kHz at temperature range below 1K and in high magnetic fields up to 7.5 T. We show that in millikelvin temperature range, the quartz tuning forks exhibit the property of the coherent oscillations. These are manifested as a temperature-dependent, extremely accurate tune-up of QTFs resonance frequencies in 9th order with very high frequency stability characterized by the low values of the Allan deviations comparable with those in lasers. Even more, we demonstrate that a normalized resonance frequency of the tuning forks manifests a universal temperature dependence, which is independent on the magnetic field strength. This feature make the QTFs a very promising low temperature thermometer in high magnetic fields in temperature range below 1 K while having the B/T ratio up to 1000. We also discuss the physical origin of the observed dependencies.

Author: MARCEL ČLOVEČKO, PETER SKYBA

Presenter: ČLOVEČKO M. (Centre of low temperature physics, Institute of Experimental Physics SAS, Košice)

Session Classification: Parallel sessions

Contribution ID: 100

Type: Oral

EFFECTS OF HIGH PRESSURE ON THE RADIOACTIVE DECAY

Tuesday 8 September 2020 14:50 (20 minutes)

It is well known that the radioactive decay can be influenced from external environment [1][2]. However we still don't have a univocal picture that describes these effects. Understanding how the radioactive decay is influenced in high density environments is useful for different fields like, for example, the relevant fusion processes in stellar environment [3] or for isotope-cycling and fractionation on Earth [4]. Therefore we prepare a test at the Nuclear Physics Institute of the Czech Academy of Science. We will use three large HPGe detectors to measure gamma-rates from a solid target of ^{22}Na under high pressure. The source of ^{22}Na decays by β^+ that presents a strong gamma line, it has reasonably long half-life ($T_{1/2} = 2.6 \text{ yr}$) and it is commercially available. This simple system may provide an optimal test-bench. Hereby we will present and discuss our preparations for the test.

REFERENCES

- [1] H. Mazaki, et. al - PHYSICAL REVIEW C VOLUME 5, NUMBER 5 MAY 1972
- [2] N. Nissim et. al - PHYSICAL REVIEW C 94, 014601 - (2016)
- [3] C. Nociforo et al. - Phys. Scr. T 150, 014028 (2012)
- [4] M.B. Andersen, et. al - Nature 517, 356 (2015)

Author: ANASTASIA CASSISA A , JAROMIR MRAZEK A , GIUSEPPE D'AGATA A , FRANCOIS DE OLIVEIRA SANTOS

Presenter: CASSISA A. (Nuclear Physics Institute of the Czech Academy of Sciences, Řež)

Session Classification: Parallel sessions

Contribution ID: **101**Type: **Oral**

PHYSICS AT FUTURE EIC EXPERIMENT

Thursday 10 September 2020 11:45 (20 minutes)

The future experimental facility EIC (Electron-Ion Collider) that will be built in Brookhaven National Laboratory in the USA introduces new opportunities for precise study in high energy nuclear physics. This contribution is an overview of the author's research related to this topic. Particularly, in this contribution, we focus on topics as nuclear shadowing, quarkonia wave function study, or exclusive and diffractive processes.

The goal is to study various nuclear effects, such as nuclear shadowing. Nuclear shadowing is a phenomenon of depletion of the cross section in comparison with a cross section on the proton target. The situation is more complicated since the interaction is not completely coherent at EIC kinematics, leading to more advanced formalism based on Green functions. The studied nuclear shadowing is universal and will be applied in fact to most of the processes on nuclear targets such as vector meson (VM) diffractive and, particularly, exclusive production. The goal in such measurement is to study gluon distribution or, generally, the universal gluon Wigner distribution. However, for that, we need to minimize the uncertainty from VM wave functions by its and its effects detailed study such as quark-antiquark potentials, Melosh spin rotation, or the structure of the VM vertex and the role of the D-wave component.

Author: KŘELINA M. (FNSPE, Czech Technical University in Prague)

Presenter: KŘELINA M. (FNSPE, Czech Technical University in Prague)

Session Classification: Parallel sessions

Contribution ID: 102

Type: **Oral**

NIHILNOVI –A NOT SO NEW APPROACH TO DATA VISUALISATION PROMOTING PORTABLE AND SIMPLE TEXT FORMATS

Thursday 10 September 2020 13:30 (20 minutes)

With an aim of promoting open-source software and text files for data storage and processing, the author presents a program [1] that attempts to resolve the less-than ideal situation in this field.

The usual approach is to either load the data into a point-and-click application for scientific plotting, or learn basics of programming and type the commands in a scientific math package. However, manually handling larger or repeated datasets is inefficient. Programming languages are inappropriate for trivial or interactive tasks. Less advanced programs re-implement only a fairly limited subset of mathematical or graphical functions used in science today. More advanced ones often bind the researcher to one particular proprietary software and data formats.

The major novelty of the program is paradoxically in that it does almost nothing new: It does not implement a new plotting library, nor does it promote its custom scripting language – Python/Matplotlib/Numpy is already a popular environment for data processing.

The talk will be practically focused, with several use cases where the concept of doing (almost) nothing new to achieve great results will be illustrated.

[1] F. Dominec: Nihilnovi source code repository (2020, 11th May), retrieved from <https://github.com/FilipDominec/nihilnovi>

Author: DOMINEC F. (Institute of Physics CAS, Prague)

Presenter: DOMINEC F. (Institute of Physics CAS, Prague)

Session Classification: Parallel sessions

Contribution ID: 103

Type: **Oral**

APPLICATION OF MACHINE LEARNING IN SINGLE CRYSTAL GROWTH

Thursday 10 September 2020 13:50 (20 minutes)

Development of a new material with required properties is a very complex task. Theoretical models for growth procedure are usually not available, at least at the beginning. Therefore, many attempts are made to achieve required properties of the material and many characterization datasets are obtained. However, the way how the physical properties of the material are affected by the growth conditions does not have to be straightforwardly evident. For such a case, machine learning can be very helpful. In this contribution, applications of several simple machine learning approaches are applied to the development process of the InGaN/GaN scintillator structure. A properly trained neural network is capable to predict luminescence properties from the growth parameters of the structure. This enables optimization of the growth parameters from empirical data only. On the other hand, understanding of underlying physics is not guaranteed but the predictions of the model can give a clue.

Author: FRANTIŠEK HÁJEK 1,2 , JIŘÍ OSWALD 1 , ALICE HOSPODKOVÁ 1 , FILIP DOMINEC 1 , ROBERT HOREŠOVSKÝ 1

Presenter: HÁJEK F. (Institute of Physics CAS, v.v.i., Prague)

Session Classification: Parallel sessions

Contribution ID: 104

Type: **Oral**

IMPACT OF MACROSCOPIC PARTICLE COMPOSITION ON GAN EPITAXIAL GROWTH MORPHOLOGY AND LUMINESCENCE

Thursday 10 September 2020 14:10 (20 minutes)

We describe macroscopic defects on InGaN/GaN multiple quantum well structures caused by accidental contamination with dust particles during the metalorganic vapour phase epitaxy. Gallium nitride and InGaN/GaN heterostructures are promising materials for many optoelectronic devices, such as light emitters, high-power and high-frequency electronics, detectors of ionizing radiation, scintillators. During the preparation of these structures, great attention is paid to optimization of the growth parameters and to reduce the density of dislocations and point defects in this material. However, only a small number of studies were performed on macroscopic defects due to particles fallen from the reactor chamber or scratches from substrate polishing. Understanding the impact of each of the contaminating elements is not only important for sample diagnostics, but it also provides insight into the complex physical and chemical processes during epitaxy. We focus on the influence of macroscopic defects on photoluminescence of GaN/InGaN multiple quantum well structures and present a Raman spectroscopy study of macroscopic defect containing regions of the samples.

Author: K. KULDOVÁ, F. DOMINEC, R. NOVOTNÝ, S. SAYEDOVÁ, F. HÁJEK, A. HOSPODKOVÁ

Presenter: KULDOVÁ K. (Institute of Physics CAS, Prague)

Session Classification: Parallel sessions

Contribution ID: 105

Type: Oral

MULTIPLE QUANTUM CRITICAL POINTS IN Ce₃PtIn₁₁

Thursday 10 September 2020 14:30 (20 minutes)

The properties of the heavy fermion compound Ce₃PtIn₁₁ are rather enigmatic. Not only the compound possesses two inequivalent Ce-sites but at ambient pressure it exhibits two successive antiferromagnetic (AFM) transitions at $T_1 = 2.2\text{K}$ and $T_N = 2\text{K}$, respectively [1]. Upon further cooling superconductivity is found with $T_c = 0.32\text{K}$. Entropy analysis conjectured the idea that the Ce₂-ions are responsible for the magnetic ordering whereas the second Ce₁-ions evokes superconductivity. Here we present our recent ¹¹⁵In NMR/NQR and specific heat results. From these we infer that Ce₃PtIn₁₁ possibly harbors two quantum critical points (QCP) i.e., zero temperature phase transitions –one close to/or at ambient pressure [2] and one to be reached by an applied hydrostatic pressure of $p_c = 1.5\text{GPa}$ [1]. Each QCP can be associated with a particular Ce-site. The critical magnetic fluctuations accompanying the QCPs are at the origin of Cooper-pairing.

*This work was supported by the Czech Science Foundation (GAČR) Grant No. 18-23606S, the EU COST Action CA16218 and by the Czech Ministry of Education under the COST project scheme, project No. LTC18024.

[1] J. Prokleška et al., Phys. Rev. B 92, 161114(R) (2015).

[2] S. Kambe et al., Phys. Rev. B. 101, 081103(R) (2020).

Author: J. FIKÁČEK^{1,2}, M. HRŮZOVÁ KRATOCHVÍLOVÁ¹, K. UHLÍŘOVÁ¹, S. KAMBE³, H. SAKAI³, Y. TOKUNAGA³, T. HATTORI³, N. HIGA³, R.E. WALSTEDT⁴, M. BRANDO⁵, AND J. CUSTERS¹

Presenter: CUSTERS J. (Faculty of Mathematics and Physics, Charles University, Prague)

Session Classification: Parallel sessions

Contribution ID: 106

Type: **Oral**

OPEN CHARM AND CHARM-TAGGED JETS PRODUCTION WITH ALICE AT LHC

Thursday 10 September 2020 14:30 (20 minutes)

Charm quarks are mostly produced in hard partonic scattering processes in the early stages of a heavy-ion collision, before the quark-gluon plasma (QGP) is formed. Because of their large mass the production cross section can be calculated using perturbative quantum chromodynamics down to zero transverse momentum. Therefore, they are ideal probes of the properties of this hot, dense, and strongly interacting medium.

Measurements of heavy-flavour tagged jets bring more relevant information of the initial parton kinematics than traditional hadron measurements and can provide information on heavy-quark energy loss in the QGP, in particular on how the radiated energy is dissipated in the medium. Moreover, the study of charm-baryon production in heavy-ion collisions provides unique information on hadronization mechanism.

This talk focuses on the latest results of charm-tagged jets and open charm production with the ALICE detector at the LHC.

Author: JAKUB KVAPIL FOR THE ALICE COLLABORATION

Presenter: KVAPIL J. (University of Birmingham, UK)

Session Classification: Parallel sessions

Contribution ID: **107**Type: **Oral**

OTEVŘENÁ DATA LHC EXPERIMENTU ATLAS

Thursday 10 September 2020 13:50 (20 minutes)

Experimenty na největším světovém urychlovači LHC v mezinárodní laboratoři CERN zaznamenávají velké množství dat ze srážek protonů a těžkých iontů. Tato data jsou ukládána ve speciálních formátech a procházejí několika kroky zpracování, než jsou vhodná pro fyzikální analýzu. Část zpracovaných dat byla některými experimenty uvolněna pro všechny zájemce. Spolu s daty jsou k dispozici i nástroje pro jejich zpracování a detailní dokumentace postupů a objasnění používaných pojmů. Na veřejně dostupných datech experimentu ATLAS ukážu možnosti jejich využití pro propagaci oboru i pro verifikaci některých výsledků nezávislými experty.

Author: CHUDOBA J.**Presenter:** CHUDOBA J.**Session Classification:** Parallel sessions

Contribution ID: **108**Type: **Oral**

ALICE ITS Upgrade

Thursday 10 September 2020 14:10 (20 minutes)

The ALICE Inner Tracking System detector is undergoing a major upgrade in order to cope with the increased data rates and to meet the requirements as set out by the physics goals of the experiment after Long Shutdown 2. The new ITS will be completely made up of ALPIDE monolithic active pixel sensors based on a CMOS 180nm process. A single sensor measures 15mm x 30mm and contains half a million pixels distributed over 512 rows and 1024 columns. These 50um thick sensors, with 27um x 29um pixel pitch, are mounted on ultra-lightweight carbon composite support structures with an embedded cooling system. This results in a considerable reduction of the material budget (down to 0.35% X_0 for the inner layers and ~1% X_0 for the outer layers) and a significant improvement of the impact parameter resolution and tracking efficiency. The innermost ITS layer will be moved as close as 23mm to the interaction point. The integration of the ITS detector assembly, made of the three innermost and four outermost layers, has been almost completed and the commissioning, first in the laboratory, is ongoing. The detector will be installed in ALICE in 2020. This talk will give a brief overview of the motivation for the upgrade and will present the first results of the detector performance obtained during the commissioning.

Author: ISAKOV ARTEM FOR THE ALICE COLLABORATION

Presenter: ISAKOV A. (Nuclear Physics Institute of the CAS, Řež)

Session Classification: Parallel sessions

Contribution ID: 109

Type: Oral

Novel high-luminosity fixed-target experiment at the LHC

Thursday 10 September 2020 13:30 (20 minutes)

Extraction of the multi-TeV proton and lead LHC beams with a bent crystal or by using an internal gas target allows one to perform the most energetic fixed-target experiment ever. pp, and pA collisions at collision energy of 115 GeV and Pb-p and Pb-A collisions at 72 GeV can be studied with high precision and modern techniques over a broad rapidity range. Using the LHCb and ALICE detectors in a fixed-target mode offers unprecedented possibilities to study, among others: the quark, gluon and heavy-quark content of the nuclei in the poorly known region of the high-momentum fractions, heavy-flavour production in a new energy domain, half way between the SPS and RHIC.

In this talk, the technical solutions to obtain a high-luminosity fixed-target experiment at the LHC will be reviewed and their possible implementations with the ALICE and LHCb detectors will be discussed. Projection studies for various observables such as Drell-Yan, charm, beauty and quarkonium production, with both detector set-ups used and with various nuclear targets and the LHC lead beams will be presented.

Author: TRZECIAK B. (Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague)

Presenter: TRZECIAK B. (Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague)

Session Classification: Parallel sessions

Contribution ID: 110

Type: Oral

Highlights in physics of two-dimensional crystals through cryomagnetic optics

Thursday 10 September 2020 14:55 (20 minutes)

The discovery of graphene –the archetype two-dimensional (2D) system, stimulated the frantic pace of research in the area of 2D materials (2DM). The rapid progress of the field, particularly in the growth technologies, is enabling expanding the research interests in other 2DM with outstanding optical, transport, magnetic and other properties giving rise to a plethora of tailored functionalities. Looking beyond the flatland, the 2D slabs can also be reassembled into designer heterostructures made layer by layer in a precisely chosen sequence. Such heterostructures (often referred to as ‘van der Waals’) reveal unusual properties driven by the stacking and mutual orientation of the 2D building blocks. In my talk, I will review the current state of knowledge on the most topical areas of 2DM research. First, the stacking phenomenon responsible for numerous exotic states ranging from superconductivity to Mott metal-insulator transition will be discussed. Next, the phenomenon of 2D magnetism will be revisited and I will review the limits imposed on boosting the material’s properties given by the Mermin–Wagner theorem. Finally, several recent experimental reports will be highlighted and future prospects in the field of 2DM will be outlined.

Author: VEJPRAVOVÁ J. (Department of condensed matter physics, Faculty of mathematics and physics, Charles University, Prague)

Presenter: VEJPRAVOVÁ J. (Department of condensed matter physics, Faculty of mathematics and physics, Charles University, Prague)

Session Classification: Plenary Talks

Contribution ID: 112

Type: **Poster**

SHORT-RANGE CORRELATION INVESTIGATION IN DEUTERON INDUCED REACTIONS

Wednesday 9 September 2020 16:20 (30 minutes)

One of the approaches to investigate the equation-of-state of dense nuclear matter is the study of the short-range correlations of nucleons in nuclei. Short-range correlations have densities comparable to the density in the center of a nucleon and they can be considered as the drops of cold dense nuclear matter. Dp elastic and dp breakup processes are investigated at Internal Target Station of Nuclotron; dp elastic process in angular range from 60 –135 degree in c.m. in the energy range from 400 - 2000 MeV; dp breakup reaction in angular the range from 19 – 56 degree from 300 - 500 MeV. Results which comes from analyzing powers of dp elastic scattering show strong sensitivity to the short range spin structure of the isoscalar nucleon-nucleon correlations. Description based on relativistic multi-scattering model provides reasonable agreement at small and large angles but the problem is related to angles between them.

Presenter: JANEK M. (Physics Department, University of Zilina)

Session Classification: Poster session

Contribution ID: 113

Type: **Poster**

The compound $[\text{Ni}(\text{2aepy})_2 \text{Cl}(\text{H}_2\text{O})] \text{Cl} \cdot \text{H}_2\text{O}$ as a candidate approaching a topological quantum critical point of a spin-1 one-dimensional antiferromagnet

Wednesday 9 September 2020 16:20 (30 minutes)

The idea of controlling the magnetic ground state of a spin-1 one-dimensional antiferromagnetic (1d AFM) quantum magnets has long been of interest to physicists. In the study of anisotropic 1d AFM systems based on Ni^{2+} compounds, a series of topologically protected quantum phases was observed, one of these phases is the topologically protected Haldane phase [1, 2]. The magnetic ground state of such a system is sensitive to the relative magnitude of the single-ion anisotropy (D) and the intrachain (J) exchange interaction parameters. The D/J ratio dictates the system's placement in one of three competing phases: a Haldane gapped phase, a quantum paramagnet, and an XY-ordered state, with a quantum critical point at their junction at $D/J = 1$. We present the study of the crystal structure and magnetic properties of compound $[\text{Ni}(\text{2aepy})_2 \text{Cl}(\text{H}_2\text{O})] \text{Cl} \cdot \text{H}_2\text{O}$ (2aepy = 2-aminoethylpyridine). Hexacoordinate Ni^{2+} ions form a zig-zag chain based on hydrogen bonds and running along crystallographic b-axis. The analysis of the experimental susceptibility using a model of spin-1 anisotropic AFM chain yielded parameter values $D/k_B = 4.05 \text{ K}$ and $J/k_B = 3.55 \text{ K}$. A theoretical prediction of $D/k_B = 3.7 \text{ K}$ using ab initio approach is very close to our experimental value.

[1] F.D.M. Haldane, Phys. Lett. A 93, 464 (1983).

[2] F.D.M. Haldane, Phys. Rev. Lett. 50, 1153 (1983).

[3] J.L. Manson, et al., Inorg. Chem. 51, 7520 (2012).

[4] D.M. Pajerowski, et al., arXiv:2001.08555.

This work was supported by VEGA 1/0426/19, APVV-18-0197, and APVV-18-0016 projects.

Presenter: HOLUB M. (Institute of Physics, Faculty of Science, P.J. Šafárik University)

Session Classification: Poster session

Contribution ID: 114

Type: **Poster**

PROPERTIES OF LIGHTNING-INDUCED ELECTROMAGNETIC WAVES DETECTED CLOSE TO JUPITER

Wednesday 9 September 2020 16:20 (30 minutes)

The Waves instrument onboard the Juno spacecraft recorded numerous cases of dispersed electromagnetic waves generated by Jovian lightning. These waves, also known as whistlers, propagated through ionospheric and magnetospheric plasmas before their detection by Juno during its close approaches to Jupiter. We present observations of nearly five thousand low-dispersion lightning whistlers detected below radial distances of 5 Jovian radii during the first half of the Juno mission. We have found that Jovian lightning discharges occur predominantly at mid-latitudes. We have found an asymmetry in the whistler occurrence in both hemispheres: the average lightning stroke rate in the northern hemisphere was approximately twice higher than in the southern hemisphere in the first quarter of the mission, but the mysterious asymmetry surprisingly disappeared in its second quarter. This effect might be explained by a random distribution of thunderstorms. A lack of whistlers in the tropics might be a consequence of their propagation in field-aligned ducts which would not allow them to reach the altitude of Juno. Generation of lightning at Jupiter's equatorial regions might be also suppressed due to the solar radiation received at Jupiter's equator, which actually inhibits the rise of warm air from within the planet into the upper atmosphere.

Presenter: KOLMAŠOVÁ I. (Department of Space Physics, Institute of Atmospheric Physics of the Czech Academy of Sciences)

Session Classification: Poster session

Contribution ID: 115

Type: **Poster**

EXOMARS 2022: PLANNED MEASUREMENTS OF ELECTROMAGNETIC RADIATION ON THE SURFACE OF MARS

Wednesday 9 September 2020 16:20 (30 minutes)

Dust grains in the Martian dust storms or dust devils may be electrically charged by triboelectric effects and laboratory experiments show that under specific conditions electric discharges might occur in the dusty Martian atmosphere. Remote measurements from the Earth using a 34-m Deep Space Network antenna have shown a non-thermal component of electromagnetic radiation from Mars which has been attributed to the effects of discharges in the dust storms but observations of the radar receiver onboard the Mars Express spacecraft showed no credible radio signals from Martian lightning between 4 and 5.5 MHz. Direct measurements of electromagnetic radiation on the surface of Mars are needed to solve this puzzle.

The Exomars 2022 Surface Platform instrumentation will include the Wave analyzer module, consisting of an assembly of magnetic and electric antennae and dedicated electronics, as a part of the Martian ground electromagnetic tool instrument. The module will be dedicated to the measurement of electromagnetic field fluctuations in the frequency band from 100 Hz to 8 MHz. We plan to experimentally investigate possible radio emissions of atmospheric origin generated by electrical discharges, as well as electromagnetic waves linked to the interactions of interplanetary plasma medium with the Martian ionosphere and magnetic anomalies.

Presenter: SANTOLÍK O. (Department of Space Physics, Institute of Atmospheric Physics of the Czech Academy of Sciences)

Session Classification: Poster session

Contribution ID: 116

Type: **Poster**

MONTE CARLO SIMULACE PRODUKCE UPSILON MEZONU

Wednesday 9 September 2020 16:20 (30 minutes)

Kvarkonia jsou důležitou sondou k vyšetřování vlastností kvark-gluonového plasmatu. Proton-protonové srážky jsou nezbytným prostředkem k ustanovení základních vlastností, které slouží ke studiu kvarkonií v proton-jaderných a jádro-jaderných srážek. Tento poster prezentuje základní charakteristiky Upsilon mezonů zjištěné pomocí Monte Carlo generátorů v proton-protonových srážkách při $\sqrt{s_{NN}} = 500$ GeV. Monte Carlo generátory srážek PYTHIA a Herwig byly využity k generování dat. Hlavním cílem těchto simulací je výzkum závislosti normalizované produkce Upsilon mezonů na normalizované multiplicitě. Závislost na normalizované multiplicitě je vhodným nástrojem k porozumění mechanismům produkce částic a souhry měkkých a tvrdých procesů kvantové chromodynamiky.

Presenter: ČEŠKA J. (FJFI, České Vysoké Učení Technické v Praze)

Session Classification: Poster session

Contribution ID: 117

Type: **Poster**

PHASE TRANSFORMATIONS IN Al-Li-BASED ALLOY STUDIED BY IN-SITU TEM.

Wednesday 9 September 2020 16:20 (30 minutes)

Al-Li based alloys of AA2195 type are designed mainly for aerospace applications. They exhibit a very high strength due to a presence of strengthening precipitates of a nanometric size. However, materials prepared by a conventional direct-chill casting and following rolling exhibit strong crystallographic texture and inhomogeneous mechanical properties. This inconvenient behavior could be suppressed by a direct twin-roll casting of the material to a final gauge and subsequent precipitation annealing. Phase transformations occurring at selected temperatures could be characterized using in-situ transmission electron microscopy (TEM) annealing experiments. Nevertheless, the processes could be significantly influenced by a constrained volume of thin foils used in TEM. A formation of platelike Li and Cu-rich non-equilibrium precipitates was studied in materials annealed conventionally in a furnace and in-situ in TEM.

Presenter: CIESLAR M. (Charles University, Faculty of Mathematics and Physics)

Session Classification: Poster session

Contribution ID: 118

Type: **Poster**

RECRYSTALLIZATION OF TWIN-ROLL CAST AA8079 ALUMINUM ALLOY AFTER HOMOGENIZATION

Wednesday 9 September 2020 16:50 (30 minutes)

Twin-roll casting of AA8079 aluminum sheets creates non-equilibrium structure as a result of high solidification rates. Homogenization treatments consisting of an exposure of the as-cast material to high temperatures close to the melting point result in a redistribution of solute atoms and a formation of a more stable structure. Newly formed equilibrium particles have different crystallographic structure, size and also their distribution is more homogeneous. Their features have a significant influence on softening processes and recrystallization occurring in sheets rolled from the homogenized material. Two different processes controlling the kinetics of recrystallization were identified by electron microscopy, light optical microscopy and microhardness measurements. It was shown, that a particle stimulated nucleation dominates in materials homogenized at high temperatures while a Zener drag is a controlling process in sheets exposed to lower homogenization temperatures.

Presenter: KRÁLÍK R. (Charles University, Faculty of Mathematics and Physics)

Session Classification: Poster session

Contribution ID: 119

Type: **Poster**

CHARACTERIZATION OF THIN FREE-STANDING AL-MG FILMS

Wednesday 9 September 2020 16:50 (30 minutes)

Small-scale thin films are frequently used in microelectronic devices and micro-electro-mechanical systems where they are commonly subjected to high strains during their dynamic motion. However, mechanical behavior of thin films significantly differs from the behavior observed in bulk materials. To understand the size dependent properties of deformation mechanisms, characterization of the grain properties of nano-scale materials is essential along with mechanical tests. A promising method is in-situ transmission electron microscopy deformation, which combines mechanical tests on a nano-scale with a direct observation of the structure even with an atomic resolution. Thin Al-based free-standing films prepared by DC magnetron sputtering were characterized by atomic force microscopy, conventional and in-situ transmission electron microscopy and automatic phase and orientation mapping in TEM.

Presenter: BAJTOŠOVÁ L. (Charles University, Faculty of Mathematics and Physics)

Session Classification: Poster session

Contribution ID: 120

Type: **Poster**

INTERDIFFUSION IN ALUMINIUM –STEEL CLAD

Wednesday 9 September 2020 16:50 (30 minutes)

Bonding of aluminium and steel into one single material –an aluminium-steel clad – represents an excellent composite material, which provides high potential for applications in automobile industry by combination of properties of respective materials –high stiffness and strength of steel and durability, corrosion resistance, thermal capacity and high specific strength of aluminium. microstructure. Optimal process conditions during clad strips production should result in the formation of a continuous diffusion layer of several Fe-Al phases. However, such layer may reduce the bonding strength and formability of the material. Diffusion and phase transformations in aluminium - steel clad sheet prepared by twin-roll casting were studied by means of light optical microscopy, electron microscopy and resistometry. Effective interdiffusion coefficient was evaluated by Boltzmann-Matano method from measured concentration profiles through the interface. Simulated results were in a direct contradiction with in-situ TEM/SEM observations showing surprisingly a formation of Fe-Al phases in the steel layer. Results confirm that a simple diffusion driven model could not cover complex processes occurring at the interface.

Presenter: KŘIVSKÁ B. (Charles University, Faculty of Mathematics and Physics)

Session Classification: Poster session

Contribution ID: 121

Type: **Poster**

INTERDIFFUSION IN ALUMINIUM –STEEL CLAD

Wednesday 9 September 2020 16:50 (30 minutes)

Bonding of aluminium and steel into one single material –an aluminium-steel clad – represents an excellent composite material, which provides high potential for applications in automobile industry by combination of properties of respective materials –high stiffness and strength of steel and durability, corrosion resistance, thermal capacity and high specific strength of aluminium. microstructure. Optimal process conditions during clad strips production should result in the formation of a continuous diffusion layer of several Fe-Al phases. However, such layer may reduce the bonding strength and formability of the material. Diffusion and phase transformations in aluminium - steel clad sheet prepared by twin-roll casting were studied by means of light optical microscopy, electron microscopy and resistometry. Effective interdiffusion coefficient was evaluated by Boltzmann-Matano method from measured concentration profiles through the interface. Simulated results were in a direct contradiction with in-situ TEM/SEM observations showing surprisingly a formation of Fe-Al phases in the steel layer. Results confirm that a simple diffusion driven model could not cover complex processes occurring at the interface.

Presenter: KŘIVSKÁ B.**Session Classification:** Poster session

Contribution ID: 122

Type: **Poster**

STATISTICAL ERROR OF FP EQUATION SOLUTION FOR COSMIC RAYS DISTRIBUTION IN THE HELIOSPHERE

Wednesday 9 September 2020 16:50 (30 minutes)

Cosmic rays propagate the heliosphere from interstellar space till orbit of planets and Sun. The problem of their distribution, described by Fokker-Planck equation is solved by a couple of methods, one of the most used is the stochastic method based on Ito lemma. Presented work is focused on the estimation of statistical error of Fokker –Planck equation solution of the 1D forward stochastic method, for evaluation cosmic rays distribution in the heliosphere. Error dependence on simulation statistics and energy for 1AU is presented. The 1% precision criterium is defined as a function of solar wind velocity and diffusion coefficient value. The systematic error of the FP equation is also discussed.

Presenter: MICHAJLENKO V.

Session Classification: Poster session

Contribution ID: 123

Type: **Poster**

COR SYSTEM FOR COSMIC RAYS TRAJECTORIES IN MAGNETOSPHERE SIMULATION

Wednesday 9 September 2020 16:50 (30 minutes)

Cosmic rays are particles, coming from space. The system COR (Cut-off rigidity) provides access for the wider scientific community to models of simulation of cosmic ray trajectories in the magnetosphere via a web interface. The system offers simulations of vertical directions or, from multiple nonvertical directions covering half sphere (2π solid angle) with the center of the sphere in the point of interest. The simulation particle tracing is realized in combined internal (IGRF) and the external geomagnetic field (Tsyganenko 96 or Tsyganenko 05) covering the years 1968 to 2020. We call this Standard simulation module. There is also a module for simulation in an earlier period called Historic simulation module that uses a couple of geomagnetic field approximations for last two millennia (years from 0 to 1968). The system also contains individual trajectory visualizations for standard modules and a magnetosphere simulation module that allows users to evaluate the magnetic field in the magnetosphere for selected space and time range. Simulations based on space and time ranges are also available for Standard and Historical cosmic ray trajectories models.

Presenter: GECASEK D.

Session Classification: Poster session

Contribution ID: 124

Type: **Poster**

SEARCH FOR PSEUDORAPIDITY FLUCTUATIONS IN HIGH ENERGY NUCLEAR COLLISIONS

Wednesday 9 September 2020 16:50 (30 minutes)

Emission of relativistic particles produced in central nuclear collisions in emulsion detector has been studied for different beam energies and primary nucleus - 16 O, 20 Ne, 28 Si, 32 S, 197 Au and 208 Pb. A preliminary study of multiplicity and target dependence of particles fluctuations on pseudorapidity distribution in terms of the quantity χ has been made. Search for the event-by-event fluctuations of observable χ and its dependence on the mass and energies of colliding nuclei, and the degree of centrality has been performed. The comparison of experimental data with model predictions has been made.

Presenter: VRLÁKOVÁ J.**Session Classification:** Poster session

Contribution ID: 125

Type: **Poster**

DEDICATED RUNAWAY ELECTRONS STUDIES AT THE TOKAMAK COMPASS

Wednesday 9 September 2020 16:50 (30 minutes)

Runaway electrons (RE) present one of the major concerns for future fusion devices (tokamaks), since the uncontrolled localized RE losses can compromise the integrity of tokamak device and its further operation. Runaway electrons can emerge within a tokamak plasma either at low density, during the discharge start-up phase or as a result of rapid plasma termination, associated with sudden cooling causing a significant electric field induction. In the presence of such a strong electric field, electrons can reach several tens of MeV (kinetic energy \sim tens of MJ). The fully effective mitigation strategy was still not identified despite the increased effort of the fusion community. Several approaches have to be explored.

The COMPASS tokamak is an ideal device for testing such strategies and developing new ones due to its high flexibility, advanced RE feedback, compact size and the high reproducibility of RE scenarios. The main studied mitigation strategies are the injection of impurities and the application of external resonant magnetic perturbations (RMPs). The combination of mentioned mitigation strategies with emphasis on RMPs will be examined. Conclusions about their efficiency at COMPASS will be elaborated and basic assumptions toward larger fusion devices will be derived.

Presenter: MACUSOVA E.

Session Classification: Poster session

Contribution ID: 126

Type: **Poster**

DESIGN OF NOVEL MATERIALS: ARPES EXPERIMENTS AND THEORY

Wednesday 9 September 2020 16:50 (30 minutes)

Angle-resolved photoemission spectroscopy (ARPES) is a leading experimental probe for studying the electronic structure and complex phenomena in quantum materials. Modern experimental arrangements consisting of new photon sources, analyzers and detectors supply not only spin resolution but also extremely high angle and energy resolution [1]. Furthermore, the use of photon energies from few eV up to several keV makes this experimental technique a rather unique tool to investigate the electronic properties of solids and surfaces [2]. On the theoretical side, it is quite common to interpret measured ARPES data by simple comparison with calculated band structure. However, various important effects, like matrix elements, the photon momentum or phonon excitation, are in this way neglected. Here, we present a generalization of the state of the art description of the photoemission process, the so called one-step model that describes excitation, transport to the surface and escape into the vacuum in a coherent way [3,4]. Nowadays, the one-step model allows for photocurrent calculations for photon energies ranging from a few eV to more than 10 keV, for finite temperatures and for arbitrarily ordered and disordered systems, and considering in addition strong correlation effects within the dynamical mean-field theory. Application of this formalism in order to understand ARPES response of new materials like low-dimensional magnetic structures [6], Rashba systems [5], topological insulator materials [1], materials relevant for photo-catalysis [8] or ultrafast femtosecond spin dynamics [7] will be shown. In this presentation I review some of the recent ARPES results and discuss the future perspective in this rapidly developing field. In addition I will introduce our new spin polarised ARPES laboratory.

Presenter: MINAR J.**Session Classification:** Poster session

Contribution ID: 127

Type: **Poster**

Výzkum přechodu do režimu zlepšeného udržení plazmatu na tokamaku COMPASS

Wednesday 9 September 2020 16:50 (30 minutes)

Provoz v režimu s vysokým udržením energie plazmatu (H-mód) je jedním z klíčových předpokladů pro úspěšné dosažení hlavního cíle tokamaku ITER, jímž je desetinásobné fúzní zesílení výkonu. Přechod do H-módu však může být významně ztížen v důsledku přítomnosti chybových polí, které v tokamaku nevyhnutelně vznikají jako důsledek nepřesností v umístění magnetických cívek či centrálního solenoidu. Tokamak COMPASS je vybaven světově unikátní sadou cívek pro kontrolu chybových polí, umístěných na vnější i vnitřní straně tokamaku, a v nedávných experimentech prokázal, že za určitých okolností vede přítomnost těchto chybových polí během přechodu do H-módu k zamknutí rotace magnetických ostrovů a následnému kolapsu celého plazmatu. Dále se experimentálně podařilo prokázat vliv polohy divertorové singularity magnetického pole na dosažení H-módu skrze ovlivnění síly toků generovaných elektrickým polem, které rozbíjí turbulentní struktury zodpovědné za většinu energetických ztrát. V neposlední řadě se připravují unikátní experimenty pro měření potenciální vorticity, která souvisí se samo-organizací zonálních toků, které regulují míru turbulence a mohou způsobit přechod do H-módu.

Presenter: PETERKA M.**Session Classification:** Poster session

Contribution ID: 128

Type: **Poster**

STUDY OF RING-LIKE STRUCTURES IN PARTICLE EMISSION IN RELATIVISTIC NUCLEAR INTERACTIONS

Wednesday 9 September 2020 16:50 (30 minutes)

The substructures in the emission of relativistic particles in central ^{197}Au and ^{208}Pb interactions with Ag, Br targets in nuclear emulsions at beam momenta 11.6 and 158 A GeV/c, respectively, have been investigated. The nonstatistical ring-like substructures of produced particles in azimuthal plane of a collision as result of hydrodynamic waves in nuclear matter have been searched and their parameters have been determined. The experimental results have been compared with the results simulated by Monte Carlo method.

Presenter: KRAVCAKOVA A.

Session Classification: Poster session

Contribution ID: 129

Type: **Poster**

NANOFLUID BASED ON A NEW GENERATION OIL

Wednesday 9 September 2020 16:50 (30 minutes)

Magnetic fluids belong to special nanomaterials with many application possibilities. One of the fields of their use is the energy industry, where the demands on the performance of electrical equipment are constantly rising. As a result, there is a need to use modern insulation systems. Adding nanoparticles to conventional media could be one way to obtain them. In this study, we focused on the complex characterization of potential magnetic nanofluids based on isoparaffin hydrocarbons extracted from natural gas. These new generation gas-to-liquid (GTL) derived electrical insulation fluids are cleaner, chemically stable, and the thermal conductivity values of these electrical insulating fluids are higher, indicating improved heat transfer properties. For the preparation of magnetic nanofluids, we used sterically stabilized nanoparticles of iron oxide with a volume fraction in samples in the range of 0.3–3.0%. The saturation magnetization, magnetic susceptibility, density, and viscosity were investigated showing a significant enhancement with an increase of the volume fraction. Our experimental results show that the nanoliquid prepared on the basis of the GTL technology has a stable colloidal character that is necessary for its application as a potential cooling medium in electro-energetics.

Presenter: PAULOVICHOVÁ K.**Session Classification:** Poster session

Contribution ID: 130

Type: **Poster**

DIELECTRIC SPECTROSCOPY OF TRANSFORMER OIL-BASED FERROFLUIDS WITH MN-ZN FERRITE NANOPARTICLES

Wednesday 9 September 2020 17:20 (30 minutes)

Transformers are one of the significant parts of the electrical power system. As the power transmission system is opting for high voltage high power transmission, high performance insulating materials are drawing attention of the electrical power industry. Nanofluids, formed by adding nanoscale particles to insulating oil, are stable and homogeneous suspensions that present advanced performance of electrical insulation and heat dissipation. Transformer oil-based ferrofluids with Mn-Zn ferrite nanoparticles have been characterized at different concentrations of particles. The dielectric-spectroscopy experiments were performed on the LCR meter in the frequency range up to 2 MHz for different AC and BC voltage. We demonstrate changes in the observed relaxation process by applying electric field with different intensity. Dielectric dissipation factor of ferrofluids samples was also compared. The low-frequency relaxation process features observed in experiments was assigned to the electric double layer polarization. How to reduce the negative effect of nanoparticles on dielectric loss of nanofluids, needs to be investigated further.

CANCELLED!

Presenter: KARPETS M.

Session Classification: Poster session

Contribution ID: 131

Type: **Poster**

Super-Klein tunneling of Dirac fermions through electrostatic gratings in graphene

Wednesday 9 September 2020 17:20 (30 minutes)

We use the Wick-rotated time-dependent supersymmetry to construct models of two-dimensional Dirac fermions in presence of an electrostatic grating. We show that there appears omnidirectional perfect transmission through the grating at specific energy. Additionally to being transparent for incoming fermions, the grating hosts strongly localized states.

Presenter: JAKUBSKÝ V.**Session Classification:** Poster session

Contribution ID: 132

Type: **not specified**

H-function Calculation for Selected Spectral Bands of Oxygen

Wednesday 9 September 2020 17:20 (30 minutes)

The H-function is the most important part for determining the absorption coefficient. This article deals with the calculation of the H-function of selected bands of molecular oxygen. Program calculations are performed on the basis of freely available data from world databases (NIST) and book sources (e.g. Glushko, Rosen) providing the possibility of comparison. For the structural calculations of oxygen bands, the Orca program, created at the Max-Planck Institute by Frank Nees, et al., was selected. The outputs of this program are used by our program NKrov2, which provides the final mathematical and graphical outputs.

Presenter: POKORNÝ J.

Session Classification: Poster session

Contribution ID: 133

Type: **not specified**

PHASE DIAGRAMS OF Ce₂Pd₂In INTERMETALLIC COMPOUND

Wednesday 9 September 2020 17:20 (30 minutes)

Thanks to the specific electronic structure the rare earth-based compounds, especially those containing Yb, Ce or Eu, often exhibit exceptional magnetic properties. In our study we have focused on cerium-based compound Ce₂Pd₂In belonging to the family of R₂T₂X compounds crystallizing in tetragonal Mo₂FeB₂-type structure. Previous studies revealed presence of two magnetic phase transitions ($T_C \approx 4.1$ K and $T_N \approx 4.5$ K) and strong dependence

of magnetic ground state on the changes of chemical composition [Giovannini, PhysRevB2000].

We present the pressure-temperature phase diagrams both for hydrostatic and uniaxial pressure. Hydrostatic pressure acts in the same way on the whole lattice, while the uniaxial one allows to act solely in the chosen direction. The results are put into context with temperature evolution of crystal lattice investigated by low temperature X-ray diffraction. Based on these results, hydrostatic pressure is supposed to act more on the a-parameter, which leads to approaching of atoms in the basal plane, affecting the exchange interactions in the system and preference antiferromagnetic phase over the ferromagnetic one. On the other hand, the uniaxial pressure acts on the parameter c showing no significant effect on the temperatures of phase transition.

Presenter: KRÁL P.**Session Classification:** Poster session

Contribution ID: 134

Type: **Poster**

Pressure induced superconductivity in a CeRhSi₃ single crystal –high pressure study

Wednesday 9 September 2020 17:20 (30 minutes)

Pressure-induced superconductivity in CeRhSi₃ and CeIrSi₃ has attracted a significant attention for unconventional nature of the superconductivity in a non-centrosymmetric lattice. All previous results, however, were limited to maximum 3 GPa of applied pressure. We focus on a high-pressure-region behavior of the pressure-induced superconductivity in CeRhSi₃. Our study was performed employing the good-quality Sn-flux-grown single-crystal (electrical current along [110]) and Bridgman anvil cell allowing to apply pressures up to 6 GPa. The initial shift of antiferromagnetic transition to higher temperatures with applied pressure, up to 1.1 GPa; emergence of superconductivity at this pressure; subsequent decrease of Néel temperature and increase of SC temperature with further pressure application were followed. The critical SC temperature reaches a maximum at 2.9 GPa. No signs of magnetic transition are observed. Further application of pressure shifts SC to lower temperatures, forming a typical SC dome. The superconductivity is expected to be completely suppressed between 5 and 6 GPa. Measurements in magnetic fields revealed a considerable decrease of critical field above 3 GPa, which exceeds the value of 19 T at the top of SC dome. Presented results are summarized in the completed T-p and H-T phase diagrams, complementing previous results.

Presenter: STAŠKO D.**Session Classification:** Poster session

Contribution ID: 135

Type: **Poster**

A SETUP FOR MEASUREMENT OF THE TOTAL REACTION CROSS SECTION

Wednesday 9 September 2020 17:20 (30 minutes)

Measurements of energy dependence of the total reaction cross section with exotic, neutron-rich nuclei are used to study the phenomena related to the structure of weakly bound neutrons in neutron halo and neutron skin. A spectrometer „MULTI-2“ for direct measurement of the total reaction cross section with radioactive beams is presented, together with some results obtained on the setup. The spectrometer consists of a multi-detector telescope for the beam projectile identification and a 4π gamma-ray spectrometer for detection of prompt gamma-rays and neutrons accompanying nuclear reactions. The characteristics of the gamma-ray spectrometer important for the evaluation of the measurements –registration efficiency as a function of energy and multiplicity of emitted gamma rays, spectrometer response and a total energy deposition were evaluated with Monte Carlo method. Results were confirmed by measurement with ^{60}Co spectroscopic source.

Presenter: SIVÁČEK I.**Session Classification:** Poster session

Contribution ID: 136

Type: **Oral**

Current development in the study of magnetically induced crystal reorientation of Ni-Mn-Ga ferromagnetic shape memory alloy

Tuesday 8 September 2020 17:35 (20 minutes)

In the past two decades Ni-Mn-Ga Heusler alloys have received great attention as these exhibit magnetic shape memory effects, first described in 1999. Modulated Ni₂MnGa martensite has exceptionally low stress for pseudoplastic deformation or structural reorientation by twinning mediated by highly mobile twinning boundaries [1]. Owing to large magnetic anisotropy and low twinning stress the large pseudoplastic deformation can be induced by a magnetic field. Thus, these materials can replace giant magnetostrictive materials with fast external actuation control and strain exceeding 10 % [1].

In this study, the energy needed for the deformation by a magnetic field and by mechanical force was measured and compared in monocrystalline Ni₅₀Mn₂₈Ga₂₂ modulated martensite. Measurements were conducted on five samples by vibration sample magnetometer and stress-strain device to compare directly different modes of loading. In contrast with the phenomenological model [2] the calculated energy of magnetically induced reorientation or pseudoplastic deformation were considerably higher compared to the energy needed using mechanical force. The switching field of samples with nucleated twinning boundaries was also measured and compared to single variant crystals. Results only partly support modelling of the twinning stress and switching field [1-3]. The model-data discrepancies are an unresolved issue suggesting a need for modification of the model.

Author: ŠIMON SUKUP^{1,2*}, SAMUEL HECZKO³ AND OLEG HECZKO

Presenter: HECZKO O. (Department of Magnetic Measurements and Materials, Institute of Physics CAS, Prague)

Session Classification: Parallel sessions

Contribution ID: 137

Type: **not specified**

Registration for teachers

Tuesday 8 September 2020 13:00 (30 minutes)

Registrace učitelů

Contribution ID: **138**

Type: **Poster**

Zahájení vernisáže BeInspired

Wednesday 9 September 2020 15:30 (15 minutes)

Contribution ID: **139**

Type: **Poster**

Poster session - opening

Wednesday 9 September 2020 15:45 (5 minutes)

Session Classification: Poster session

Contribution ID: 140

Type: **Poster**

STUDY OF P3HT THIN FILM PHOTOLUMINESCENCE SPECTRA USING H-J AGGREGATE THEORY

Wednesday 9 September 2020 17:20 (30 minutes)

The usual interband transitions S₀-S₀, S₀-S₁, S₀-S₂ can be clearly identified in the recently published photoluminescence measurements on P3HT layers. However, in many cases the spectra are characterized by weaker ripples besides these main features. We point out their possible interpretation by means of the H and J aggregates theory proposed by Spano (F. C. Spano, J. Chem. Phys. 122, 234701-234715, 2005). Such an explanation correlates with the fact that the P3HT material in a spin-coated layer consists of the amorphous phase and one or two crystalline phases depending on the substrate surface properties.

Author: Dr KOTOROVÁ, S. (Faculty of Electrical Engineering and Information Technology, Slovak University of Technology, Ilkovičova 3, Bratislava, Slovakia)

Co-authors: CIRÁK, J. (Faculty of Electrical Engineering and Information Technology, Slovak University of Technology, Ilkovičova 3, Bratislava, Slovakia); Mr VÁRY, T. (Faculty of Electrical Engineering and Information Technology, Slovak University of Technology, Ilkovičova 3, Bratislava, Slovakia); Mr CHLPÍK, J. (Faculty of Electrical Engineering and Information Technology, Slovak University of Technology, Ilkovičova 3, Bratislava, Slovakia); Mr NÁDAŽDY, V. (Institute of Physics, Slovak Academy of Sciences, Dúbravská 1, Bratislava, Slovakia)

Presenter: Dr KOTOROVÁ, S. (Faculty of Electrical Engineering and Information Technology, Slovak University of Technology, Ilkovičova 3, Bratislava, Slovakia)

Session Classification: Poster session

Contribution ID: 143

Type: **not specified**

Surprise for the persistent individuals - goodbye drink

Thursday 10 September 2020 15:35 (20 minutes)