

Searching for Dark Photons at the LHeC & FCC-he

Zeren Simon Wang

1909.02312

in collaboration with M. D'Onofrio and O. Fischer

December 4, 2019

IBS-PNU Joint Workshop on Physics beyond the Standard Model
Busan, Korea



asia pacific center for
theoretical physics

- Dark Matter $\left\{ \begin{array}{l} \text{Galaxy rotation curves} \\ \text{Structure formation} \\ \dots \end{array} \right.$
- Hidden sector theories predict new particles to interact with the Standard Model (SM) field content via feebly coupled mediator particles
- Different portals between the dark sector and the SM: focus on the vector portal and dark photon
- Studied at the HL-LHC, CEPC, FCC-ee/hh, beam-dump, and fixed-target experiments
- Sensitivity of electron-proton colliders may be unique:
LHeC & FCC-he

The dark photon model

- Extend the SM gauge group by an additional (broken) gauge group:
 $U(1)_X$

$$\mathcal{L} \supset -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} - \frac{\epsilon}{2}X_{\mu\nu}F^{\mu\nu}$$

- Applying a field re-definition to get rid of the kinetic mixing term
 $X_{\mu\nu}F^{\mu\nu}$

$$\mathcal{L} \supset -\sum_f \bar{f} \epsilon e q_f A' f$$

- Mass can stem from e.g. a Higgs mechanism
- SM particles assumed *uncharged* under $U(1)_X$
- Two parameters: $\{m_{\gamma'}, \epsilon\}$
- One parameter ϵ controls both production and decay
- Focus on MeV-GeV mass range in this work: lepton and pion pair final states

Portals between the SM and the dark sector

The only non-gravitational portals possible connecting the SM and dark sector:


- scalar: dark Higgs
- pseudoscalar: axion-like particles
- vector: dark photons
- neutrino: heavy neutral leptons

Dark photon could be the messenger to the dark sector or even constitute dark matter, motivating search for dark photons




	E_e (GeV)	E_p (TeV)	\sqrt{s} (TeV)	\mathcal{L}_{int} (ab^{-1})
LHeC	60	7	1.3	1
FCC-he	60	50	3.5	3

- Ideal laboratory to study common features of electrons and quarks
- Advantages:
 - Small background (no QCD interaction between e and p)
 - Very low pileup
 - Less problem of synchrotron radiation that occurs typically in circular lepton colliders
 - ...
- Disadvantages:
 - Relatively smaller scattering cross section
 - ...

Dark photon production




DIS: $e^- \text{ parton} \rightarrow e^- \text{ parton} \gamma'$



DIS: $Q^2 \gg m_p^2 \simeq 1 \text{ GeV}^2$

Signal estimation



$$N_{\text{dv}}(\sqrt{s}, \mathcal{L}, m_X, \epsilon)$$

$$= \sigma(M, \epsilon) \mathcal{L} \times \int D(\vartheta, \gamma) P_{\text{dv}}(x_{\min}(\vartheta), x_{\max}(\vartheta), \Delta x_{\text{lab}}(\tau, \gamma)) d\vartheta d\gamma$$


$$P_{\text{dv}} = \text{Exp}\left(\frac{-x_{\min}}{\Delta x_{\text{lab}}}\right) - \text{Exp}\left(\frac{-x_{\max}}{\Delta x_{\text{lab}}}\right)$$

$$x_{\min} = 200\mu m, x_{\max} = \infty, \Delta x_{\text{lab}} = \tau_{\text{lab}} |\vec{v}| = \beta_{\gamma'} \gamma' \tau c$$

	$\eta(e/\mu)$	$\eta(\text{jets})$
LHeC	(-4.3, 4.9)	(-5, 5)
FCC-he	(-5.0, 5.2)	(-5.5, 5.5)

In practice, apply $|\eta| < 4.7$

Dark photon decays




[Raggi, Kozhuharov, '15]

$$\Gamma(\gamma' \rightarrow l^+l^-) = \frac{1}{3} \alpha_{\text{QED}} m_{\gamma'} \epsilon^2 \sqrt{1 - \frac{4m_l^2}{m_{\gamma'}^2}} \left(1 + \frac{2m_l^2}{m_{\gamma'}^2}\right)$$


$$\Gamma_{\text{total}}(\gamma') = \frac{\Gamma(\gamma' \rightarrow e^-e^+)}{\text{BR}(\gamma' \rightarrow e^-e^+)}$$

- Real low-energy photons interacting with the detector material or the beam pipe $\rightarrow e^-e^+$ pairs
 - location of the secondary vertex coincides with the detector material or the beam pipe
 - easily rejected
- Long-lived hadrons such as K_S , K_L , and Λ
 - lifetime far away from IP (3cm, 15m, and 8cm)
 - hadronic activity is aligned with the proton beam and propagates mostly into the forward hemisphere of the detector
 - their primary decay channels are only marginally consistent with our signal signature
 - their masses are well known


Results I: 90% C.L. sensitivity reaches




Results II: 90% C.L. sensitivity reaches



- Blue solid line: Nbkg=0, signal efficiency = 100%
- Blue dashed line: Nbkg=0, signal efficiency = 20%
- Red solid line: Nbkg=100, signal efficiency = 100%
- Red dashed line: Nbkg=100, signal efficiency = 20%



Results III: Comparison



[Physics Briefing Book: Input for the European Strategy for Particle Physics Update 2020 [1910.11775]]

- DM may reside in a dark sector
- Different portals between the SM and the dark sector
- The vector portal concerns a dark photon which may provide connection to DM or even constitute the DM (kinetic mixing)
- New and existing searches ongoing hunting such dark photons
- Electron-proton colliders may explore a unique territory in the parameter space for MeV-GeV dark photons

Thank You!