

Studies of baryon-baryon interaction at J-PARC

Recent results and future plans at J-PARC on baryon-baryon interaction studies are presented.

We recently observed several new events of double-strange hypernuclei in the emulsion-counter hybrid experiment (E07) [1]. Observed events indicate a rather deep binding energy of Ξ in the ^{14}N nucleus. One of the events is interpreted as a bound state of Ξ in the nuclear 0s orbit, suggesting that the ΞN - $\Lambda\Lambda$ interaction is unexpectedly weak [2]. Further Ξ -hypernuclear studies by missing mass spectroscopy of the (K^- , K^+) reaction (E70) will be conducted soon.

We also successfully carried out a high-statistics Σ^\pm - p scattering experiment (E40) [3]. In the Σ^+p channel, the obtained cross section was converted to the phase shift, showing that the Σ^+p interaction has a much stronger repulsive core than the NN interaction, which is interpreted as a result of Pauli principle in the quark level [4].

In the future, we plan to investigate the ΛNN three-body force via high-resolution Λ -hypernuclear spectroscopy. Together with high quality Λ - p scattering experiments, it allows us to solve the “hyperon puzzle” in neutron stars. Such studies will be performed using new secondary beam lines in the J-PARC Hadron Facility extension project [5].

[1] H. Ekawa et al., Prog. Theor. Exp. Phys. 2019 (2019) 021D02;

S. H. Hayakawa et al., Phys. Rev. Lett. 126 (2021) 062501.

[2] M. Yoshimoto et al., Prog. Theor. Exp. Phys. 2021 (2021) 7.

[3] K. Miwa et al., Phys. Rev. C 104 (2021) 045204; Phys. Rev. Lett. 128 (2022) 072501.

[4] T. Nanamura et al., Prog. Theor. Exp. Phys. 2022 (2022) 093D01,

[5] K. Aoki et al., arXiv: 2110.04462 [nucl-ex] (2021).

Primary author: Prof. TAMURA, Hirokazu (Tohoku University)

Presenter: Prof. TAMURA, Hirokazu (Tohoku University)

Track Classification: Hadron-hadron interactions