



Contribution ID: 120

Type: oral

## Probing the nucleus with linearly polarized photons

Wednesday 13 January 2021 19:25 (20 minutes)

The intense electromagnetic fields produced by ultra-relativistic heavy nuclei have been proposed as a source of quasi-real photons i.e. in the Weizsacker-Williams equivalent photon method. A photon from one nucleus can fluctuate into a quark antiquark pair and interact directly with the other nucleus to produce a vector meson (e.g.  $\rho^0$ ). Recent STAR measurements of the Breit-Wheeler pair production process -  $\gamma\gamma \rightarrow e^+e^-$  have demonstrated that the interacting photons are linearly polarized, and that photon polarization induces angular modulations in the final state particle distribution.

In this talk we present STAR measurements of diffractive photo-production of the  $\rho^0$ -meson (and direct  $\pi^+\pi^-$  pairs) in ultra-peripheral Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. We measure azimuthal angular distributions of the final state  $\pi^+\pi^-$  pairs and observe  $\cos 2\Delta\phi$  and  $\cos 4\Delta\phi$  modulations (where  $\Delta\phi = \Delta\phi[(\pi^+ + \pi^-), (\pi^+ - \pi^-)]$ ). Theoretical predictions suggest that such modulations may provide new insight into nuclear structure and may shed light on the transverse momentum dependent (TMD) distributions of gluons within large nuclei - a topic of great interest both at existing experiments and at a future Electron Ion Collider.

[1] C. Weizsäcker, Zeitschrift für Physik 88 (1934): 612–25.

[2] H. Xing, C. Zhang, J. Zhou, and Y.-J. Zhou, ArXiv:2006.06206 [Hep-Ph] (2020).

[3] A. Metz, and J. Zhou. Physical Review D 84 5 (2011).

**Primary author:** Dr BRANDENBURG, James (Brookhaven National Laboratory)

**Presenter:** Dr BRANDENBURG, James (Brookhaven National Laboratory)

**Session Classification:** PDF

**Track Classification:** Partonic structure of protons and nuclei